HEALTH BASICS

Body temperature norms

Body temperature norms describe the usual range of temperature readings in healthy people.

WHAT IT IS
The usual range of healthy body temperature readings

WHERE IT COMES UP
Home checks, medical assessments, illness, exercise, and hot weather

WHY IT MATTERS
Helps identify concerning changes and measurement errors

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
A pharmacist, nurse, primary care clinician, or urgent care service

HOW IT IS CHECKED
A thermometer used at a specified body site

Overview

Body temperature is a measurement of the balance between heat made and heat lost by your body. A "normal" temperature is not one exact number. The familiar figure of 37°C, or 98.6°F, is a useful reference point, not a rule that applies to every person at every time.

Many healthy readings fall roughly between 36°C and 37.5°C. The expected reading is affected by the body site, thermometer type, time of day, age, activity, hormones, and surrounding temperature. Research reviewing normal temperatures has also found meaningful differences between people and measurement methods.

A fever is usually defined as a temperature of 38°C, or 100.4°F, or higher. However, a reading should be interpreted alongside the person's age, symptoms, measurement site, and overall condition. A person may feel feverish below 38°C, while a single unusual reading can result from technique or equipment.

Body temperature norms are not a disease or diagnosis. They are reference ranges used to understand a thermometer reading.

How it works

A part of the brain called the hypothalamus helps regulate body temperature. It works like a control centre, adjusting heat production and heat loss.

Your muscles and organs produce heat. Your body loses heat through the skin, breathing, sweating, and contact with cooler surroundings. Blood vessels near the skin can widen to release heat or narrow to conserve it. Shivering creates additional heat through muscle activity.

Temperature normally follows a daily rhythm. It is often lower in the early morning and higher in the late afternoon or evening. Exercise, eating, sleep, stress, and menstrual-cycle changes can also shift a reading.

The measurement site matters because thermometers do not all measure the same part of the body:

There is no reliable adjustment that converts every armpit, mouth, ear, or rectal reading into an exact equivalent. Record the site rather than automatically adding or subtracting a fixed number.

Different devices also work differently. A digital contact thermometer must touch the body correctly. An infrared thermometer measures heat coming from the ear or forehead and must be positioned as its instructions describe. Skin-temperature strips are less reliable for deciding whether someone has a fever.

- Rectal readings generally come closest to internal, or "core," temperature
- Oral readings measure temperature inside the mouth
- Armpit readings often run lower and are more affected by positioning
- Ear readings can be affected by placement, earwax, and ear-canal shape
- Forehead readings can be influenced by sweat, distance, and surroundings

Why it matters for your health

Temperature can help you judge whether your body is responding to infection, inflammation, environmental heat, or cold exposure. It can also help a clinician assess how urgently you need care.

The number is only one part of that assessment. How you feel and function may be more important than a small difference in the reading. For example, someone who is alert, drinking normally, and breathing comfortably may need different care from someone with the same

temperature who is confused or struggling to breathe.

Age is important. Babies younger than 3 months with a temperature of 38°C or higher need urgent medical assessment because serious infections can be difficult to recognise at that age.

Some people may not develop a high temperature even with significant illness. This can occur in older adults and in people whose immune system is weakened by a health condition or treatment. Their usual temperature may provide useful context, but a low baseline does not create a separate, fixed fever threshold without clinical advice.

Repeatedly checking your temperature when you feel well is usually unnecessary. A personal record may be useful if a clinician has asked you to monitor it or if you are investigating repeated unusual readings.

What can go wrong

An abnormal reading may reflect a health condition, an environmental problem, or inaccurate measurement.

Fever is a regulated rise in temperature, often caused by infection. Inflammation, some medicines, and certain medical conditions can also cause fever. Fever is not the same as overheating: during fever, the brain temporarily raises the body's temperature target.

Heat exhaustion and heatstroke happen when the body gains more heat than it can lose. Heatstroke is an emergency, especially when overheating is accompanied by confusion, collapse, a seizure, or loss of consciousness. A thermometer reading alone should not delay emergency help when these signs are present.

Hypothermia means a core body temperature below 35°C, or 95°F. It can result from cold exposure, wet clothing, immersion in cold water, or health problems that reduce heat production. Suspected hypothermia needs urgent medical care and may be an emergency.

Measurement error is common. A reading may be misleading if the thermometer is poorly positioned, used at a different site than intended, or affected by direct sunlight, sweat, recent exercise, or a hot or cold drink. A damaged device or low battery may also give inconsistent results.

Seek emergency help for severe difficulty breathing, confusion, a seizure, collapse, blue or grey skin, inability to wake normally, suspected heatstroke, or suspected serious hypothermia.

For a reading that is only mildly unusual, repeat the measurement correctly and consider how the person feels. Contact a healthcare service promptly if abnormal readings continue,

symptoms are worsening, or you are concerned. A baby younger than 3 months with a temperature of 38°C or higher needs urgent assessment rather than routine follow-up.

What you can do

If you need to check a temperature, use a digital thermometer designed for the chosen body site. Follow the device instructions because positioning and timing differ between models.

You can make a result easier to interpret by recording:

If a result seems unexpected, check the device placement and repeat the measurement after the person has settled in a comfortable indoor environment. Use the same site and method when comparing readings over time.

Ask a pharmacist, nurse, or primary care clinician which thermometer and measurement site are suitable for your age and situation. If monitoring has been recommended, ask what result should prompt a call and which service to contact.

If you feel well and have no reason to monitor your temperature, there is usually nothing you need to do. Normal daily shifts do not require treatment.

- The temperature and unit, either Celsius or Fahrenheit
- The body site used
- The date and time
- The thermometer type
- Relevant concerns, such as chills, poor drinking, or unusual drowsiness
- Any recent exercise, bathing, or hot or cold drinks

Common questions

Is 37°C always normal?

It is a common reference value, but it is not a universal personal baseline. A healthy reading can be somewhat lower or higher depending on the person, time, site, and device.

Is 37.5°C a fever?

Not necessarily. It can be within the expected range in some circumstances, especially later in the day or after activity. Fever is commonly defined as 38°C or higher, but the full situation still matters.

Can I compare an armpit reading with an oral reading?

Only with caution. Different sites produce different results, and fixed conversion formulas are not dependable. Compare readings taken at the same site with the same method.

Which unit should I use?

Either Celsius or Fahrenheit is acceptable. Make sure the unit is recorded. A reading of 38°C equals 100.4°F, while 35°C equals 95°F.

Should I know my personal baseline?

Most people do not need one. It may help if a clinician has asked you to monitor temperature or if your usual readings are consistently outside common ranges.

Does feeling hot prove that I have a fever?

No. Warm surroundings, exercise, hormonal changes, anxiety, and skin blood flow can make you feel hot without a measured fever. A thermometer provides more useful information.

Sources

- MedlinePlus Medical Encyclopedia: Body temperature norms , US National Library of Medicine
- Geneva II, Cuzzo B, Fazili T, Javaid W. Normal Body Temperature: A Systematic Review , Open Forum Infectious Diseases , 2019
- NICE guideline NG143: Fever in under 5s-assessment and initial management , National Institute for Health and Care Excellence
- NHS: High temperature in adults , National Health Service
- NHS: High temperature in children , National Health Service
- NHS: Hypothermia , National Health Service
- FDA: Non-contact infrared thermometers , US Food and Drug Administration