HEALTH BASICS

Exercises for COPD

Exercises for COPD are planned activities that build fitness, strength, and confidence with movement.

WHAT IT IS
Planned physical activity adapted for people with COPD

WHERE IT COMES UP
COPD care, pulmonary rehabilitation, and recovery after flare-ups

WHY IT MATTERS
Improved stamina, strength, breathlessness control, and daily function

WHAT IT IS NOT
Not a disease or a replacement for medical treatment

WHO TO ASK
Pulmonary rehabilitation team, respiratory clinician, or physiotherapist

HOW IT IS CHECKED
Walk testing, oxygen levels, symptoms, and exercise response

Overview

Exercises for COPD are activities chosen to improve fitness, muscle strength, balance, and confidence. They are not a disease or a medical diagnosis.

COPD, or chronic obstructive pulmonary disease, is a long-term lung disease that limits airflow. It can make physical activity feel difficult. Some people then move less to avoid breathlessness. Over time, their muscles become less efficient, so everyday tasks require more effort.

Regular activity can help interrupt this cycle. It does not reverse the lung damage caused by COPD, but it can help your heart and muscles use oxygen more efficiently. This may make walking, dressing, shopping, or climbing stairs more manageable.

Pulmonary rehabilitation is often the safest and most useful place to begin. This is a supervised programme combining exercise, education, and support. The team assesses your health and adjusts activities to your abilities, other medical conditions, and oxygen needs.

How it works

Exercise for COPD usually includes several types of activity.

Endurance exercise trains your heart, lungs, and muscles to work together for longer. Examples include walking, using a stationary bicycle, or exercising while seated. Some programmes use interval training, which alternates short periods of activity with rest or easier movement.

Strength exercise works specific muscle groups. It may use resistance bands, light weights, exercise machines, or your own body weight. Stronger leg and arm muscles need less effort for daily tasks.

Balance and flexibility exercises support safer movement and joint mobility. They may be especially relevant if you have fallen, feel unsteady, or have reduced mobility.

Breathing techniques may help you manage breathlessness during activity. Pursed-lip breathing involves breathing in gently and breathing out slowly through narrowed lips. It can help some people slow their breathing and empty their lungs more comfortably. Breathing techniques support exercise but do not replace physical conditioning.

A warm-up allows your breathing and circulation to increase gradually. A cool-down gives them time to settle. Stopping suddenly after harder activity can make some people feel light-headed.

Breathlessness during exercise is not automatically a sign of harm. A pulmonary rehabilitation team may use a breathlessness scale, a walking test, and oxygen measurements to find an appropriate level. Spirometry, the breathing test used to assess airflow, does not by itself show how much exercise you can safely manage.

Why it matters for your health

Exercise training is a central part of pulmonary rehabilitation for COPD. Evidence shows that pulmonary rehabilitation can improve exercise capacity, breathlessness, and health-related quality of life.

The benefits are not limited to the lungs. Activity can strengthen muscles, support balance, and help maintain independence. It may also improve confidence in managing breathlessness.

Pulmonary rehabilitation is particularly important after a hospital admission for a COPD flare-up, also called an exacerbation. A flare-up is a period when COPD-related breathing problems become worse than usual and may require additional treatment. The timing and setting of rehabilitation should be planned with the clinical team involved in your recovery.

Exercise works alongside treatments such as inhalers, smoking cessation support,

vaccinations, and prescribed oxygen. It does not replace them. Likewise, oxygen is not a substitute for activity when exercise is otherwise safe.

Benefits usually require continued activity. A rehabilitation programme can help you find forms of exercise that are realistic at home or in your community after supervised sessions end.

What can go wrong

Exercise plans need adjustment when other health conditions or changes in COPD affect safety.

A COPD exacerbation can reduce exercise tolerance. During a flare-up, your usual session may be too demanding. A clinician or rehabilitation team can advise when and how to restart.

Exercise-related oxygen desaturation means the oxygen level in your blood falls during activity. It may not be obvious from breathlessness alone. A supervised walk test with a pulse oximeter can check for this. If you use prescribed oxygen, do not change the flow rate unless your oxygen team has given you a specific plan.

Heart conditions can also cause breathlessness, chest discomfort, palpitations, or faintness during exercise. Muscle or joint conditions may require lower-impact activities. Poor balance, unsuitable footwear, loose rugs, or oxygen tubing can increase the risk of falls.

Muscle tiredness and some breathlessness can be expected during training. Seek routine clinical advice if your exercise ability is steadily declining, your usual activity produces much more breathlessness, or you repeatedly feel dizzy or notice palpitations.

Stop exercising and seek urgent medical advice if you develop a marked change in breathing that does not settle with rest or with your agreed action plan.

Call emergency services for severe breathing difficulty, fainting, new confusion, blue or grey lips, or chest pain that is severe, persistent, or accompanied by sweating or nausea. Do not try to exercise through these symptoms.

What you can do

Ask whether pulmonary rehabilitation is suitable for you. Programmes may be offered in hospitals, community settings, at home, or remotely. Availability differs by area.

Before starting or changing a programme, useful questions include:

Keep a simple record of the activity, how long you managed, how breathless you felt, and whether you had chest discomfort, dizziness, or unusual fatigue. This can help a rehabilitation

professional adjust your plan.

Choose activities that are practical and repeatable. Walking indoors, using a stationary bicycle, or doing seated exercises may be easier than outdoor activity during very hot, cold, or polluted conditions.

Start at the level set for you and build gradually. Do not copy another person's exercise target, oxygen setting, or resistance level. COPD severity, fitness, other medical conditions, and recent flare-ups all affect what is appropriate.

If you have an action plan for COPD flare-ups, keep it available. An exercise plan should also state when to reduce activity, when to stop, and who to contact.

- Do I need a supervised walking or exercise test?
- Should my oxygen level be checked during activity?
- How should I use my prescribed inhalers or oxygen around exercise?
- Which activities fit my joint, heart, balance, or mobility needs?
- What signs mean I should pause and contact the clinical team?

Common questions

Can exercise improve lung function?

Exercise does not repair the structural lung damage caused by COPD. Its main effects are on muscle efficiency, cardiovascular fitness, confidence, and the effort required for daily activities.

Is walking enough?

Walking can provide endurance training, especially when its pace or duration is planned. Strength and balance work may add benefits that walking alone does not provide. A pulmonary rehabilitation assessment can identify the most useful combination.

Should I avoid exercise if I become breathless?

Some breathlessness is expected with physical activity. The aim is to exercise at a manageable level, not to avoid movement completely or push through severe distress. A breathlessness scale and supervised testing can help define that level.

Can I exercise while using oxygen?

Many people exercise with prescribed oxygen. The equipment and flow setting need to follow your oxygen prescription and safety instructions. Never increase the flow on your own to exercise harder.

Are breathing exercises the same as pulmonary rehabilitation?

No. Breathing techniques may help control breathing during activity, but pulmonary rehabilitation also includes physical training, education, and an individual assessment.

What if I cannot stand or walk far?

Seated cycling, supported movements, and chair-based strength exercises may be possible. A physiotherapist or rehabilitation team can adapt activities around mobility limits rather than using a standard routine.

Sources

- American Thoracic Society: Pulmonary Rehabilitation for Adults with Chronic Respiratory Disease-Official Clinical Practice Guideline , American Journal of Respiratory and Critical Care Medicine , 2023
- National Institute for Health and Care Excellence: Chronic obstructive pulmonary disease in over 16s-diagnosis and management, NG115 , recommendations on pulmonary rehabilitation and COPD care
- National Heart, Lung, and Blood Institute: Pulmonary Rehabilitation , overview of programme assessment, exercise training, and education
- McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y: Pulmonary rehabilitation for chronic obstructive pulmonary disease , Cochrane Database of Systematic Reviews , 2015
- Spruit MA and colleagues: An Official American Thoracic Society/European Respiratory Society Statement-Key Concepts and Advances in Pulmonary Rehabilitation , American Journal of Respiratory and Critical Care Medicine , 2013