HEALTH BASICS

Intrauterine

Intrauterine means located, placed, or happening inside the uterus.

WHAT IT IS
Located, placed, or happening inside the uterus

WHERE IT COMES UP
Pregnancy scans, contraceptive devices, fertility procedures, and medical records

WHY IT MATTERS
Clarifies pregnancy location, device position, or treatment route

WHAT IT IS NOT
Not a disease or diagnosis by itself

WHO TO ASK
Obstetrician-gynaecologist, midwife, fertility clinician, or sexual health service

HOW IT IS CHECKED
Usually ultrasound when location or position needs confirmation

Overview

Intrauterine means "within the uterus." The uterus, also called the womb, is the muscular organ in the pelvis where a pregnancy normally develops.

Intrauterine is an adjective, not a disease or a diagnosis on its own. Its meaning becomes clear only when you read the word that follows it. For example:

The term may appear in an ultrasound report, pregnancy record, contraception appointment, fertility plan, or operation note. Seeing it does not automatically mean that something is wrong.

- An intrauterine pregnancy is located inside the uterus
- An intrauterine device is a contraceptive placed inside the uterus
- Intrauterine insemination places prepared sperm directly into the uterus
- An intrauterine procedure takes place within the uterine cavity

How it works

The uterus has a central space called the uterine cavity. "Intrauterine" usually describes something located in, placed in, or directed into this space.

During pregnancy, a fertilised egg normally attaches to the lining of the uterus. An ultrasound can help determine where the pregnancy is developing. Early in pregnancy, a scan may show a pregnancy sac, yolk sac, embryo, or heartbeat, depending on timing. A very early scan may not yet show enough to confirm the location.

A blood test for human chorionic gonadotropin, or hCG , can show that pregnancy tissue is present. However, an hCG result by itself cannot prove that a pregnancy is intrauterine. Clinicians may use repeat blood tests together with ultrasound when the location remains uncertain.

An intrauterine contraceptive is placed through the cervix, which is the opening of the uterus. Copper intrauterine devices prevent pregnancy without hormones. Hormonal devices release a progestogen inside the uterus. Terminology differs by country: "IUD" may describe all intrauterine contraception, or it may refer specifically to a copper device while "IUS" means a hormonal system.

In intrauterine insemination, or IUI , prepared sperm is passed through the cervix into the uterus around the time of ovulation. The term describes where the sperm is placed. It does not show whether fertilisation or pregnancy will occur, or where a later pregnancy will implant.

Why it matters for your health

Location can change what follow-up or treatment is appropriate.

In early pregnancy, confirming an intrauterine location helps distinguish a pregnancy in the uterus from an ectopic pregnancy . An ectopic pregnancy develops outside the main uterine cavity, most often in a fallopian tube. It cannot be moved into the uterus and needs medical assessment.

An ultrasound report that confirms a definite intrauterine pregnancy is generally reassuring about location. It does not answer every other question about the pregnancy. The report may separately describe development, heartbeat, bleeding, the placenta, or other findings.

For contraception, an intrauterine device needs to sit in an appropriate position to work as intended. Ultrasound is not routinely required for everyone after insertion, but it may be used when there is pain, unusual bleeding, missing threads, concern about expulsion, or uncertainty about position.

In fertility care, "intrauterine" identifies the route of a procedure. It does not describe the cause of fertility problems or guarantee a particular result.

The word also appears in older terms. For example, intrauterine growth restriction has often been used for a fetus that is not growing as expected. Many current medical sources use fetal

growth restriction instead. That is a separate clinical concern, not a meaning of "intrauterine" by itself.

What can go wrong

The word itself does not cause health problems. However, it may appear when clinicians are checking for particular conditions or situations.

A pregnancy of unknown location means a pregnancy test is positive, but ultrasound has not yet shown where the pregnancy is. This may happen because it is too early to see the pregnancy, because a pregnancy has ended, or because it is ectopic. Follow-up commonly involves another ultrasound, repeat hCG tests, or both.

An ectopic pregnancy is outside the main uterine cavity. Get urgent medical advice if you may be pregnant and develop one-sided or worsening pelvic pain, vaginal bleeding, or new pain at the tip of your shoulder.

Call emergency services or go to an emergency department for severe abdominal or pelvic pain, fainting, marked dizziness, collapse, or heavy bleeding, especially if pregnancy is possible. These symptoms can have several causes, but they need prompt assessment.

An intrauterine contraceptive can sometimes move downward, come out, become embedded in the uterine wall, or pass through the wall during insertion. Contact the service that fitted it if you cannot feel threads you were previously able to feel, can feel the hard device itself, or think it has moved. Use the backup contraception advised by that service until its position and pregnancy protection are clarified.

Seek prompt care if you have a positive pregnancy test while an intrauterine contraceptive is in place. These devices greatly reduce the overall chance of pregnancy, including ectopic pregnancy. However, when pregnancy does occur with a device in place, its location needs to be checked.

Fever, worsening pelvic pain, unusual discharge, or feeling significantly unwell after insertion also calls for timely medical advice. Sudden severe pain, fainting, or heavy bleeding requires emergency assessment.

Intrauterine adhesions are bands of scar tissue inside the uterus. They may develop after some uterine procedures or infections and can affect menstrual bleeding or fertility. This is a named condition involving the uterine cavity; it is not another meaning of the word intrauterine.

What you can do

If you see "intrauterine" in a report, read the full phrase rather than interpreting the word alone. There is no test or action needed simply because this term appears.

For an early pregnancy report, you can ask:

Keep a record of pregnancy test dates, the first day of your last menstrual period if known, bleeding, and pain. These details can help the healthcare team interpret early findings. Do not use a single hCG number to decide the location or health of a pregnancy yourself.

If you use an intrauterine contraceptive, keep a note of the device type and insertion date. Ask how long that specific product is intended to remain in place and when it should be reviewed, replaced, or removed. If you were taught to check the threads, ask what to do if they feel different or cannot be found.

After IUI, follow the fertility service's instructions about pregnancy testing and follow-up. The procedure's name does not confirm that implantation occurred inside the uterus.

Routine questions can usually wait for the clinician or service involved in your pregnancy, contraception, scan, or fertility care. New significant pain, bleeding, faintness, fever, or a positive pregnancy test with an intrauterine device calls for more prompt assessment.

- Was the pregnancy definitely seen inside the uterus?
- What structures were visible on the scan?
- Is another scan or blood test needed?
- When should follow-up happen?

Common questions

Does "intrauterine" mean the finding is normal?

Not necessarily. It only describes location, placement, or route. The rest of the report explains whether the finding is expected and whether follow-up is needed.

Does an intrauterine pregnancy rule out ectopic pregnancy?

A clearly confirmed intrauterine pregnancy generally rules out a single ectopic pregnancy. Rarely, pregnancies can exist both inside and outside the uterus, called a heterotopic pregnancy. This is more relevant after some fertility treatments. Ongoing severe or worsening symptoms still need assessment.

Is an intrauterine device the same as an IUD?

Usually. "IUD" means intrauterine device. In some healthcare systems, copper contraception is called an IUD and a hormonal device is called an intrauterine system, or IUS.

Does IUI make sure a pregnancy implants in the uterus?

No. IUI describes placing sperm inside the uterus. It does not guarantee fertilisation, implantation, or an intrauterine pregnancy.

Can a pregnancy be too early to call intrauterine?

Yes. A pregnancy may not yet be visible on ultrasound. Follow-up timing is based on the scan findings, symptoms, pregnancy dates, and sometimes repeat hCG measurements.

Sources

- National Cancer Institute. "Definition of intrauterine." NCI Dictionary of Cancer Terms
- American College of Obstetricians and Gynecologists. "Ectopic Pregnancy."
- American College of Obstetricians and Gynecologists. "Long-Acting Reversible Contraception: Intrauterine Device and Implant."
- Centers for Disease Control and Prevention. U.S. Selected Practice Recommendations for Contraceptive Use, 2024 . Morbidity and Mortality Weekly Report
- National Institute for Health and Care Excellence. Ectopic pregnancy and miscarriage: diagnosis and initial management . Guideline NG126
- NHS. "Intrauterine insemination (IUI)."