HEALTH BASICS

Intrinsic factor

Intrinsic factor is a stomach protein that helps your body absorb vitamin B12.

WHAT IT IS
A stomach protein needed for normal vitamin B12 absorption

WHERE IT COMES UP
Vitamin B12 deficiency and pernicious anemia testing

WHY IT MATTERS
Enables the gut to absorb enough vitamin B12

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Primary care clinician or gastroenterology team

HOW IT IS CHECKED
Intrinsic factor antibody blood test when pernicious anemia is suspected

Overview

Intrinsic factor is a protein made by specialized cells in the lining of your stomach. Its main job is to help your small intestine absorb vitamin B12.

Vitamin B12 supports red blood cell production, nerve function, and DNA formation. Most vitamin B12 from food cannot enter the body efficiently without intrinsic factor.

Intrinsic factor is not a disease, diagnosis, vitamin, or digestive enzyme. You may hear about it when blood tests show possible vitamin B12 deficiency or when a clinician is investigating pernicious anemia. Pernicious anemia is an autoimmune condition in which the immune system interferes with the stomach cells or intrinsic factor needed for vitamin B12 absorption.

A low vitamin B12 level does not automatically mean that intrinsic factor is missing. Diet, medicines, stomach or intestinal conditions, and previous surgery can also affect vitamin B12 levels or absorption.

How it works

Vitamin B12 in food is usually attached to protein. Stomach acid and digestive enzymes help

release it. The free vitamin first attaches to another transport protein, sometimes called haptocorrin or R-protein.

At the same time, stomach cells called parietal cells release intrinsic factor. These are also the cells that produce stomach acid.

In the first part of the small intestine, pancreatic enzymes separate vitamin B12 from its initial transport protein. The vitamin can then bind to intrinsic factor. This protected vitamin B12-intrinsic factor complex travels through the small intestine.

Near the end of the small intestine, in an area called the terminal ileum , receptors recognize the complex and bring the vitamin into the body. Intrinsic factor is broken down, while vitamin B12 enters the bloodstream attached to transport proteins.

The body stores vitamin B12, mainly in the liver. Because these stores may last for a long time, reduced absorption may not become apparent soon after a stomach or intestinal problem begins.

A small amount of vitamin B12 can be absorbed without intrinsic factor. This is one reason clinician-directed oral vitamin B12 treatment may work for some people who cannot use intrinsic factor normally. Others may need injections. The suitable route depends on the cause and severity of deficiency.

Why it matters for your health

Without enough working intrinsic factor, the body may not absorb enough vitamin B12 from food. Over time, this can contribute to vitamin B12 deficiency.

Deficiency can interfere with red blood cell production and cause megaloblastic anemia , in which red blood cells become unusually large and do not develop normally. It can also affect nerves and thinking. Nerve-related problems may occur even when a blood count does not show anemia.

Intrinsic factor matters when a clinician is trying to understand why vitamin B12 is low. The cause can affect which tests are useful, whether treatment is taken by mouth or injection, and how long follow-up may be needed.

Testing also helps distinguish an absorption problem from low dietary intake. Simply eating more vitamin B12 may not correct a deficiency if the stomach cannot produce working intrinsic factor or the terminal ileum cannot absorb the vitamin-intrinsic factor complex.

What can go wrong

Several conditions and situations can interfere with this absorption pathway:

Other causes of low vitamin B12 do not necessarily involve intrinsic factor. These include limited dietary intake, some medicines, and other digestive conditions.

An intrinsic factor antibody blood test can support a diagnosis of autoimmune vitamin B12 deficiency. A positive result is meaningful in the right clinical setting, but a negative result does not rule it out. Not everyone with pernicious anemia has a detectable antibody.

- Autoimmune gastritis occurs when the immune system attacks the stomach lining, including parietal cells. This may reduce both stomach acid and intrinsic factor.
- Pernicious anemia is vitamin B12 deficiency caused by autoimmune loss or blockage of intrinsic factor. Despite its name, nerve effects can occur before anemia is found.
- Intrinsic factor antibodies may attach to intrinsic factor and prevent vitamin B12 from binding to it or being absorbed.
- Stomach surgery , including removal of part or all of the stomach or some forms of bariatric surgery, can reduce the cells that produce intrinsic factor.
- Terminal ileum disease or removal can prevent absorption even when the stomach makes enough intrinsic factor. Examples include Crohn's disease affecting the ileum and surgery that removes this section.
- Congenital intrinsic factor deficiency is an uncommon inherited condition in which the body does not make functional intrinsic factor.
- Imerslund-Gräsbeck syndrome is an inherited disorder affecting the intestinal receptor that normally absorbs the vitamin B12-intrinsic factor complex.

What you can do

There is no reason to test intrinsic factor routinely if you feel well and have no relevant blood-test findings or risk factors.

If vitamin B12 deficiency is suspected or confirmed, ask what may be causing it. Useful questions include:

A methylmalonic acid test can provide supporting evidence because this substance may rise when cells do not have enough usable vitamin B12. Kidney problems can also raise it, so the result must be interpreted in context. A full blood count can check for anemia and unusually large red blood cells, but a normal result does not exclude vitamin B12 deficiency.

Keep a record of previous stomach or bowel surgery, digestive diagnoses, medicines,

supplements, and earlier vitamin B12 results. Tell the clinician if close biological relatives have pernicious anemia or autoimmune conditions.

Arrange prompt medical assessment for new numbness, weakness, balance problems, difficulty walking, confusion, or changes in vision. These symptoms have many possible causes and should not be assumed to come from vitamin B12.

Seek emergency care for severe breathing difficulty, chest pain, fainting, sudden one-sided weakness, or sudden trouble speaking. These are emergency warning signs regardless of whether vitamin B12 is involved.

- Was the result based on a vitamin B12 level alone?
- Do I need a full blood count or methylmalonic acid test?
- Could an intrinsic factor antibody test clarify the cause?
- Could stomach surgery, bowel disease, diet, or medicines be relevant?
- Will I need repeat blood tests or long-term vitamin B12 replacement?

Common questions

Is intrinsic factor found in food or supplements?

No. Your stomach makes intrinsic factor. Foods and supplements may contain vitamin B12, but they do not replace intrinsic factor itself.

Does a low vitamin B12 level prove that I lack intrinsic factor?

No. Low vitamin B12 has several possible causes. Test results, medical history, diet, medicines, and sometimes antibody testing help identify the likely explanation.

Can intrinsic factor itself be measured?

Direct measurement is not part of routine assessment. Clinicians usually check vitamin B12 status and may order an intrinsic factor antibody test when autoimmune absorption problems are suspected.

Can supplements restore intrinsic factor?

Vitamin B12 treatment does not make the stomach produce intrinsic factor again. Instead, treatment supplies vitamin B12 in a way that can bypass or overcome the absorption problem. The appropriate form and follow-up should be based on the cause.

Can there be nerve problems without anemia?

Yes. Vitamin B12 deficiency can affect nerves even when hemoglobin and red blood cell measurements are not clearly abnormal. New neurological changes deserve prompt

assessment.

Is pernicious anemia the same as any vitamin B12 deficiency?

No. Pernicious anemia refers specifically to autoimmune interference with intrinsic factor and vitamin B12 absorption. Vitamin B12 deficiency from diet, medicines, or other intestinal conditions is not automatically pernicious anemia.

Sources

- National Institutes of Health Office of Dietary Supplements: Vitamin B12 Fact Sheet for Health Professionals
- NICE guideline NG239: Vitamin B12 deficiency in over 16s-diagnosis and management , 2024
- Nielsen MJ, Rasmussen MR, Andersen CBF, Nexø E, Moestrup SK. "Vitamin B12 transport from food to the body's cells-a sophisticated, multistep pathway." Nature Reviews Gastroenterology & Hepatology . 2012;9:345-354
- Green R, Allen LH, Bjørke-Monsen AL, et al. "Vitamin B12 deficiency." Nature Reviews Disease Primers . 2017;3:17040
- Shah SC, Piazuelo MB, Kuipers EJ, Li D. "AGA Clinical Practice Update on the Diagnosis and Management of Atrophic Gastritis: Expert Review." Gastroenterology . 2021;161(4):1325-1332.e7
- Mayo Clinic Laboratories: Intrinsic Factor Blocking Antibody, Serum