HEALTH BASICS

Overcoming breastfeeding problems

Practical ways to assess and manage common problems with breastfeeding, milk expression, and milk transfer.

WHAT IT IS
Practical support for common breastfeeding challenges

WHERE IT COMES UP
Feeding, pumping, recovery after birth, and infant weight checks

WHY IT MATTERS
Supports milk transfer, comfort, growth, and feeding choices

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Lactation specialist, midwife, pediatrician, or obstetric clinician

HOW IT IS CHECKED
Observed feeding, infant weight, diaper output, and breast assessment

Overview

Breastfeeding problems are difficulties with feeding at the breast, expressing milk, milk supply, or comfort. They are not a single disease or diagnosis. Different problems have different causes, and more than one may happen at the same time.

Common concerns include painful attachment, sore nipples, breast fullness, worries about milk supply, difficulty pumping, or a baby who struggles to take in milk. Some improve with small changes. Others need an assessment of the parent, the baby, and a complete feed.

Breastfeeding does not have to be all or nothing. Direct breastfeeding, expressed milk, donor milk where available, infant formula, or a combination may be appropriate. Your health, your baby's health, and your feeding goals all matter.

Early support can be useful, especially when feeding is painful or you are unsure whether your baby is getting enough milk. Asking for help does not mean you have failed.

How it works

Milk production works mainly through milk removal. When a baby feeds effectively, or milk is expressed, the breast receives signals to keep producing milk. Frequent, effective feeding is especially important while milk production is becoming established.

Milk supply cannot be judged reliably by breast fullness, leaking, pumping volume, or how long a feed lasts. Breasts often feel softer as feeding becomes established. A pump may also remove less milk than a baby can, particularly if the flange does not fit or the settings are uncomfortable.

Attachment, also called the latch, describes how the baby takes the nipple and surrounding breast tissue into the mouth. Effective attachment helps the baby remove milk and reduces pressure on the nipple. Some initial pulling may be noticeable, but pain that continues through a feed deserves attention.

Babies do not all feed on a fixed schedule. Feeding cues include bringing hands to the mouth, turning toward touch, and becoming restless. Crying is a later cue and can make attachment harder. Cluster feeding-several feeds close together-can be normal, particularly during the early weeks.

A clinician assesses milk transfer using the whole picture. This may include watching a feed, listening for swallowing, checking the baby's weight pattern and diaper output, and examining the baby's mouth and the parent's breasts or nipples.

Why it matters for your health

Persistent nipple or breast pain can make feeding difficult and may lead to damaged skin. Cracks and bleeding are often linked to shallow attachment, pump friction, skin irritation, or other causes that can be assessed and treated.

Breasts can become uncomfortably full when milk production increases or milk is not removed as expected. This is called engorgement. Severe swelling may flatten the area around the nipple, making attachment harder.

Breast inflammation can cause a tender, swollen area. Mastitis is inflammation of breast tissue and may include fever or flu-like feelings. Some cases involve bacterial infection and need antibiotics like amoxicillin or doxycycline . Continuing to breastfeed is generally safe, but forceful massage and repeated pumping to "empty" the breast can worsen swelling.

For the baby, effective milk transfer supports hydration and growth. A weight check is more useful than guessing from feed length or pumping output. Some babies need temporary extra milk while the cause of poor transfer is addressed.

Feeding difficulties can also affect sleep, mood, and confidence. Tell a healthcare professional

if distress, anxiety, or low mood is persistent or makes daily life hard. Urgent mental health support is needed if you may harm yourself or your baby.

What can go wrong

Painful or damaged nipples may result from shallow attachment, an awkward feeding position, a poorly fitting pump flange, excessive suction, dermatitis, or blood-vessel spasm. Burning pain is sometimes labelled as thrush, but pain alone cannot confirm a yeast infection.

Engorgement causes firm, swollen, painful breasts, often when milk production increases. Cold packs between feeds and gentle handling may reduce discomfort. Avoid deep massage or forceful squeezing.

Inflammatory mastitis can cause a red or discoloured, hot, painful area with fever or body aches. Symptoms that are severe, worsening, or not improving need medical assessment. An abscess is a pocket of infected fluid that may require imaging and drainage.

Low milk production or poor milk transfer can relate to infrequent milk removal, ineffective attachment, illness, some medicines, hormonal conditions, previous breast surgery, or a baby's feeding difficulty. Perceived low supply is not always true low supply, so assessment should come before assuming that more milk is needed.

Oversupply or a forceful milk flow may cause leaking, breast discomfort, coughing or pulling away during feeds, and repeated fullness. Extra pumping can maintain oversupply, so changes are best planned with skilled support.

Tongue-tie is a tight band of tissue under the tongue. It does not always cause feeding problems. Appearance alone does not show whether treatment is needed; function, milk transfer, pain, and growth should be assessed.

Seek same-day medical advice if:

Call emergency services if your baby is difficult to wake, has trouble breathing, appears blue or grey, has a seizure, or is not responding normally.

- Your baby is unusually sleepy, feeds poorly, or has clearly reduced wet diapers
- Jaundice-the yellowing of skin or eyes-is increasing
- Your baby is not gaining weight as expected
- You have fever with a painful, red or discoloured breast area
- Breast pain, swelling, or nipple damage is worsening
- A breast lump persists or becomes soft and fluid-filled

What you can do

Ask for an observed feed. A lactation specialist, midwife, health visitor, pediatric clinician, or maternity clinician can assess positioning, attachment, swallowing, comfort, and milk transfer.

Keep a short feeding record if there is concern about intake. Note feeds, expressed milk or formula offered, wet and dirty diapers, and any weight measurements. Do not repeatedly weigh your baby at home unless a clinician has explained how and when to do it.

During feeding, bring your baby close with the head and body aligned. Wait for a wide-open mouth before bringing the baby toward the breast. If pain continues, gently break the suction with a clean finger and try again rather than enduring the full feed.

For pumping, check that the flange fits without rubbing or pulling in excessive breast tissue. Use comfortable suction; stronger suction does not necessarily remove more milk. If pumping is being used to protect supply, ask how often to express and when the plan should be reviewed.

If supplementation is recommended, ask:

For mild fullness, feed according to your baby's cues and express only enough for comfort if needed. Cold packs may help swelling. Ask a pharmacist or clinician which pain medicine is safe for you, particularly if you have other conditions or take medicines.

Arrange routine follow-up when feeding is improving but you still need help with comfort, pumping, combination feeding, or your longer-term feeding plan. There is nothing you must do to continue breastfeeding at any cost. A safe feeding plan that protects both parent and baby is the goal.

- Why extra milk is needed
- What type and amount are appropriate
- How your baby's weight and hydration will be followed
- Whether expressing could help maintain milk production
- When the feeding plan should be reassessed

Common questions

Does nipple pain mean the latch is wrong?

Not always. Attachment is a common cause, but pump injury, skin conditions, infection, and blood-vessel spasm can also cause pain. An observed feed and nipple examination can help

identify the cause.

Does a small pumping volume mean I have low supply?

No. Pump output is affected by timing, equipment fit, pump type, stress, and how your body responds to the pump. Your baby's weight pattern, diaper output, and observed milk transfer are more informative.

Can I breastfeed with mastitis?

Breastfeeding is generally safe and may continue according to comfort. Avoid aggressive massage or excessive pumping. Seek medical care for fever, worsening symptoms, or lack of improvement because antibiotics or an ultrasound may be needed.

Will giving formula end breastfeeding?

Not necessarily. Some families use temporary or ongoing combination feeding. If maintaining milk production is important to you, ask whether expressing is needed when a breastfeed is replaced.

Should tongue-tie always be divided?

No. Many tongue-ties do not impair feeding. Treatment decisions should follow a skilled assessment of tongue function, attachment, pain, milk transfer, and growth.

Sources

- American College of Obstetricians and Gynecologists. Breastfeeding Challenges , Committee Opinion No. 820 , 2021
- Academy of Breastfeeding Medicine. Clinical Protocol #36: The Mastitis Spectrum, Revised 2022 , Breastfeeding Medicine , 2022
- Academy of Breastfeeding Medicine. Clinical Protocol #3: Supplementary Feedings in the Healthy Term Breastfed Neonate, Revised 2017 , Breastfeeding Medicine , 2017
- Centers for Disease Control and Prevention. Low Milk Supply , Breastfeeding Special Circumstances
- NHS. Breast pain and breastfeeding
- NHS. Sore or cracked nipples when breastfeeding