HEALTH BASICS

Sex-linked dominant

Sex-linked dominant describes how some gene variants on the X chromosome can pass through families.

WHAT IT IS
A pattern of inheriting a gene variant on the X chromosome

WHERE IT COMES UP
Family histories, genetic diagnoses, and reproductive planning

WHY IT MATTERS
Changes inheritance chances and who may be affected

WHAT IT IS NOT
Not a disease or diagnosis

WHO TO ASK
Genetic counselor or clinical genetics service

HOW IT IS CHECKED
Family pedigree and targeted genetic testing

Overview

Sex-linked dominant is an inheritance pattern. It describes how a gene variant may pass from parent to child when the gene is on a sex chromosome.

The term usually means X-linked dominant . The X and Y chromosomes are called sex chromosomes. Most people have two X chromosomes or one X and one Y, although other chromosome patterns occur.

"Dominant" means that one altered copy of a gene may be enough to affect a person. This does not mean that the related condition is always severe. It also does not mean that everyone with the variant will be affected in the same way.

Sex-linked dominant is not a disease or a diagnosis. A genetic report should name the gene, the specific variant, and the condition it may be associated with. The inheritance label alone cannot tell you what medical care is needed.

How it works

Genes are sections of DNA that provide instructions for the body. Each chromosome contains many genes. People usually inherit one copy of each chromosome from each biological parent.

A person with two X chromosomes usually has two copies of most X-linked genes. If one X carries a dominant disease-causing variant, that person may develop features of the related condition.

A person with one X and one Y chromosome usually has only one copy of most X-linked genes. A disease-causing variant on that X may therefore affect them. Some X-linked conditions are more severe in people with one X, but this is not true for every condition.

The usual inheritance calculations are:

Each pregnancy is a separate event. Having one child who inherits the variant does not change the chance for the next pregnancy.

These calculations assume a straightforward X-linked dominant pattern. The real pattern may look different because of a new variant, chromosome differences, egg or sperm mosaicism, or the way a particular gene works.

X-inactivation also matters. In people with two X chromosomes, each cell usually switches off much of one X. If cells switch off the two X chromosomes in an uneven pattern, people with the same variant can have different features or levels of severity.

Not everyone with a disease-causing variant necessarily develops noticeable features. This is called reduced penetrance . When people do have features but they differ in type or severity, this is called variable expression .

Y-linked inheritance is different. Because people usually have only one Y chromosome, the words dominant and recessive are generally not useful for Y-linked genes.

- A parent with two X chromosomes, one carrying the variant, has a 1 in 2 chance of passing that variant in each pregnancy
- A parent with one X and one Y who carries the variant on the X passes it to every child who inherits that X, typically daughters
- That parent does not pass the X-linked variant to children who inherit the Y, typically sons

Why it matters for your health

An inheritance pattern can help explain who in a family may have inherited a particular variant. It may also guide decisions about genetic testing, health monitoring, and pregnancy planning.

The pattern does not determine your health by itself. Medical meaning comes from the specific gene and variant involved. A variant in one X-linked gene may require regular blood

tests or specialist reviews, while a variant in another may call for different follow-up.

A confirmed result can sometimes allow relatives to have targeted testing . This looks specifically for the family's known variant rather than examining many genes without a clear target.

The result may also affect reproductive options. Depending on the situation, available tests can include testing during pregnancy or preimplantation genetic testing with in vitro fertilisation. These options have practical limits and medical risks, so discussing them does not commit you to using them.

What can go wrong

The phrase can be misunderstood as a diagnosis. It cannot identify a condition or predict its severity without the gene, variant, medical history, and family history.

Conditions commonly described as X-linked dominant include:

These are separate conditions with different health effects. Sharing an inheritance pattern does not make them medically similar.

A family may appear not to follow the expected pattern. A variant can arise for the first time in an egg, sperm, or early embryo; this is called a de novo variant . A parent may also have a variant in only some egg or sperm cells, known as germline mosaicism , even when standard blood testing does not find it.

Genetic test results can also be uncertain. A variant of uncertain significance , often shortened to VUS, means there is not enough evidence to decide whether the change affects health. It should not be treated as the same as a confirmed disease-causing variant.

Older records may use "dominant" or "recessive" labels differently from current reports. Some X-linked conditions do not fit neatly into either category because people with one or two X chromosomes can both be affected in different ways.

- X-linked hypophosphatemia , usually related to variants in the PHEX gene
- Incontinentia pigmenti , usually related to variants affecting IKBKG
- MECP2-related disorders , including Rett syndrome in many affected people

What you can do

If this phrase appears in your records, obtain a copy of the full genetic report. Record the gene

name, exact variant, laboratory classification, test date, and laboratory name. The word "positive" alone is not enough to interpret a genetic result.

Useful questions for a genetics professional include:

A family health history can also help. Include biological parents, siblings, children, grandparents, aunts, uncles, and cousins when that information is available. Note confirmed diagnoses and genetic results rather than relying only on descriptions such as "similar problems."

If pregnancy planning is relevant, a genetic counselor or clinical genetics service can explain the chance of passing on the specific variant and which testing options exist. The inheritance label by itself is not enough to select a test.

There is usually nothing to treat or monitor simply because the words "sex-linked dominant" appear in a record. Follow-up is based on the associated condition and the strength of the genetic evidence.

This inheritance term does not create an emergency. A newly found result can usually be discussed through routine follow-up. If a known associated condition has its own urgent warning signs, follow the emergency advice provided for that condition. Time-sensitive pregnancy decisions may justify a prompt genetics appointment, but they are not usually a medical emergency.

- Is this variant classified as pathogenic, likely pathogenic, or uncertain?
- What condition is this variant associated with?
- Does the condition consistently follow an X-linked dominant pattern?
- Could relatives benefit from targeted testing?
- Does the result change routine screening or follow-up?
- Could the classification change as new evidence becomes available?

Common questions

Is sex-linked dominant the same as X-linked dominant?

It usually is. "Sex-linked" is broader language, but dominant inheritance most often refers to a gene on the X chromosome. Y-linked traits are not usually called dominant or recessive.

Can a father pass an X-linked dominant variant to a son?

Not through the usual chromosome pattern. A child who inherits the father's Y chromosome does not receive his X chromosome. Biological relationships, sex chromosome differences, or

an incorrect original result can complicate a family history.

Can someone carry the variant without being affected?

Possibly. Some variants have reduced penetrance, meaning not everyone who inherits them develops clear features. Others cause mild features that were never recognised. The answer depends on the specific gene and variant.

Does dominant mean more serious?

No. "Dominant" describes inheritance, not severity. Some dominant conditions are mild, while others can have major health effects.

Can a family have no previous history?

Yes. The variant may be new in that person, relatives may have mild or unrecognised features, or family information may be incomplete.

Can ancestry testing confirm this pattern?

Consumer ancestry tests are not a substitute for clinical genetic testing. If a result could affect health care or family planning, confirmation through an appropriate clinical laboratory may be needed.

Sources

- MedlinePlus Genetics: What are the different ways a genetic condition can be inherited?
- MedlinePlus Genetics: What do the results of genetic tests mean?
- GeneReviews: X-Linked Hypophosphatemia
- GeneReviews: Incontinentia Pigmenti
- GeneReviews: MECP2 Disorders
- Richards and colleagues: Standards and guidelines for interpreting sequence variants, Genetics in Medicine , 2015