HEALTH BASICS

# Sex-linked recessive

Sex-linked recessive describes how some genetic variants on the X chromosome are inherited.

**WHAT IT IS**
An inheritance pattern involving a recessive X-chromosome variant

**WHERE IT COMES UP**
Genetic reports, family histories, carrier screening, and genetic counselling

**WHY IT MATTERS**
Helps estimate who may have or pass on a condition

**WHAT IT IS NOT**
Not a disease or diagnosis

**WHO TO ASK**
A genetic counsellor or clinical genetics service

**HOW IT IS CHECKED**
Family history review and condition-specific genetic testing

## Overview

Sex-linked recessive is a pattern of inheritance. It describes how certain genetic conditions can pass through families. It is not a disease or a diagnosis.

In most medical settings, "sex-linked recessive" means X-linked recessive . The relevant genetic change, called a variant, is on the X chromosome. The X and Y chromosomes are often called sex chromosomes because they are involved in typical patterns of sexual development.

A person with one X chromosome has only one copy of most genes on that chromosome. If the copy contains a disease-causing variant, there may be no second working copy to make up for it.

A person with two X chromosomes usually has two copies of each X-linked gene. If only one copy contains a recessive variant, the other copy may provide enough gene function. This person is often called a carrier . However, some carriers can have health effects, depending on the gene and how their X chromosomes are used by their cells.

Chromosome patterns do not always fit XX or XY. This means inheritance estimates may need to be adjusted for people with chromosome differences. Gender identity does not determine which chromosomes or genetic variants someone has.

## How it works

Genes are sections of DNA that provide instructions for the body. A pathogenic variant is a genetic change known to interfere with a gene's function. "Recessive" means that a working copy of the gene can often compensate for the changed copy.

People usually inherit one X chromosome from the egg and either an X or Y chromosome from the sperm. A parent cannot pass an X chromosome to a child who receives that parent's Y chromosome.

For a person with two X chromosomes who carries one X-linked recessive variant, and a reproductive partner whose X chromosome does not contain that variant:

If a person with one X chromosome has the condition, they pass that X chromosome to every child who receives their X. They do not pass it to children who receive their Y. If the other biological parent does not carry the variant, children who inherit the affected parent's X are generally carriers rather than affected.

These are standard inheritance estimates, not predictions about a particular pregnancy. The calculation can change if both biological parents carry relevant variants, a parent has an unusual chromosome pattern, or the genetic finding has not been clearly classified.

A person with two X chromosomes can sometimes have an X-linked condition. Possible reasons include inheriting a disease-causing variant on both X chromosomes, having only one complete X chromosome, or having skewed X-inactivation . X-inactivation is the normal process by which cells switch off much of one X chromosome. If cells mainly switch off the working copy, a carrier may develop health effects.

- Each child who receives the partner's Y chromosome has a 1 in 2 chance of inheriting the variant
- Each child who receives the partner's X chromosome has a 1 in 2 chance of being a carrier
- Each pregnancy is a separate event, so earlier outcomes do not change later chances

## Why it matters for your health

Recognising this inheritance pattern can help explain why a condition appears mostly in people with one X chromosome, while passing through relatives with two X chromosomes who may have few or no related health effects.

Examples of X-linked recessive conditions include haemophilia A, haemophilia B, Duchenne muscular dystrophy, some forms of red-green colour vision deficiency, and

glucose-6-phosphate dehydrogenase deficiency. These are separate conditions with different tests, treatments, and health effects. The inheritance label does not show how severe a condition will be.

The pattern may help a genetics service estimate which relatives could carry a known family variant. It can also guide testing before or during pregnancy. Testing is usually most informative when the specific variant found in an affected relative is known.

Carrier status can sometimes matter for your own health. For example, some carriers of haemophilia variants have reduced clotting-factor levels and an increased tendency to bleed. Some carriers of dystrophin gene variants have a higher risk of heart muscle disease. Whether follow-up is useful depends on the specific gene and variant, not simply on the words "sex-linked recessive."

## What can go wrong

The inheritance pattern itself does not cause symptoms or become an illness. Problems can arise from the genetic condition linked to it or from misunderstanding a test result.

A family history may look inconsistent with X-linked recessive inheritance. Relatives may have been misdiagnosed, may not know their genetic relationship, or may have mild features that were never recognised. A new variant can also arise in an egg, sperm, or early embryo without being found in either parent's routine test.

A report may identify a variant of uncertain significance , often shortened to VUS. This means there is not enough evidence to decide whether the variant causes disease. A VUS does not by itself confirm a diagnosis or carrier status and should not be treated as though it were a known pathogenic variant.

"Carrier" can also be misunderstood as meaning "unaffected." Some carriers have measurable or important health effects. The relevance must be assessed for the named condition.

Testing only one gene or one family variant may miss other genetic causes. A negative result can be reassuring about the change tested, but it may not rule out every related condition.

The phrase "sex-linked recessive" is not an emergency finding. Routine follow-up with the clinician who ordered the test or a genetics service is usually appropriate. If you are pregnant or planning a pregnancy, arrange follow-up promptly because some testing choices are time-sensitive.

Emergency care should be based on actual symptoms or an existing condition's care plan, not this inheritance label. For example, bleeding that will not stop, severe breathing difficulty, collapse, or sudden severe weakness needs urgent assessment.

## What you can do

If this term appears on a report, identify the condition, gene, and exact variant being discussed. The inheritance pattern alone is not enough to interpret your health risk.

Useful questions for the ordering clinician or genetics service include:

Keep a copy of the laboratory report rather than relying on a remembered gene name. Reports may contain the exact DNA change needed for accurate testing in relatives.

A blood or saliva test can often look for a known family variant. Broader testing may be considered when the family's variant is unknown, but it can produce uncertain findings. A genetic counsellor can explain the possible results before testing.

Pregnancy-related options may include testing during pregnancy or testing embryos created through in vitro fertilisation. These choices have medical, practical, and personal implications. A genetics and reproductive care team can explain what is available and what each test can and cannot establish.

If the term appears only in general reading and there is no relevant personal or family history, there may be nothing you need to do.

- Is the variant classified as pathogenic, likely pathogenic, or uncertain?
- Does this result diagnose a condition, show carrier status, or only suggest testing?
- Could this finding affect my own health?
- Which relatives, if any, could benefit from targeted testing?
- Would the result change pregnancy or health-screening options?

## Common questions

Is sex-linked recessive the same as X-linked recessive?

Usually, yes. In routine genetics, sex-linked recessive generally refers to a recessive variant on the X chromosome. Y-linked traits follow a different pattern and are not usually described as recessive.

Can a carrier develop health problems?

Yes, for some conditions. X-inactivation, clotting-factor levels, and the function of the specific gene can affect a carrier's health. Carrier status should not automatically be assumed to have no medical effect.

Can a father pass an X-linked variant to a son?

Not through the usual X-linked inheritance route. A child who inherits the father's Y chromosome does not receive his X chromosome. Other relatives may still carry the same variant through different family lines.

Does a negative carrier test rule out all risk?

Not always. A targeted negative test means the known family variant was not found. Its meaning depends on the test's scope, the accuracy of the family diagnosis, and whether another genetic cause is possible.

Can the pattern skip generations?

It can appear to. A variant may pass through carriers who have no recognised health effects before appearing in a later relative with one X chromosome.

## Sources

- National Human Genome Research Institute. Sex Linked . Genetics glossary.
- MedlinePlus Genetics. What are the different ways a genetic condition can be inherited? . U.S. National Library of Medicine.
- MedlinePlus Genetics. What are the uses of genetic testing? . U.S. National Library of Medicine.
- Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants . Genetics in Medicine . 2015;17:405-424.
- Konkle BA, Nakaya Fletcher S. Hemophilia A . GeneReviews . University of Washington, Seattle.
- Darras BT, Urion DK, Ghosh PS. Dystrophinopathies . GeneReviews . University of Washington, Seattle.