CONDITION

Acoustic trauma

Learn how sudden loud sounds can injure the ear, what symptoms need urgent care, how hearing is tested, and which treatments and prevention steps may help.

WHAT IT IS
Ear injury from sudden extremely loud sound

BODY SYSTEM
Ears and hearing

MAIN SYMPTOMS
Muffled hearing, tinnitus, ear pain, dizziness

WHAT CAUSES IT
Sudden extremely loud sound

HOW IT IS DIAGNOSED
Otoscopic examination and audiogram

MAIN TREATMENT
Noise avoidance and hearing-loss treatment

CAN IT BE CURED?
Sometimes - temporary hearing changes may recover

IS IT AN EMERGENCY?
Sometimes - severe dizziness making standing unsafe

LONG-TERM OUTLOOK
Variable recovery with possible permanent hearing loss or tinnitus

Overview

Acoustic trauma is an ear injury caused by a sudden, extremely loud sound. Common examples include gunfire, explosions, fireworks, industrial accidents, and loud sound from speakers or headphones close to the ear.

The pressure wave from an intense sound can damage several parts of the ear. It may tear the eardrum, disturb the tiny middle-ear bones, or injure sensory cells and nerve connections in the inner ear. A blast can also push debris into the ear or cause head and balance-system injuries.

Acoustic trauma may cause conductive hearing loss , meaning sound cannot move normally through the outer or middle ear. It may instead cause sensorineural hearing loss , which results from damage to the inner ear or hearing nerve. Some people have both types.

Hearing may recover after a temporary change in sensitivity, sometimes called a temporary threshold shift. However, inner-ear sensory cells in humans do not grow back after they are destroyed. Some hearing loss or tinnitus may therefore be permanent.

Prompt assessment matters because sudden inner-ear hearing loss may benefit from treatment started soon after the injury. Blast exposure can also cause injuries outside the ear that need immediate attention.

Symptoms

Symptoms often begin immediately after the sound, although some may become more noticeable over the next several hours.

Possible symptoms include:

A torn eardrum can cause sharp pain followed by hearing loss, drainage, or bleeding. Pain may lessen quickly even though the injury remains.

Seek same-day medical care for sudden hearing loss, especially if it affects one ear more than the other. Sudden sensorineural hearing loss is time-sensitive and cannot be confirmed by symptoms alone.

Call emergency services or go to an emergency department after an explosion or major accident if there is severe head or neck injury, trouble breathing, confusion, loss of consciousness, weakness, facial drooping, repeated vomiting, clear fluid from the ear, heavy bleeding, or severe dizziness that makes standing unsafe. A blast can injure the lungs, brain, and other organs even when the ear symptoms seem most obvious.

- Reduced or muffled hearing in one or both ears
- Tinnitus, such as ringing, buzzing, hissing, or roaring
- Ear pain or a feeling of pressure or fullness
- Sound distortion or unusual sensitivity to everyday sounds
- Bleeding or fluid from the ear
- Dizziness, spinning, nausea, or poor balance
- Difficulty understanding speech, especially around background noise

Causes and risk factors

A single impulse sound can produce acoustic trauma. Gunshots and explosions are frequent

causes because they create a sharp pressure wave with very high sound intensity. Firecrackers, air horns, vehicle air bags, power tools, and feedback from amplified sound can also cause injury when they are close to the ear.

Risk depends on more than loudness alone. Important factors include:

Enclosed spaces can increase risk because sound reflects from walls and other hard surfaces. One ear may be more affected depending on body position or how a firearm was held.

People who work around weapons, aircraft, construction equipment, manufacturing machinery, or explosives may face repeated exposure. Musicians, event staff, and people who use loud personal audio equipment may also be at risk. Repeated noise exposure usually causes gradual noise-induced hearing loss rather than one clearly recognized injury, but an intense event can produce acute trauma on top of earlier damage.

Hearing protection lowers exposure but does not make all noise safe. Earplugs must fit properly, and earmuffs need a complete seal. In very loud settings, workplace safety programs may require both. Engineering controls, barriers, distance, and reduced exposure time remain important.

Some medicines and chemicals can harm the inner ear. These do not usually cause acoustic trauma by themselves, but they may increase concern about hearing damage when combined with noise exposure. A clinician may ask about medicines, solvents, and prior ear disease during the evaluation.

- How close the person is to the sound
- Whether the event occurs indoors or in another enclosed area
- How long the exposure lasts
- Whether loud sounds occur repeatedly
- Whether correctly fitted hearing protection is used
- The direction of the sound in relation to each ear
- Previous noise exposure or existing hearing problems

Diagnosis

A clinician will ask what happened, when symptoms began, whether one or both ears are affected, and whether hearing has changed since the event. Details about an explosion, head impact, ear drainage, dizziness, prior hearing loss, and hearing protection help guide the examination.

The outer ear canal and eardrum are usually examined with an otoscope, a lighted instrument used to look into the ear. This may reveal blood, debris, a torn eardrum, or other injury. Nothing should be pushed into an injured ear to clean it.

A hearing test, or audiogram , measures the quietest sounds a person can hear at different pitches. It also compares hearing through headphones with hearing measured through a small vibrator placed behind the ear. This helps distinguish middle-ear problems from inner-ear injury.

Speech testing may check how softly words can be heard and how clearly they are understood. Tympanometry measures how the eardrum and middle ear respond to changes in air pressure. It may not be appropriate in every fresh injury, particularly when the eardrum appears perforated.

When symptoms persist, a repeat audiogram can show whether hearing is recovering or whether a stable loss remains. Testing should not be delayed when there is a sudden, significant hearing change.

Imaging is not routinely needed for every loud-noise exposure. A CT scan may be considered when there is major head trauma, suspected damage to the middle-ear bones, a penetrating injury, or signs of a skull fracture. MRI may be used in selected cases of sensorineural hearing loss to evaluate the hearing nerve and nearby structures.

Because earwax, infection, middle-ear fluid, and other conditions can also reduce hearing, an examination and formal testing are needed to identify the likely type of injury.

Treatment and management

The first step is to move away from hazardous sound. Avoid further loud-noise exposure until the ear has been assessed and symptoms have stabilized. If noise cannot be avoided, properly fitted hearing protection is important.

Sudden hearing loss after acoustic trauma should be evaluated promptly. An ear, nose, and throat specialist, also called an otolaryngologist or ENT clinician, may be involved. An audiologist performs detailed hearing tests and can help with hearing rehabilitation.

Corticosteroids like prednisone or hydrocortisone are medicines that reduce inflammation. They may be offered for sudden sensorineural hearing loss after a discussion of potential benefits, timing, and risks. Depending on the situation, treatment may be given by mouth or placed through the eardrum into the middle ear. This is called an intratympanic injection. These treatments are not appropriate for every hearing change, and the choice depends on the examination, audiogram, medical history, and time since the injury.

Hyperbaric oxygen treatment allows a person to breathe oxygen in a pressurized chamber. It may be considered together with corticosteroids in selected cases of sudden sensorineural hearing loss. Access is limited, and it has its own risks and practical burdens. It is not routine treatment for every case of acoustic trauma.

Many traumatic eardrum perforations heal without surgery. The ear usually needs to be kept dry while healing. A clinician may advise avoiding swimming and protecting the ear during bathing. Do not place cotton swabs, nonprescribed ear drops, or other objects into the ear. Some drops can be unsafe when the eardrum is open.

Antibiotics like amoxicillin or doxycycline are not automatically needed for every acoustic injury or eardrum tear. They may be used when an injury is contaminated or an infection develops. Persistent perforations or disruption of the middle-ear bones may require surgical repair.

Dizziness and balance problems may improve as the ear recovers. Persistent symptoms may require vestibular testing or balance therapy. Medicines that suppress dizziness are sometimes used briefly, but prolonged use can interfere with the brain's adjustment to a balance injury.

If hearing loss remains, hearing aids or other hearing devices may improve communication. Severe loss may require evaluation for implantable hearing technology. Tinnitus management can include hearing correction, sound therapy, sleep support, and counseling approaches that reduce the distress caused by the sound. No supplement has been proven to restore sensory cells destroyed by noise.

Practical communication steps can help during recovery. Reduce background noise, face the person speaking, and ask others to speak clearly rather than shout. Shouting may distort speech and can be uncomfortable when sound sensitivity is present.

Complications

Possible complications include permanent hearing loss, ongoing tinnitus, sound sensitivity, and difficulty understanding speech in noisy places. Even when a standard hearing test improves, some people continue to notice trouble following conversation or locating where sounds come from.

A persistent eardrum opening can increase the risk of middle-ear infection and may cause continuing conductive hearing loss. Scarring or movement of the middle-ear bones can also affect sound transmission.

Inner-ear damage may cause lasting dizziness or imbalance. Tinnitus and hearing loss can

interfere with sleep, concentration, work, relationships, and mood. These effects are treatable even when the underlying hearing damage cannot be reversed.

Repeated noise exposure can add to existing injury. Preventing another exposure is an important part of limiting further hearing loss.

Follow up

Follow-up timing depends on the injury and initial test results. A clinician may arrange a repeat hearing test within days or weeks to document recovery. Continued ENT or audiology care is appropriate when hearing loss, tinnitus, drainage, pain, or balance symptoms persist.

Return for prompt medical review if hearing suddenly worsens, drainage develops, pain increases, fever appears, or dizziness returns. These changes can suggest infection or another complication.

Emergency care is appropriate for new facial weakness, severe neurologic symptoms, inability to walk safely, heavy bleeding, clear ear drainage after head trauma, or breathing problems after a blast.

For routine prevention, ask about workplace noise monitoring, properly fitted hearing protection, and periodic hearing tests if loud sound is part of work or recreation. Hearing protectors should be worn consistently and replaced when damaged or poorly fitting.

Sources

- National Institute on Deafness and Other Communication Disorders. Noise-Induced Hearing Loss . National Institutes of Health.
- Chandrasekhar SS, Tsai Do BS, Schwartz SR, et al. Clinical Practice Guideline: Sudden Hearing Loss (Update) . Otolaryngology-Head and Neck Surgery . 2019;161(1 Suppl):S1-S45.
- Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Preventing Occupational Noise-Induced Hearing Loss .
- Occupational Safety and Health Administration. Occupational Noise Exposure, 29 CFR 1910.95 .
- Le TN, Straatman LV, Lea J, Westerberg B. Current insights in noise-induced hearing loss: a literature review of the underlying mechanism, pathophysiology, asymmetry, and management options . Journal of Otolaryngology-Head & Neck Surgery . 2017;46:41.

- World Health Organization. World Report on Hearing . 2021.