CONDITION

Agenesis of commissura magna cerebri

Learn about agenesis of the corpus callosum, including possible symptoms, causes, diagnosis, treatment, developmental support, and follow-up.

WHAT IT IS
Absence of all or part of the corpus callosum

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Developmental delays, learning difficulties, coordination problems, seizures

WHAT CAUSES IT
Genetic, prenatal infection or exposure, often unknown

HOW IT IS DIAGNOSED
Fetal or postnatal brain MRI

MAIN TREATMENT
Supportive therapies and symptom management

CAN IT BE CURED?
No, no treatment can create it

IS IT AN EMERGENCY?
Sometimes, seizure lasting about five minutes

LONG-TERM OUTLOOK
Highly variable and usually nonprogressive

Overview

Agenesis of commissura magna cerebri is another name for agenesis of the corpus callosum . The corpus callosum is a thick band of nerve fibers that connects the brain's right and left halves. It helps the two sides exchange and combine information.

"Agenesis" means that a structure did not form. In complete agenesis, the corpus callosum is absent. In partial agenesis, only part of it developed. Related terms include hypoplasia, meaning the structure is unusually thin, and dysgenesis, meaning it developed differently from expected.

This condition begins before birth. It is not the same as damage to a previously normal corpus callosum from a stroke, injury, infection, or surgery.

The effect on a person's health varies greatly. Some people with an isolated finding have average intelligence, few physical symptoms, and no major limitations. Others have difficulties with learning, movement, communication, seizures, or daily activities. Challenges may be more significant when other brain differences, chromosome changes, or genetic conditions are also present.

Even when early development seems typical, subtle differences may become clearer as school, work, and social situations grow more complex. These may involve problem-solving, understanding indirect language, adjusting to unfamiliar situations, or recognizing social cues. Brain imaging alone cannot predict an individual child's exact development.

Symptoms

Some people have no obvious symptoms, and the condition is found unexpectedly during brain imaging. When symptoms occur, they depend partly on whether agenesis is isolated or occurs with other medical conditions.

Possible features include:

Infants with more complex brain or genetic conditions may have poor feeding, low muscle tone, trouble gaining weight, or marked developmental delays. However, these findings are not specific to corpus callosum agenesis and require a broader medical evaluation.

Symptoms can change with age. A young child may meet early milestones but later have difficulty when tasks require several types of information to be processed at once. Teenagers and adults may need support with time management, social judgment, independent living, or adapting to rapid change.

Having the imaging finding does not mean that every listed symptom will occur. It also does not establish the cause of a person's learning, behavioral, or neurological concerns.

- Delays in sitting, walking, speaking, or other developmental skills
- Learning difficulties, especially with complex or unfamiliar information
- Problems with planning, organization, flexible thinking, or problem-solving
- Difficulty understanding humor, sarcasm, emotions, or unspoken social rules
- Poor coordination, balance problems, or unusual muscle tone
- Feeding or swallowing difficulties, particularly during infancy
- Seizures
- Vision or hearing problems

- Differences in head growth
- Sleep, attention, anxiety, or behavior concerns

Causes and risk factors

The corpus callosum forms during early brain development before birth. Agenesis occurs when the nerve fibers that normally cross between the brain's two halves do not develop or follow their expected path.

Many different biological processes can interfere with this development. Possible causes include:

Agenesis can occur by itself or as one feature of a syndrome affecting several parts of the body. The likelihood of additional medical or developmental concerns is generally higher when imaging shows other brain abnormalities.

In many families, no specific cause is found. Parents should not assume that the condition resulted from something they did or did not do during pregnancy.

The chance of the condition occurring again in a future pregnancy depends on its cause. It may be low when testing finds no genetic explanation, or it may be higher when an inherited or chromosome-related condition is identified. A genetic counselor can review the family history, test results, and available reproductive options without assuming that one recurrence estimate applies to every family.

- A change in a single gene involved in brain development
- Missing, extra, or rearranged chromosome material
- A broader genetic or metabolic condition
- Another fetal brain malformation
- Certain prenatal infections or exposures
- An interruption in early fetal brain development for which no cause can be identified

Diagnosis

Agenesis may be suspected before birth during a prenatal ultrasound. Certain changes in the shape or position of structures inside the fetal brain can suggest that the corpus callosum has not developed normally. A detailed ultrasound, sometimes called targeted neurosonography, is usually needed to examine the brain more closely.

Fetal magnetic resonance imaging, or fetal MRI, may provide additional detail. MRI uses a strong magnetic field rather than ionizing radiation. It can help confirm the finding and look for other brain differences that may be difficult to see on ultrasound. The usefulness and timing of fetal MRI depend on the pregnancy and local expertise.

After birth, brain MRI is generally the most detailed imaging test for evaluating the corpus callosum. It can show whether agenesis is complete or partial and whether other structures are affected. Ultrasound through an infant's soft spot or computed tomography may identify some abnormalities, but these tests do not provide the same detail as MRI.

Diagnosis often includes more than imaging. Depending on the person's age and findings, evaluation may involve:

Genetic testing may include chromosome analysis, chromosomal microarray, or sequencing of individual genes or many genes. The choice depends on the imaging findings, physical features, family history, and testing already completed. Testing does not identify a cause in every case.

Prenatal imaging can provide important information, but it cannot precisely predict future intelligence, behavior, or independence. Prognosis is usually clearer over time through developmental observation and formal assessment.

- A neurological and physical examination
- Review of pregnancy, birth, development, and family history
- Developmental, psychological, speech, or school-based assessment
- Vision and hearing testing
- An electroencephalogram, or EEG, if seizures are suspected
- Genetic counseling and appropriate genetic testing

Treatment and management

There is no treatment that can create a missing corpus callosum. Management focuses on the person's symptoms, development, health, education, and ability to participate in daily life.

Babies and young children may benefit from early-intervention services. These can include physical therapy for movement and balance, occupational therapy for daily skills and sensory needs, and speech-language therapy for communication or feeding. Support should be based on the child's actual needs rather than the MRI result alone.

School-age children may need a learning assessment and an individualized education plan or

other classroom accommodations. Helpful approaches can include clear instructions, visual schedules, extra time, repetition, and breaking complex tasks into smaller steps. Some students understand familiar facts well but need more help applying them in a new setting.

Speech-language therapy may address more than pronunciation. It can also support understanding of indirect language, conversation, storytelling, and social communication. Occupational therapy may help with handwriting, coordination, self-care, organization, or adapting the environment.

Seizures are treated with antiseizure medicine selected by a clinician according to seizure type, age, other health conditions, and possible side effects. Not everyone with agenesis has seizures or needs medication. Persistent seizures may require care from a neurologist or epilepsy specialist.

Feeding problems may be evaluated by a feeding therapist, speech-language pathologist, dietitian, or gastroenterology team. A swallowing study may be appropriate if there is coughing, choking, recurrent chest illness, or concern that food or liquid is entering the airway.

Associated conditions may need separate treatment. For example, hydrocephalus-an abnormal buildup of fluid in the brain-may occasionally require neurosurgical care. Surgery does not treat the absent corpus callosum itself.

Counseling or behavioral health care can help with anxiety, frustration, attention problems, social stress, or adjustment to a lifelong condition. Support should build on the person's strengths and goals rather than focusing only on deficits.

Complications

Possible complications vary and are often related to associated brain or genetic conditions. They may include epilepsy, persistent learning disabilities, motor impairment, feeding problems, or reduced independence with complex daily tasks.

Communication and social difficulties can affect friendships, school, employment, and mental health. A person may speak fluently but still have trouble interpreting context, recognizing another person's viewpoint, or responding to an unexpected situation. These challenges can be missed unless assessment includes higher-level language and practical functioning.

Swallowing impairment can increase the risk of poor nutrition or food entering the airway. Poorly controlled seizures may cause injury and interfere with learning or sleep. Hydrocephalus and other structural brain abnormalities may produce complications separate from those caused by callosal agenesis.

The diagnosis itself is not usually a progressive disease. The corpus callosum does not

continue to disappear. However, support needs may change as life demands become more complex.

Follow up

Routine follow-up should be tailored to symptoms and age. Children commonly need developmental monitoring at regular health visits, with formal reassessment if parents, teachers, or clinicians notice new concerns. Vision, hearing, growth, feeding, movement, learning, behavior, and seizure control may all need review.

Neurology follow-up is often appropriate for seizures, abnormal movements, significant muscle-tone differences, developmental regression, or complex MRI findings. Genetics follow-up may be useful when testing changes over time or when a person is considering pregnancy. Adults may benefit from reassessment when planning education, employment, transportation, finances, or independent living.

Seek prompt medical advice for a new seizure, repeated unexplained staring or jerking episodes, loss of previously gained skills, worsening headaches, persistent vomiting, or a major change in alertness or behavior.

Call emergency services for:

A clinician can provide an individualized seizure action plan when seizures are part of the condition. Routine questions about learning, behavior, therapies, or school supports can usually be addressed at scheduled follow-up visits.

- A seizure lasting about five minutes or longer
- Repeated seizures without recovery between them
- Trouble breathing, blue or gray skin, or severe choking
- Sudden weakness, loss of consciousness, or inability to awaken
- A severe head injury or other immediately life-threatening concern

Sources

- National Institute of Neurological Disorders and Stroke. "Agenesis of the Corpus Callosum." National Institutes of Health, health information resource.



- Paul LK, Brown WS, Adolphs R, et al. "Agenesis of the Corpus Callosum: Genetic, Developmental and Functional Aspects of Connectivity." Nature Reviews Neuroscience . 2007;8(4):287-299. doi:10.1038/nrn2107.
- Edwards TJ, Sherr EH, Barkovich AJ, Richards LJ. "Clinical, Genetic and Imaging Findings Identify New Causes for Corpus Callosum Development Syndromes." Brain . 2014;137(6):1579-1613. doi:10.1093/brain/awt358.
- Palmer EE, Mowat D. "Agenesis of the Corpus Callosum: A Clinical Approach to Diagnosis." American Journal of Medical Genetics Part C: Seminars in Medical Genetics . 2014;166C(2):184-197. doi:10.1002/ajmg.c.31405.
- D'Antonio F, Pagani G, Familiari A, et al. "Outcomes Associated With Isolated Agenesis of the Corpus Callosum: A Meta-analysis." Pediatrics . 2016;138(3):e20160445. doi:10.1542/peds.2016-0445.
- International Society of Ultrasound in Obstetrics and Gynecology. "ISUOG Practice Guidelines (Updated): Sonographic Examination of the Fetal Central Nervous System. Part 2: Performance of Targeted Neurosonography." Ultrasound in Obstetrics & Gynecology . 2021;57(4):661-671. doi:10.1002/uog.23616.