CONDITION

Alcoholic ketoacidosis

Learn how alcoholic ketoacidosis develops, what symptoms require emergency care, and how clinicians diagnose, treat, and help prevent another episode.

WHAT IT IS
Serious metabolic problem causing acidic ketone buildup

BODY SYSTEM
Metabolism

MAIN SYMPTOMS
Persistent vomiting, upper abdominal pain, dizziness, fast deep breathing

WHAT CAUSES IT
Heavy alcohol use, starvation, vomiting, dehydration

HOW IT IS DIAGNOSED
History, examination, blood ketones and acid-base tests

MAIN TREATMENT
Intravenous fluids, glucose, thiamine, electrolyte replacement

CAN IT BE CURED?
Yes, prompt treatment usually corrects it

IS IT AN EMERGENCY?
Yes, vomiting and poor intake after heavy drinking

LONG-TERM OUTLOOK
Most people improve with prompt treatment

Overview

Alcoholic ketoacidosis is a serious metabolic problem that can develop when a person who has been drinking heavily stops eating and continues to vomit. It is also called alcohol-related ketoacidosis.

The condition happens when the body does not have enough available glucose, its main source of energy. The liver begins turning fat into chemicals called ketones. Ketones are acidic. When they build up in the blood, they can cause a high-anion-gap metabolic acidosis, meaning that the blood becomes too acidic because of unmeasured acids.

Alcoholic ketoacidosis often develops after several days of heavy alcohol use followed by a sudden decrease in drinking because of nausea, vomiting, abdominal pain, or illness. During this period, the person may eat very little and become dehydrated. The blood alcohol level may

be low or undetectable by the time medical care is sought.

Unlike diabetic ketoacidosis, alcoholic ketoacidosis often occurs with a normal, low, or only mildly elevated blood glucose level. However, the two conditions can sometimes be difficult to distinguish, especially in someone who also has diabetes.

Alcoholic ketoacidosis requires prompt assessment in an emergency department. Treatment is usually effective, but clinicians must also look for related problems such as alcohol withdrawal, pancreatitis, infection, internal bleeding, liver disease, or exposure to a toxic alcohol.

Symptoms

Symptoms commonly begin after heavy alcohol use followed by poor food intake and repeated vomiting. They may develop over hours to a few days.

Possible symptoms include:

Many people with alcoholic ketoacidosis remain awake and able to answer questions, even when the chemical changes in their blood are significant. Confusion or a reduced level of alertness is less typical and may point to another or additional problem. Possibilities include low blood glucose, a head injury, infection, severe alcohol withdrawal, liver-related brain dysfunction, or another substance.

Symptoms that may suggest alcoholic ketoacidosis need same-day emergency evaluation, particularly when vomiting and poor food intake follow heavy alcohol use. Do not try to correct the condition by drinking more alcohol.

Call emergency services for fainting, a seizure, severe confusion, inability to wake, breathing difficulty, vomiting blood, black stools, severe or worsening abdominal pain, chest pain, or signs of shock such as cold or clammy skin with marked weakness. A person who may be seriously ill should not drive to the hospital.

- Persistent nausea and vomiting
- Pain or tenderness in the upper abdomen
- Extreme thirst or a dry mouth
- Weakness, dizziness, or difficulty standing
- Fast, deep breathing
- A rapid heartbeat
- Reduced urination

- Agitation, shakiness, sweating, or anxiety from alcohol withdrawal

Causes and risk factors

Alcoholic ketoacidosis usually results from several changes occurring at the same time.

Heavy alcohol use can reduce food intake. Vomiting then causes further loss of fluid, salt, and nutrients. As the liver's stored carbohydrate is used up, the body has less glucose available for energy. Insulin levels fall, while stress hormones signal the body to release fat. The liver converts this fat into ketones.

Alcohol metabolism also changes the liver's chemical balance. This favors production of beta-hydroxybutyrate, the main ketone found in alcoholic ketoacidosis, and can interfere with the liver's ability to make new glucose. Dehydration reduces the kidneys' ability to clear ketones, allowing the acids to accumulate.

Factors associated with greater risk include:

Alcoholic ketoacidosis does not affect everyone who drinks alcohol. It reflects a specific combination of alcohol exposure, starvation, vomiting, dehydration, and hormone changes. It can also occur alongside other urgent conditions rather than as the only diagnosis.

- A period of heavy or prolonged alcohol use
- Little or no food intake
- Repeated vomiting
- Dehydration
- An alcohol use disorder
- A recent illness, injury, infection, or major physical stress
- Conditions such as pancreatitis that cause pain, vomiting, and poor intake

Diagnosis

There is no single test that confirms alcoholic ketoacidosis in every case. Clinicians use the person's recent history, physical examination, and laboratory results while checking for other causes of illness.

The medical team may ask about alcohol use, the last alcoholic drink, recent meals, vomiting, diabetes, medicines, substance exposure, and previous withdrawal symptoms. Honest information helps guide safe care. It should be used to support treatment, not to judge the

person.

Blood testing commonly shows a high-anion-gap metabolic acidosis and increased ketones. A beta-hydroxybutyrate blood test is especially useful. Standard urine ketone tests mainly detect a different ketone called acetoacetate, so a urine result may appear only mildly positive even when beta-hydroxybutyrate is high.

Blood glucose is often normal or low, but it may be elevated. A high glucose result does not automatically establish diabetic ketoacidosis, particularly in someone with diabetes or recent stress. Clinicians consider the full pattern before choosing treatment.

Other tests may assess:

Vomiting can cause a second chemical disturbance that partly hides the acidosis. Because of this mixed pattern, the blood pH may not be as low as expected.

Depending on the circumstances, imaging or additional blood tests may be needed to look for pancreatitis, pneumonia, internal bleeding, head injury, sepsis, or ingestion of methanol or ethylene glycol. Diabetic ketoacidosis, starvation ketoacidosis, lactic acidosis, kidney failure, and certain poisonings can produce similar laboratory findings.

- Sodium, potassium, magnesium, phosphate, and kidney function
- Blood glucose and beta-hydroxybutyrate
- Blood acidity and carbon dioxide levels
- Lactate, which can rise with dehydration or other illness
- Liver and pancreas tests
- A blood count and markers of infection
- Blood alcohol level
- An electrocardiogram to check the heart rhythm

Treatment and management

Alcoholic ketoacidosis is treated in a hospital or emergency department. The main goals are to restore fluid and glucose, stop ketone production, replace missing electrolytes, and manage the cause of vomiting or poor intake.

Clinicians usually give intravenous fluids containing saline and glucose. Glucose encourages the body to release insulin, which helps stop fat breakdown and ketone production. Fluids improve circulation and help the kidneys remove acids.

Thiamine, or vitamin B1, is commonly given before or along with glucose because people with

prolonged heavy alcohol use may have low thiamine stores. Thiamine helps the brain and body use carbohydrates. If blood glucose is dangerously low, glucose treatment should not be delayed while waiting for thiamine.

Potassium, magnesium, and phosphate may be low at presentation or may fall during treatment. They are checked repeatedly and replaced when needed. Heart monitoring may be used when electrolyte abnormalities could affect the heart rhythm.

Insulin is generally not the main treatment for alcoholic ketoacidosis. It may cause dangerously low blood glucose if given when glucose is already normal or low. Clinicians may use insulin when diabetic ketoacidosis cannot be excluded or when significant hyperglycemia requires it.

Bicarbonate, a medicine that reduces acidity, is not routinely needed because fluids and glucose usually correct the underlying problem. Its use is reserved for selected situations based on the severity and cause of the acidosis.

The care team also treats associated problems. Anti-nausea treatment may help control vomiting. When the person can safely eat, food provides carbohydrates and other nutrients. Pancreatitis, infection, bleeding, injury, or another illness requires its own treatment.

Alcohol withdrawal can begin while alcoholic ketoacidosis is being treated. Staff monitor for tremor, sweating, anxiety, hallucinations, seizures, and severe confusion. Medication may be given according to a monitored withdrawal plan.

After the immediate illness improves, clinicians may discuss treatment for alcohol use disorder. Options can include counseling, peer support, structured treatment programs, and medications that reduce alcohol cravings or help support abstinence. The best approach depends on the person's goals, health, and access to care.

Complications

Most people improve when the condition is recognized and treated promptly. The ketone level and acid-base balance often begin to correct after fluids and glucose are given. However, complications can result from the metabolic disturbance or from illnesses occurring with it.

Potential complications include:

A change in alertness should not simply be blamed on alcohol. It requires assessment for these and other possible causes.

- Severely low blood glucose
- Abnormal heart rhythms related to low potassium or magnesium

- Low blood pressure and poor circulation from dehydration
- Acute kidney injury
- Aspiration of vomit into the lungs
- Thiamine deficiency and Wernicke encephalopathy, a brain disorder affecting awareness, balance, and eye movements
- Alcohol withdrawal seizures or delirium
- Complications of pancreatitis, infection, bleeding, or liver disease

Follow up

Before discharge, the medical team generally confirms that vomiting is controlled, food and fluids can be taken safely, and important laboratory abnormalities are improving. The underlying reason for poor intake or abdominal pain should also be addressed.

Follow the discharge plan for repeat blood tests, nutrition, medicines, and appointments. A primary care clinician can review recovery, liver and pancreas health, nutrition, diabetes risk, and any ongoing symptoms. If alcohol use has become difficult to control, follow-up can also connect the person with addiction medicine, counseling, or community services.

Stopping alcohol suddenly can be dangerous for someone who is physically dependent. Symptoms such as shaking, sweating, anxiety, or insomnia should be discussed with a clinician. A history of withdrawal seizures, hallucinations, or delirium is especially important when planning a reduction or stopping alcohol.

Return for urgent medical assessment if vomiting comes back, food or fluids cannot be kept down, urination decreases, abdominal pain worsens, or withdrawal symptoms develop. Call emergency services for a seizure, fainting, severe confusion, hallucinations with unsafe behavior, breathing difficulty, vomiting blood, black stools, or inability to wake.

Longer-term prevention focuses on avoiding another cycle of heavy drinking, starvation, vomiting, and dehydration. A practical plan may include regular meals, early care for persistent vomiting, and evidence-based treatment for alcohol use disorder.

Sources

- Long B, Lentz S, Gottlieb M. "Alcoholic Ketoacidosis: Etiologies, Evaluation, and Management." The Journal of Emergency Medicine . 2021;61(6):658-665.

- McGuire LC, Cruickshank AM, Munro PT. "Alcoholic Ketoacidosis." Emergency Medicine Journal . 2006;23(6):417-420.
- MSD Manual Professional Edition. "Alcoholic Ketoacidosis." Merck & Co., Inc.; reviewed October 2023.
- The American Society of Addiction Medicine. The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management . Journal of Addiction Medicine . 2020;14(3S Suppl 1):1-72.
- National Institute for Health and Care Excellence. Alcohol-Use Disorders: Diagnosis and Management of Physical Complications . Clinical guideline CG100. Published 2010; updated 2017.
- Kraut JA, Madias NE. "Metabolic Acidosis: Pathophysiology, Diagnosis and Management." Nature Reviews Nephrology . 2010;6(5):274-285.