CONDITION

Allergy to natural rubber

Learn how natural rubber latex allergy causes skin, breathing, and whole-body reactions, and how it is diagnosed, treated, and managed safely.

WHAT IT IS
Immune allergy to natural rubber latex proteins

BODY SYSTEM
Immune system

MAIN SYMPTOMS
Hives, swelling, wheezing, dizziness or fainting

WHAT CAUSES IT
Immune sensitization to natural rubber proteins

HOW IT IS DIAGNOSED
Latex-specific IgE blood test, specialist skin-prick testing

MAIN TREATMENT
Latex avoidance and emergency allergy management

CAN IT BE CURED?
No, no routine permanent cure exists

IS IT AN EMERGENCY?
Sometimes, breathing difficulty, throat swelling, collapse, or rapid worsening

LONG-TERM OUTLOOK
Variable course with more frequent reactions after repeated exposure

Overview

Natural rubber latex is a milky fluid obtained mainly from the rubber tree, Hevea brasiliensis . It is used to make some medical gloves, balloons, rubber bands, elastic products, condoms, dental dams, and other items.

A natural rubber latex allergy is an immune reaction to proteins in latex. The most important form is an immediate, or IgE-mediated, allergy. IgE is an antibody that can trigger symptoms within minutes after latex touches the skin or moist surfaces such as the eyes, mouth, or airways. Reactions can range from hives to a severe whole-body reaction called anaphylaxis.

Not every rash from a rubber product is a latex allergy. Two other conditions are common:

Distinguishing these conditions matters because their risks and management are different.

Someone with immediate latex allergy may need special planning before medical or dental care. A person with contact dermatitis may instead need patch testing and avoidance of a specific rubber-processing chemical.

Many healthcare settings now use non-latex products, especially non-powdered synthetic gloves. However, natural rubber latex may still be present in medical, workplace, and household items.

- Irritant contact dermatitis is dry, cracked, or sore skin caused by friction, sweating, repeated handwashing, or chemicals. It does not involve an allergy.
- Allergic contact dermatitis is a delayed immune reaction, usually to chemicals added during rubber manufacturing rather than to latex proteins.

Symptoms

Symptoms depend on the type of reaction, the amount of exposure, and how the latex enters the body. A person's past reaction does not reliably predict the severity of a future reaction.

Immediate latex allergy can cause:

Anaphylaxis can affect several parts of the body at once. Breathing difficulty, throat or tongue swelling, collapse, or rapidly worsening symptoms after latex exposure require emergency care. If an epinephrine auto-injector has been prescribed, use it as directed and call emergency services. Antihistamines like cetirizine or loratadine do not replace epinephrine for anaphylaxis.

Airborne reactions are possible when latex proteins attach to glove powder and enter the air. Powdered medical gloves have been banned in the United States, which has reduced this type of exposure in healthcare settings.

Allergic contact dermatitis usually develops hours to one or two days after contact. It may cause an itchy, blistered, scaly, or spreading rash. Irritant dermatitis is more likely to cause burning, dryness, cracking, and soreness limited to exposed skin.

Some people with latex allergy also react to foods containing similar proteins. This is sometimes called latex-fruit syndrome. Banana, avocado, kiwi, and chestnut are well-known examples, but many people with latex allergy eat these foods without difficulty. Foods should not be removed from the diet solely because of a latex allergy label.

- Itching, redness, or hives where latex touched the skin
- Swelling of the lips, face, tongue, or throat
- Itchy, watery eyes or a runny nose

- Coughing, wheezing, chest tightness, or shortness of breath
- Stomach cramps, vomiting, or diarrhea
- Dizziness, weakness, confusion, or fainting

Causes and risk factors

Immediate latex allergy develops when the immune system becomes sensitized to natural rubber proteins. After sensitization, later exposure may release chemicals such as histamine, producing allergy symptoms.

Exposure can occur through direct skin contact, breathing airborne particles, or contact with the mouth, eyes, genitals, or internal tissues. Medical procedures can create higher-risk exposure because latex-containing equipment may contact moist surfaces or tissue.

Repeated exposure increases the chance of sensitization. Groups with a higher recognized risk include:

Having a risk factor does not mean that a person has latex allergy. Likewise, occasional contact with a rubber item does not prove that latex caused a symptom. Products may contain synthetic rubber, adhesives, dyes, powders, or other substances that can also irritate the skin or cause allergy.

- People who have had many surgeries or medical procedures, especially beginning in childhood
- People with spina bifida or certain conditions affecting the urinary tract
- Healthcare, dental, laboratory, rubber-industry, and other workers with frequent latex exposure
- People with eczema, asthma, hay fever, or other allergic conditions
- People who have previously developed symptoms around latex products
- People with immediate allergic reactions to foods associated with latex cross-reactivity

Diagnosis

Diagnosis begins with a detailed medical history. A clinician may ask which product was involved, how quickly symptoms began, whether the product contained natural rubber latex, and whether similar reactions have occurred during healthcare, dental work, balloon exposure, or condom use.

It is helpful to describe the exact symptoms and their timing. Photographs of a rash and packaging from the suspected product may provide useful clues. The clinician may also ask about work exposures, previous operations, asthma, eczema, and reactions to foods.

Testing depends on the suspected type of reaction.

A blood test for latex-specific IgE can support a diagnosis of immediate allergy. However, a positive result does not always mean that latex causes clinical symptoms, and a negative result does not completely rule out allergy.

Skin-prick testing may be used by allergy specialists in regions where suitable standardized materials and safe testing procedures are available. Because testing can itself trigger a reaction, it should not be attempted with household products or outside an appropriate medical setting.

Patch testing is different from skin-prick testing. It places small amounts of suspected chemicals on the skin for several days and is used to investigate delayed allergic contact dermatitis. It does not diagnose immediate allergy to latex proteins.

In some cases, the history remains more informative than laboratory testing. Deliberate exposure or challenge testing is rarely appropriate and, when considered, requires specialist supervision with emergency treatment available.

Treatment and management

Avoiding natural rubber latex is the main way to prevent reactions. There is no routine cure that permanently removes the allergy. The amount of avoidance needed should be based on the type of reaction and advice from an allergy specialist.

Tell doctors, dentists, nurses, pharmacists, employers, and other relevant people about a confirmed or strongly suspected allergy. Ask for the information to be placed clearly in medical and dental records. Before surgery or a procedure, contact the facility in advance so staff can prepare an appropriate latex-safe environment and check equipment.

Synthetic alternatives include nitrile, neoprene, and vinyl. A product labeled "synthetic rubber" does not contain natural rubber latex proteins, although other ingredients may still cause irritation or contact allergy. Product labels and manufacturer information may be needed because appearance alone cannot identify latex.

People who use barrier contraception should discuss non-latex condoms or other options with a healthcare professional. Alternative products may differ in protection against pregnancy and sexually transmitted infections.

A clinician may recommend an anaphylaxis action plan and prescribe an epinephrine auto-injector for someone considered at risk of a severe immediate reaction. The person and those close to them should know when and how to use it. Auto-injectors should be stored as directed, kept accessible, and replaced before expiration.

A medical identification bracelet or card can alert others to the allergy if the person cannot communicate. Carrying a written list of products that have caused reactions may also help.

Antihistamines may ease limited itching or hives, but they do not prevent or treat dangerous breathing or circulation problems. Skin treatment for irritant or allergic contact dermatitis may include moisturizers, changes in glove or skin-care practices, and clinician-directed anti-inflammatory treatment.

At work, management may involve an occupational health service. Changes can include non-latex gloves, safer equipment, education, and control of airborne or surface contamination. People should not repeatedly test suspected products on their own skin.

Food avoidance is needed only when a food has caused symptoms or a qualified clinician recommends it after assessment. Unnecessary restriction can make a balanced diet harder to maintain.

Complications

The most serious complication of immediate latex allergy is anaphylaxis. Risk may be greater when latex contacts internal tissue or moist surfaces during an operation, examination, dental procedure, or use of certain devices.

Repeated exposure may lead to more frequent reactions, although the course varies from person to person. Asthma symptoms can also become difficult to control if workplace exposure continues.

Ongoing hand dermatitis can cause painful cracks and interfere with sleep or work. Damaged skin may also make exposure to irritants more troublesome. Identifying whether the problem is irritation, delayed contact allergy, or immediate latex allergy can help prevent ineffective restrictions.

An incorrect allergy label can also have consequences. It may create anxiety or complicate medical care without improving safety. On the other hand, an unrecognized allergy may lead to preventable exposure. A careful specialist assessment is useful when the diagnosis is uncertain.

Follow up

Routine follow-up may be appropriate with a primary care clinician, allergist, dermatologist, or occupational health professional. Follow-up can review test results, workplace or household exposures, skin care, and preparation for future procedures.

People prescribed epinephrine should periodically review their action plan and injection technique. They should also check expiration dates and discuss how many devices to have available based on their clinician's advice.

Contact a clinician promptly if reactions are becoming more frequent, asthma or wheezing is occurring with exposure, dermatitis is persistent or infected, or a new medical or dental procedure is planned. New food reactions should also be assessed rather than managed through broad food avoidance.

Seek emergency help for trouble breathing, throat or tongue swelling, fainting, confusion, collapse, or rapidly worsening symptoms after possible latex exposure. Use prescribed epinephrine without delay when the action plan indicates it, then call emergency services because symptoms can return after initial improvement.

Sources

- American Academy of Allergy, Asthma & Immunology. "Latex Allergy." Patient information reviewed through the AAAAI public education resource.
- American College of Allergy, Asthma & Immunology. "Latex Allergy." Patient education resource, Allergies section.
- Australasian Society of Clinical Immunology and Allergy. "Latex Allergy." Patient and clinician information resource.
- U.S. Food and Drug Administration. "Banned Devices; Ban Powdered Surgeon's Gloves, Powdered Patient Examination Gloves, and Absorbable Powder for Lubricating a Surgeon's Glove." Federal Register . 2016;81(243):91722-91731.
- National Institute for Occupational Safety and Health. Preventing Allergic Reactions to Natural Rubber Latex in the Workplace . DHHS (NIOSH) Publication No. 97-135; 1997.
- Kelly KJ, Sussman G. "Latex Allergy: Where Are We Now and How Did We Get There?" The Journal of Allergy and Clinical Immunology: In Practice . 2017;5(5):1212-1216. doi:10.1016/j.jaip.2017.05.029.