CONDITION

Autism spectrum disorder - asperger syndrome

Learn how Asperger syndrome relates to autism spectrum disorder, including common traits, assessment, support options, complications, and follow-up care.

WHAT IT IS
Lifelong neurodevelopmental condition

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Social communication differences, repetitive behaviors, focused interests, sensory differences

WHAT CAUSES IT
Complex genetic and developmental factors

HOW IT IS DIAGNOSED
Developmental history and direct observation

MAIN TREATMENT
Individualized support and accommodations

CAN IT BE CURED?
No, lifelong and not an illness

IS IT AN EMERGENCY?
Sometimes, immediate danger of self-harm or suicide

LONG-TERM OUTLOOK
Lifelong with widely varying support needs

Overview

Autism spectrum disorder, often called autism, is a lifelong neurodevelopmental condition that affects social communication, patterns of behavior or interests, sensory processing, and responses to change.

Asperger syndrome was previously a separate diagnosis, usually for autistic people without significant language or intellectual delays. Current diagnostic systems include it within autism spectrum disorder. People previously diagnosed with Asperger syndrome do not usually need a new evaluation solely because the terminology changed.

Some people continue to describe themselves as having Asperger syndrome. Others prefer "autistic person," "person with autism," or another term. Individual preferences should be respected.

Autism is a spectrum because its features and effects vary widely. One person may need substantial daily support, while another may live independently but have difficulty with social demands, sensory overload, organization, or unexpected change. Needs can vary by setting and over time.

Autism is not an illness that needs to be cured. Assessment may help someone understand their experiences, request accommodations, and obtain support for communication, education, employment, health, or daily living.

Symptoms

Autistic traits begin during early development, although they may not become clear until social or organizational demands exceed a person's ability to cope. Some people are not identified until adolescence or adulthood.

Autism involves differences in social communication and restricted or repetitive patterns of behavior, interests, or sensory responses.

Social communication differences may include:

These differences do not mean that an autistic person lacks feelings or empathy. Some experience other people's emotions strongly but have difficulty recognizing or responding to them in expected ways.

Restricted or repetitive features may include:

Sensory differences are also common. A person may be unusually sensitive or less responsive to sounds, lights, smells, textures, temperatures, pain, or body sensations. Crowds, fluorescent lighting, certain clothing, or background noise may become overwhelming. Others may seek particular movements, textures, sounds, or pressure because they are calming.

Some people hide or compensate for autistic traits, often called masking or camouflaging. They may copy social behavior, rehearse conversations, or force eye contact. Masking can require significant effort and contribute to exhaustion or distress.

No single trait confirms autism. Similar experiences can occur with attention-deficit/hyperactivity disorder (ADHD), anxiety, language disorders, learning differences, trauma, or other conditions.

- Difficulty interpreting facial expressions, tone, implied meanings, humor, or unwritten social rules
- A direct, formal, detailed, or literal communication style

- Difficulty starting, maintaining, or ending conversations
- Finding group conversations hard to follow
- Preferring a few close relationships or time alone
- Needing more time to process spoken information
- Using less eye contact or finding it uncomfortable
- Strong, focused interests
- A preference for routine, predictability, and clear expectations
- Distress or fatigue when plans change unexpectedly
- Repeated movements or sounds, sometimes called "stimming," that may help regulate emotions or sensory input
- Repeating words or phrases
- Organizing objects or information in specific ways
- Close attention to details, patterns, systems, or particular topics

Causes and risk factors

Autism does not have one known cause. Research indicates that it usually results from a complex combination of genetic and developmental factors affecting brain development.

Autism can run in families, but inheritance is not simple. Many genetic variations may contribute, and someone can be autistic without another known autistic family member. Certain genetic or chromosomal conditions are associated with autism, although most autistic people do not have an identified genetic syndrome.

Factors associated with a greater likelihood of autism include having an autistic sibling and certain events or exposures before or around birth. These associations do not mean that a parent caused autism. A specific cause cannot usually be identified.

Vaccines do not cause autism. Large studies have found no link between autism and the measles, mumps, and rubella vaccine or vaccines containing thimerosal. Parenting style, emotional warmth, diet, and ordinary childhood experiences also do not cause autism.

Diagnosis

There is no blood test, brain scan, or online questionnaire that can diagnose autism by itself.

Diagnosis is based on developmental history, current traits, direct observation, and effects on everyday life.

A childhood assessment may involve a pediatrician, psychologist, speech-language therapist, occupational therapist, or another clinician with autism expertise. The team may ask about early language, play, social interaction, routines, sensory responses, behavior, learning, and family history. Information from caregivers and school can help.

Developmental screening is not a diagnosis. It may indicate that a child needs fuller evaluation. A child can also be referred whenever a caregiver, teacher, or health professional has concerns.

Adult assessment examines current experiences and childhood development. School reports or information from someone who knew the person as a child may help, but assessment may still be possible without them.

Clinicians consider other explanations and co-occurring conditions, including ADHD, anxiety, depression, obsessive-compulsive disorder, language or learning disorders, intellectual disability, sleep problems, epilepsy, and hearing or vision difficulties.

A thorough evaluation should identify strengths and support needs. It may assess communication, learning, daily living skills, sensory needs, mental health, education, employment, and relationships.

Treatment and management

Autism itself does not require a cure. Management focuses on reducing barriers, building useful skills, supporting well-being, and treating co-occurring medical or mental health concerns. Plans should be individualized with the autistic person and, when appropriate, their family or caregivers.

Helpful communication adjustments may include written instructions, predictable schedules, extra processing time, quieter waiting areas, direct language, and advance notice of changes.

Children may benefit from speech and language therapy, occupational therapy, educational support, and structured approaches that build communication, play, daily living, or emotional regulation skills. Goals should be meaningful to the child rather than focused on making them appear non-autistic. Caregivers should be included without blame.

School accommodations may include consistent routines, sensory breaks, visual schedules, written directions, extra time, transition support, or a quieter workspace. Available services depend on local systems and individual needs.

Adults may benefit from workplace adjustments, organizational coaching, independent-living support, communication therapy, sensory adaptations, or help with education and relationships. Counseling may address anxiety, low mood, identity, burnout, or masking. Therapy may need clear language, concrete examples, predictable sessions, and sensory accommodations.

No medicine treats the core features of autism. Medication may be considered for a co-occurring condition or symptom causing significant difficulty, such as ADHD, anxiety, depression, severe irritability, or sleep problems. Decisions should include careful assessment, non-medication options, expected benefits, side effects, and monitoring.

Restrictive diets, supplements, chelation, and treatments marketed as removing "toxins" have not been shown to cure autism. Some can cause harm or interact with medicines. Discuss dietary changes with a qualified clinician, particularly if eating is already limited.

Complications

Complications may result from co-occurring conditions, unmet support needs, or environments that do not accommodate communication and sensory differences.

Possible concerns include anxiety, depression, ADHD, sleep disorders, epilepsy, digestive symptoms, feeding difficulties, and problems recognizing or describing pain and illness. A limited range of accepted foods may affect nutrition and should be assessed rather than assumed to be behavioral.

Social exclusion, bullying, unemployment, relationship difficulties, and barriers to health care can affect quality of life. Repeated sensory or social overload may contribute to shutdowns, intense distress, or autistic burnout-prolonged exhaustion and reduced ability to manage daily tasks after sustained demands or masking.

Mental health symptoms should not automatically be attributed to autism. New withdrawal, loss of skills, sleep changes, reduced eating, or major behavior changes need clinical assessment.

Thoughts of self-harm or suicide require urgent attention. Contact a local crisis service or health professional promptly. If there is immediate danger, an attempt has occurred, or the person cannot stay safe, call emergency services or go to the nearest emergency department.

Follow up

Follow-up should reflect the person's age, goals, health, and support needs. Autism does not

usually require repeated testing to confirm it is still present. Reviews identify changing needs and check whether supports remain useful.

Routine follow-up may cover:

Autistic people should receive the same preventive health care as anyone else. Appointment adjustments may make examinations, vaccinations, dental visits, and screening tests easier.

Arrange a routine appointment if current supports no longer fit, daily tasks are becoming harder, or concerns develop gradually. Seek prompt medical review for loss of established skills, persistent inability to eat or drink enough, severe sleep disruption, suspected seizures, or a major change in mental state.

- Communication, learning, work, relationships, and independent living
- Sensory needs and accommodations
- Sleep, eating, physical activity, dental care, and general health
- Anxiety, mood, attention, behavior changes, and caregiver well-being
- Medicine effectiveness and side effects
- School transitions, adulthood, employment, or living arrangements

Sources

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders . 5th ed., text revision. American Psychiatric Association Publishing; 2022.
- World Health Organization. International Classification of Diseases 11th Revision: 6A02 Autism Spectrum Disorder . WHO; 2024 release.
- Hyman SL, Levy SE, Myers SM; Council on Children With Disabilities, Section on Developmental and Behavioral Pediatrics. Identification, evaluation, and management of children with autism spectrum disorder. Pediatrics . 2020;145(1):e20193447.
- National Institute for Health and Care Excellence. Autism Spectrum Disorder in Under 19s: Recognition, Referral and Diagnosis . Clinical guideline CG128. NICE; published 2011, updated 2017.
- National Institute for Health and Care Excellence. Autism Spectrum Disorder in Adults: Diagnosis and Management . Clinical guideline CG142. NICE; published 2012, updated 2021.
- National Institute for Health and Care Excellence. Autism Spectrum Disorder in Under 19s: Support and Management . Clinical guideline CG170. NICE; published 2013, updated 2021.

- Centers for Disease Control and Prevention. Clinical Testing and Diagnosis for Autism Spectrum Disorder . National Center on Birth Defects and Developmental Disabilities; reviewed 2024.