CONDITION

Baking powder overdose

Learn how swallowing too much baking powder can affect the stomach, body salts, and acid-base balance, and when to contact Poison Control or seek emergency care.

WHAT IT IS
Harmful effects from swallowing excessive baking powder

BODY SYSTEM
Digestive system and body salt balance

MAIN SYMPTOMS
Nausea, vomiting, abdominal pain or bloating, unusual weakness

WHAT CAUSES IT
Swallowing too much baking powder

HOW IT IS DIAGNOSED
Exposure history, electrolyte tests, blood gas test

MAIN TREATMENT
Supportive care

CAN IT BE CURED?
Yes - supportive treatment can correct abnormalities

IS IT AN EMERGENCY?
Sometimes - seizure, collapse, breathing difficulty or unresponsiveness

LONG-TERM OUTLOOK
Usually no lasting harm after small accidental exposures

Overview

Baking powder is a leavening product that helps baked goods rise. It usually contains sodium bicarbonate, one or more powdered acids, and a starch that keeps the ingredients dry. Formulas vary, so the exact ingredients should be checked on the package.

A baking powder overdose happens when someone swallows enough baking powder to cause harmful effects. This is different from eating food prepared with the usual amount of baking powder. Normal cooking use is not considered an overdose.

Baking powder is also different from baking soda, which is pure sodium bicarbonate. However, baking powder contains sodium bicarbonate and can cause some of the same problems when swallowed in a large amount. Its other ingredients may also affect symptoms.

When baking powder reacts with liquid and stomach acid, it produces carbon dioxide gas. This can cause bloating, nausea, vomiting, or abdominal pain. A large exposure may also disturb the body's electrolytes, which are minerals such as sodium and potassium that help control muscles, nerves, and heart rhythm.

Too much bicarbonate may make the blood overly alkaline. This condition is called metabolic alkalosis . Severe alkalosis can affect breathing, heart rhythm, muscle function, and alertness. Vomiting and diarrhea may add to fluid and electrolyte problems.

If someone may have swallowed more than a small taste, contact Poison Control rather than waiting for symptoms. In the United States, call 1-800-222-1222 or use the official online Poison Control service. The service is free and confidential. Outside the United States, contact the local poison information center.

Call emergency services immediately if the person collapses, has a seizure, has trouble breathing, or cannot be awakened.

Symptoms

Symptoms depend on the amount swallowed, the product's ingredients, the person's age and size, and whether the exposure happened once or repeatedly. Health conditions affecting the kidneys, heart, or fluid balance may also change the effects.

Early digestive symptoms may include:

A larger exposure may cause signs of fluid, sodium, potassium, or acid-base disturbances. These can include unusual weakness, muscle cramps, twitching, confusion, irritability, or marked sleepiness. Some people may urinate more than usual.

Severe poisoning may affect the brain, breathing, or heart. Possible signs include an irregular heartbeat, severe confusion, seizures, slowed or difficult breathing, or loss of consciousness. These require emergency care.

Symptoms may not show how serious an exposure is. A person can have important electrolyte or acid-base changes before the signs become obvious. This is one reason to obtain poison-center advice promptly after a significant or uncertain ingestion.

Powder in the mouth or throat may also cause coughing or irritation. If someone inhales a large amount of dust and develops ongoing coughing, wheezing, or breathing difficulty, they need medical assessment.

- Thirst or a dry mouth
- Nausea or vomiting

- Diarrhea
- Abdominal pain, fullness, or bloating
- Increased burping from carbon dioxide gas

Causes and risk factors

Most exposures happen when baking powder is swallowed directly rather than used in a recipe. A child may taste it after finding an open container. An adult may mistake it for another powder or intentionally consume it as a home remedy.

Repeated spoonfuls can be harmful even if no single dose seems very large. This may occur when someone uses a sodium bicarbonate-containing product for indigestion or another symptom without realizing how much sodium and bicarbonate they are taking. Baking powder should not be used as a substitute for an antacid or other medicine.

Risk may be higher when:

The product formula matters. Baking powders may contain ingredients such as monocalcium phosphate, sodium acid pyrophosphate, sodium aluminum sulfate, or cream of tartar. Clinicians and poison specialists use the package label to assess the likely effects.

Do not assume that "aluminum-free," "low sodium," or another label claim makes a large ingestion safe. It may change the ingredients, but it does not remove the risk from swallowing an excessive amount.

- The amount swallowed is large or unknown
- The product was consumed repeatedly
- The person is a young child or an older adult
- Kidney disease limits the body's ability to remove sodium or regulate electrolytes
- Heart disease makes excess sodium or fluid harder to tolerate
- The person is dehydrated from vomiting, diarrhea, fever, or poor fluid intake
- Other sodium bicarbonate products, antacids, diuretics like furosemide or hydrochlorothiazide , or medicines affecting electrolytes are also being used
- Confusion, impaired judgment, an eating disorder, or pica leads to repeated nonfood ingestion

Diagnosis

There is no single test that confirms every baking powder overdose. Assessment begins with the product name, ingredient list, estimated amount, time of ingestion, and symptoms. The person's age, weight, health conditions, and medicines are also important.

Keep the container or take a clear photograph of its label. Do not delay calling Poison Control while trying to calculate an exact amount. An estimate such as "one spoonful," "part of the container," or "an unknown amount" can still help.

In a clinic or emergency department, the medical team may check breathing, heart rate, blood pressure, temperature, alertness, hydration, and the abdomen. An electrocardiogram, or ECG, may be used to look for heart-rhythm changes.

Blood tests may measure sodium, potassium, chloride, bicarbonate, glucose, and kidney function. A blood gas test can show whether the blood is too alkaline and how breathing is responding. Calcium, magnesium, or phosphate may be checked when symptoms, test results, or product ingredients make these measurements relevant.

Testing is based on the exposure. A small accidental taste in a person without symptoms may not require hospital tests after poison-center review. A large, repeated, or uncertain ingestion is more likely to need medical evaluation and monitoring.

Treatment and management

Do not make the person vomit. Vomiting can lead to choking or allow material to enter the lungs. Do not try to neutralize baking powder with vinegar, lemon juice, or another acid. Mixing acids and bicarbonate can produce more gas and may worsen stomach pressure or vomiting.

Remove any remaining powder from the mouth. If the person is awake and able to swallow normally, follow the instructions given by Poison Control. Do not give food, drinks, medicines, or home remedies to a drowsy person or anyone who is vomiting repeatedly.

Contact Poison Control promptly if more than a small taste may have been swallowed, the amount is unknown, symptoms are present, or repeated use has occurred. Specialists can determine whether observation at home is reasonable or whether the person needs urgent medical evaluation.

Have the following information ready when possible:

There is no specific antidote for baking powder overdose. Treatment is supportive, meaning it focuses on breathing, circulation, hydration, symptoms, and abnormal laboratory results.

In a medical setting, clinicians may monitor heart rhythm and repeat blood tests. Intravenous

fluids may be used when appropriate. Electrolyte abnormalities are corrected carefully because changing sodium, potassium, or acid-base balance too quickly can cause additional problems. Treatment is adjusted for kidney function, heart health, and the specific laboratory findings.

Medicines may be given for nausea, seizures, or other complications when needed. People with severe metabolic alkalosis, major electrolyte disturbances, kidney failure, or unstable heart rhythms may require intensive monitoring. Dialysis is rarely considered in severe cases when the body cannot correct the abnormalities or when kidney function is seriously impaired.

Activated charcoal is not a home treatment and should not be given unless a poison specialist or clinician specifically recommends it. Hospital decontamination procedures are not routinely appropriate for every ingestion and can carry their own risks.

- The person's age and approximate weight
- The product name and ingredient list
- The estimated amount and time swallowed
- Current symptoms
- Medical conditions and regular medicines
- Any first aid already given

Complications

Most small accidental exposures do not lead to lasting harm, especially when they are assessed early. The chance of complications rises with a large amount, repeated ingestion, delayed care, or underlying kidney or heart disease.

Possible complications include dehydration from vomiting or diarrhea and abnormal sodium, potassium, or chloride levels. High sodium or low potassium can affect the brain, muscles, and heart.

Severe metabolic alkalosis may cause muscle spasms, confusion, abnormal heart rhythms, seizures, or reduced breathing. Excess sodium and fluid may worsen swelling, blood pressure, or heart failure in someone who is vulnerable.

Gas formation can cause marked stomach swelling. Serious injury from extreme stomach distention is uncommon, but severe or worsening abdominal pain, a hard or swollen abdomen, repeated vomiting, vomiting blood, or fainting requires emergency assessment.

Vomiting may also lead to aspiration, which means stomach contents enter the lungs. New

breathing difficulty, choking, chest discomfort, or persistent coughing after vomiting needs urgent medical attention.

Follow up

Follow the instructions from Poison Control or the treating medical team. A person who remains well after a minor exposure may only need observation for a specified period. Do not resume intentional use of baking powder as a home remedy.

After a medically significant exposure, follow-up may include repeat electrolyte or kidney-function tests. This is more likely after severe symptoms, abnormal hospital results, repeated ingestion, or in people with kidney or heart disease.

Arrange routine medical follow-up if the ingestion was related to ongoing indigestion, attempts to change body pH, pica, an eating disorder, confusion, or difficulty using medicines safely. The goal is to address the reason for the exposure and identify safer treatment or support.

Seek urgent medical advice for continuing vomiting, increasing weakness, muscle cramps, unusual sleepiness, confusion, reduced urination, or worsening abdominal symptoms.

Call emergency services for a seizure, collapse, severe breathing difficulty, inability to wake the person, or signs of a serious heart-rhythm problem.

Sources

- MedlinePlus Medical Encyclopedia. Baking powder overdose. U.S. National Library of Medicine; accessed March 8, 2025.
- Health Resources and Services Administration. Poison Help. U.S. Department of Health and Human Services; accessed March 8, 2025.
- America's Poison Centers. Get Poison Help. National poison-center guidance; accessed March 8, 2025.
- Al-Abri SA, Kearney T. Baking soda misuse as a home remedy: case experience of the California Poison Control System. Journal of Clinical Pharmacy and Therapeutics. 2014;39(1):73-77.
- Thomas SHL, Stone CK. Acute toxicity from baking soda ingestion. American Journal of Emergency Medicine. 1994;12(1):57-59.

- Merck Manual Consumer Version. Metabolic alkalosis. Merck & Co.; accessed March 8, 2025.