CONDITION

Black lung disease

Learn how coal mine dust can damage the lungs, what symptoms may occur, how black lung disease is diagnosed, and how exposure control and supportive care can help.

WHAT IT IS
Coal dust-related lung disease

BODY SYSTEM
Lungs and breathing

MAIN SYMPTOMS
Long-lasting cough, shortness of breath, chest tightness, wheezing

WHAT CAUSES IT
Inhaled coal mine dust

HOW IT IS DIAGNOSED
Work history and chest X-ray or CT

MAIN TREATMENT
Exposure control and supportive care

CAN IT BE CURED?
No, established scarring is permanent

IS IT AN EMERGENCY?
Sometimes, with severe trouble breathing

LONG-TERM OUTLOOK
May remain stable or progress to respiratory failure

Overview

Black lung disease is a group of lung problems caused by breathing coal mine dust. The condition most often called black lung is coal workers' pneumoconiosis , or CWP. Pneumoconiosis means lung disease caused by inhaled mineral dust.

Coal mine dust can settle deep in the lungs. The immune system reacts to the particles, which may lead to inflammation and permanent scarring. This damage makes it harder for the lungs to move oxygen into the blood. Coal mine dust can also contribute to chronic bronchitis and emphysema, two forms of chronic obstructive pulmonary disease, or COPD.

CWP is often divided into two forms. Simple CWP causes small areas of abnormal tissue in the lungs. It may cause few or no symptoms. Complicated CWP , also called progressive massive fibrosis or PMF, causes larger areas of dense scarring. PMF can seriously affect breathing.

The term "black lung" comes from the dark coal dust found in affected lung tissue. It does not mean that a person's lungs will look black on an imaging test.

Black lung disease matters because lung damage may be permanent and can progress even after dust exposure ends. Early detection can help a worker reduce further exposure, receive appropriate monitoring, and address other conditions that affect breathing.

Symptoms

Early CWP may not cause noticeable symptoms. Some people first learn that they have lung changes during workplace health screening.

When symptoms occur, they may include:

Symptoms do not always match the amount of change seen on a chest X-ray. A person with simple CWP may have few symptoms, while another person with similar imaging findings may have significant breathing difficulty because of COPD, asthma, heart disease, or another condition.

PMF is more likely to cause increasing shortness of breath, a persistent cough, low blood oxygen, and difficulty completing usual activities. These symptoms can resemble those of other lung or heart conditions, so testing is needed to determine the cause.

Arrange a routine medical appointment if a cough, wheezing, or breathlessness persists or gradually worsens. Contact a clinician promptly for a sudden change in breathing, fever with worsening cough, or blood in the mucus.

Call emergency services for severe trouble breathing, blue or gray lips, new confusion, fainting, or chest pressure that does not quickly improve.

- A long-lasting cough
- Shortness of breath, especially during activity
- Reduced exercise tolerance
- Chest tightness
- Mucus production
- Wheezing

Causes and risk factors

Black lung disease develops after coal mine dust is inhaled into the lungs. This dust contains

coal particles and may also contain silica and other minerals. Crystalline silica is a particularly harmful mineral found in rock. Miners may encounter high silica exposure while cutting through rock around coal seams.

The body cannot easily remove particles that reach the deepest parts of the lungs. Immune cells attempt to contain them, but the ongoing response can damage nearby tissue and create scars.

Risk depends mainly on the total amount of dust inhaled over time. Important factors include:

Both underground and surface miners can be exposed. Workers outside mining may also encounter coal or silica dust, but their lung conditions may be classified differently depending on the exposure.

Smoking does not cause CWP. However, smoking can cause COPD and lung cancer and can further reduce breathing capacity in someone exposed to coal dust. Coal mine dust itself can also contribute to COPD, even in a person who has never smoked.

Black lung is not contagious. It cannot be passed from one person to another. It is not caused by a short-term infection.

- The concentration of respirable dust in the workplace
- The number of years spent in dusty work
- Work involving drilling, blasting, cutting, or handling rock
- Exposure to dust with a high silica content
- The effectiveness of ventilation, water sprays, dust monitoring, and other controls
- Continued exposure after early lung changes appear

Diagnosis

Diagnosis begins with a detailed work history. A clinician may ask about job titles, specific tasks, years of exposure, dust-control practices, respirator use, and whether similar illness has occurred among coworkers. Past jobs remain relevant because symptoms and imaging changes can appear after mining work has ended.

A medical assessment may include:

A lung biopsy is not usually needed when the exposure history and imaging are typical. It may be considered if the diagnosis remains unclear or if doctors need to rule out another disease.

An abnormal X-ray does not show exactly how much breathing is affected. Likewise, normal spirometry does not necessarily rule out early CWP. Clinicians consider the occupational

history, symptoms, imaging, and lung function together.

In the United States, eligible coal miners can receive confidential health screening through the National Institute for Occupational Safety and Health's Coal Workers' Health Surveillance Program. Medical screening and an evaluation for disability or compensation are separate processes.

- Chest X-ray: X-rays can show the small rounded shadows associated with simple CWP or the larger areas seen in PMF. For occupational surveillance, images may be classified using a standardized international system. A NIOSH-certified B Reader is a physician trained to apply this system.
- Chest CT scan: CT produces more detailed images and may identify abnormalities not clearly visible on an X-ray. It may be used when findings are uncertain or another condition is suspected.
- Spirometry: This breathing test measures how much air a person can exhale and how quickly. It can identify airflow obstruction or reduced lung capacity, but it cannot diagnose CWP by itself.
- Oxygen measurement: A pulse oximeter estimates blood oxygen through a sensor placed on a finger. Some people also need testing during walking or exercise.
- Other tests: Blood tests, sputum testing, heart tests, or additional imaging may be needed to look for infection, tuberculosis, cancer, heart disease, or another explanation for the findings.

Treatment and management

There is no treatment that removes coal dust or reverses established lung scarring. Management focuses on preventing additional exposure, easing symptoms, maintaining activity, and treating related health problems.

Reducing dust exposure is central to care. Workplaces should use engineering controls such as ventilation, water sprays, enclosed equipment, and dust monitoring. Respiratory protection may be part of a broader safety program, but it should not replace effective workplace dust controls.

A miner with signs of dust-related lung disease should discuss exposure with an occupational medicine clinician and the workplace health or safety program. Under certain US federal protections, some miners may be eligible to work in a lower-dust area without loss of pay. A union representative, occupational health clinic, NIOSH, or the Mine Safety and Health Administration can provide information about applicable programs and rights.

Treatment may include:

Corticosteroids and medicines that suppress the immune system are not established treatments for CWP itself. Antibiotics like amoxicillin or doxycycline only treat bacterial infections; they do not treat black lung disease.

Practical steps such as pacing activities, taking rest breaks, staying physically active within safe limits, and avoiding secondhand smoke or additional dust may also help with daily symptoms.

- Inhaled medicines: Bronchodilators like albuterol or salmeterol relax the muscles around the airways and may help when asthma-like narrowing or COPD is also present. These medicines do not treat lung scarring itself.
- Pulmonary rehabilitation: This supervised program combines exercise, breathing strategies, and education. It may improve stamina and help people manage breathlessness.
- Oxygen therapy: Supplemental oxygen may be prescribed when testing shows that blood oxygen is too low. Oxygen should be used according to the prescribed flow and safety instructions.
- Vaccination and infection care: Staying current with clinician-recommended respiratory vaccines can reduce the risk of some serious infections. New infections should be assessed and treated appropriately.
- Smoking cessation: For people who smoke, stopping can slow additional lung damage and lower the risk of cancer, heart disease, and COPD complications.
- Advanced care: Severe disease may require specialist management. Lung transplantation may be considered for a small number of people with advanced respiratory failure.

Complications

Some people with simple CWP remain stable, while others develop worsening disease. Progression may occur despite leaving mining work because dust already in the lungs can continue to trigger inflammation and scarring.

Possible complications include:

Exposure to silica is also linked to silicosis, tuberculosis, and lung cancer. A person's individual risk depends on the type and amount of exposure and other health factors. Coughing blood, unexplained weight loss, fever, or a major change in symptoms needs medical assessment rather than being assumed to be part of black lung disease.

- Progressive massive fibrosis: Large areas of scarring distort lung tissue and can severely limit airflow and oxygen transfer.
- COPD: Chronic bronchitis and emphysema can cause cough, mucus, wheezing, and persistent airflow obstruction.
- Low blood oxygen: Advanced lung damage may prevent enough oxygen from reaching the bloodstream.
- Pulmonary hypertension: High pressure can develop in the blood vessels that carry blood through the lungs.
- Right-sided heart strain: Over time, pulmonary hypertension may weaken the right side of the heart, a condition called cor pulmonale.
- Respiratory failure: Severely damaged lungs may no longer provide enough oxygen or remove enough carbon dioxide.
- Infections: Damaged lungs may have less reserve during pneumonia or another respiratory infection.

Follow up

Follow-up should reflect the person's symptoms, test results, exposure status, and other medical conditions. A primary care clinician may coordinate care with a pulmonologist, who specializes in lung disease, or an occupational medicine clinician.

Routine visits may review breathing symptoms, work exposure, smoking status, physical activity, inhaler technique, vaccination needs, and oxygen use. Spirometry, imaging, or exercise oxygen testing may be repeated when clinically appropriate. Imaging is not always needed at every visit.

Current and former miners should keep copies of job histories, workplace exposure records, chest images, breathing tests, and medical reports. These records can help clinicians recognize changes over time. Eligible miners may also choose to participate in federal surveillance screening even when they feel well.

Seek earlier medical review if breathing becomes noticeably worse, daily activities become harder, oxygen readings are repeatedly below the range set by a clinician, or a new cough or fever develops. Emergency symptoms include severe breathlessness at rest, blue or gray coloring, confusion, fainting, or persistent chest pressure.

Sources

- Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Coal Workers' Health Surveillance Program . CDC/NIOSH.
- National Institute for Occupational Safety and Health. Coal Mine Dust Exposures and Associated Health Outcomes: A Review of Information Published Since 1995 . Current Intelligence Bulletin 64. DHHS (NIOSH) Publication No. 2011-172; 2011.
- National Institute for Occupational Safety and Health. Criteria for a Recommended Standard: Occupational Exposure to Respirable Coal Mine Dust . DHHS (NIOSH) Publication No. 95-106; 1995.
- Hall NB, Blackley DJ, Halldin CN, Laney AS. Current review of pneumoconiosis among US coal miners. Current Environmental Health Reports . 2019;6(3):137-147.
- Laney AS, Weissman DN. Respiratory diseases caused by coal mine dust. Journal of Occupational and Environmental Medicine . 2014;56(10 Suppl):S18-S22.
- Electronic Code of Federal Regulations. Title 42, Part 37: Specifications for Medical Examinations of Coal Miners . US Government Publishing Office.