CONDITION

Developmental differences of the female genital tract

Developmental differences of the female genital tract are structural differences present from birth that may affect the vagina, cervix, uterus, or fallopian tubes.

WHAT IT IS
Congenital differences in reproductive tract structure

BODY SYSTEM
Reproductive and urinary systems

MAIN SYMPTOMS
Absent periods, pelvic pain, fertility problems, pregnancy complications

WHAT CAUSES IT
Early developmental changes, sometimes genetic

HOW IT IS DIAGNOSED
Examination, ultrasound, MRI, sometimes keyhole surgery

MAIN TREATMENT
Monitoring, symptom relief, dilation, or corrective surgery

CAN IT BE CURED?
Sometimes - selected structural differences can be corrected

IS IT AN EMERGENCY?
Sometimes - severe pain or inability to pass urine

LONG-TERM OUTLOOK
Most people lead full lives with appropriate care

Overview

Developmental differences of the female genital tract are structural differences that form before birth. They can affect the vagina, cervix, uterus, or fallopian tubes. Sometimes the ovaries or urinary system are also involved.

These conditions are also called congenital reproductive tract anomalies or Müllerian anomalies. The Müllerian ducts are early embryonic structures that normally develop into the fallopian tubes, uterus, cervix, and upper vagina.

Possible patterns include:

Some differences cause no symptoms and are found during pregnancy imaging, fertility

assessment, or surgery for another reason. Others block menstrual flow, affect sexual function, or increase pregnancy risks.

A developmental difference does not define your sex, gender, sexuality, or ability to have relationships. Its effect on fertility depends on the structures involved.

- A uterus divided by a wall, called a septate uterus
- A heart-shaped or two-sided uterus, such as a bicornuate uterus
- One-sided uterine development, called a unicornuate uterus
- Two uteruses, sometimes with two cervixes or a divided vagina
- An underdeveloped or absent uterus, cervix, or upper vagina
- A vaginal wall or hymen that blocks menstrual flow

Symptoms

Symptoms often appear around puberty, although some conditions are not found until adulthood.

Possible signs include:

Monthly pain without bleeding may mean blood is trapped behind a blockage in the hymen, vagina, or cervix. Pressure can increase as blood collects in the vagina or uterus.

People with an absent or underdeveloped uterus may develop breasts and pubic hair normally but not have periods because the ovaries can still produce reproductive hormones. Uterine shape differences usually do not affect hormone levels or puberty, and some cause no symptoms.

- Periods not starting despite other signs of puberty
- Monthly or repeated pelvic pain without bleeding
- Unusual menstrual bleeding
- Difficulty inserting a tampon
- Pain or difficulty with vaginal penetration
- Pressure or a lump near the vaginal opening
- Difficulty emptying the bladder or repeated urinary infections
- Difficulty becoming pregnant
- Repeated miscarriage, early birth, or unusual fetal position

Causes and risk factors

These differences arise while the reproductive tract is forming early in pregnancy. Structures may not develop fully, join together, or open as expected.

Most have no single identifiable cause. Genetic changes can contribute, particularly when other organs are affected, and a difference may rarely be part of a wider genetic syndrome.

Because the reproductive and urinary systems develop closely together, a uterine or vaginal difference may occur with a kidney or urinary tract difference. Examples include having one kidney or a kidney in an unusual position. Some conditions are also associated with differences involving the spine, ribs, hearing, or heart.

Exposure before birth to diethylstilbestrol, or DES, can cause changes in the uterus and cervix. This synthetic estrogen was once prescribed during pregnancy but is no longer used for that purpose.

These conditions are not caused by exercise, sexual activity, contraception, or something you did. Usually, nothing a pregnant parent did could have prevented them.

Diagnosis

Diagnosis begins with symptoms and medical history. A clinician may ask about puberty, periods, pain, urinary symptoms, sexual activity, pregnancies, and previous operations.

An external genital examination may identify a blocked hymen or vaginal opening. An internal examination is not always needed, especially in younger people or when imaging provides enough information. Examinations should be explained, require your consent, and stop if you ask.

Tests may include:

Blood tests may check hormones or chromosomes when periods have not begun and the cause is unclear.

Hysteroscopy uses a thin camera passed through the cervix to view the uterus. Laparoscopy uses a camera inserted through small abdominal cuts. These procedures are generally reserved for treatment planning or questions that imaging has not answered.

- Ultrasound to examine the uterus, ovaries, kidneys, and trapped fluid or blood. It may be performed over the abdomen or, when appropriate and agreed, through the vagina.

- Three-dimensional ultrasound to show the internal and outer shape of the uterus in more detail.
- MRI to provide detailed images of complex vaginal, cervical, or uterine anatomy.
- Kidney imaging because urinary tract differences can occur alongside genital tract differences.

Treatment and management

Treatment depends on symptoms, anatomy, age, and personal goals, including menstrual flow, sexual function, fertility, and pregnancy. A difference found by chance may need only an explanation and follow-up.

A blockage trapping menstrual blood generally requires treatment to create a safe drainage route. Surgery depends on whether the blockage affects the hymen, vagina, or cervix. Complex procedures should be planned by a team experienced in congenital reproductive anatomy.

A short or absent vagina does not require treatment because of its appearance. If comfortable vaginal penetration is wanted, options include gradual dilation or surgery. Dilation uses smooth devices to gently lengthen vaginal tissue over time and should be taught with privacy, support, and attention to comfort. Surgery may be considered if dilation is unsuitable, unsuccessful, or not preferred after informed discussion.

Most uterine shape differences do not automatically require surgery. An operation may be considered for selected problems, such as a uterine septum associated with adverse reproductive outcomes or a separate uterine remnant causing pain or pregnancy risk.

Other care may include:

Hormone treatment cannot create a missing uterus or correct a blockage. It may help if a separate hormonal or ovarian condition is present.

- Pain relief
- Treatment for endometriosis or infection
- Fertility assessment or treatment
- Specialist pregnancy monitoring
- Pelvic floor physiotherapy
- Psychosexual counselling or psychological support

Complications

Blocked menstrual flow can cause blood to collect in the vagina, uterus, or fallopian tubes. Over time, this may contribute to persistent pain, infection, or endometriosis.

Some uterine differences increase the chance of miscarriage, early birth, restricted fetal growth, or a baby lying sideways or bottom-first. Risk varies by anatomy.

A pregnancy in a small, separate uterine horn can rupture and cause internal bleeding. This is uncommon but serious. The uterus should be clearly mapped before or early in pregnancy when this anatomy is known or suspected.

Fertility can be affected if the uterus, cervix, or fallopian tubes are absent, blocked, or significantly underdeveloped. However, many people with uterine differences conceive without fertility treatment. Someone without a functioning uterus cannot carry a pregnancy, although their ovaries may still produce eggs.

Diagnosis may also affect body image, sexual confidence, relationships, and feelings about fertility. These emotional effects deserve support alongside physical care.

Prevention

Most developmental differences of the female genital tract cannot be prevented. They form early before birth, usually without a known cause.

Diet, exercise, sexual activity, and routine behaviour cannot change an existing structural difference. Treatment focuses on symptoms, complications, and personal or reproductive goals.

If a genetic condition is identified, genetic counselling can explain inheritance and reproductive implications. The condition is not anyone's fault.

Living with it and what to expect

The anatomy usually remains stable, but its effects may change through life. A blockage may become apparent when periods begin, while questions about vaginal function, fertility, or pregnancy may arise later.

Some people need one procedure and no long-term treatment. Others with complex anatomy need follow-up across adolescence and adulthood. Care may involve gynecology, urology, fertility services, maternal-fetal medicine, pelvic floor therapy, or mental health support.

Follow-up can review pain, bleeding, urinary symptoms, sexual comfort, and reproductive

plans. Before trying to become pregnant, a specialist review can confirm uterine anatomy, check whether kidney imaging is needed, and identify whether specialist pregnancy care is recommended.

A vaginal or uterine difference does not prevent intimacy or a satisfying sex life. Treatment should reflect your wishes rather than pressure to make your body look or function in a particular way.

Fertility outcomes vary. Some people conceive and carry a pregnancy without difficulty. Others may need fertility treatment, pregnancy monitoring, or information about gestational surrogacy, adoption, or choosing not to become a parent.

When to get medical help

Arrange a routine medical appointment if:

Seek urgent medical care for severe or rapidly worsening pelvic pain, especially with fever, vomiting, abdominal swelling, or difficulty passing urine.

Get emergency help for sudden severe abdominal pain with faintness, collapse, shoulder pain, or heavy bleeding. If you could be pregnant, these symptoms may indicate internal bleeding or an ectopic pregnancy and require immediate assessment.

- Periods have not started by age 15 or within about three years of breast development
- You have repeated monthly pelvic pain without bleeding
- Tampon insertion or vaginal penetration is unexpectedly difficult or painful
- You have ongoing pelvic pain, unusual bleeding, or fertility concerns
- A scan suggests a difference in the uterus, cervix, or vagina

Common questions

Can I still have periods with a developmental difference?

Often, yes. Periods may occur normally if there is a functioning uterus and an open route through the cervix and vagina. They may be absent if the uterus is absent or underdeveloped. A blockage can cause monthly pain without visible bleeding.

Can I get pregnant?

Many people with uterine shape differences can become pregnant. Fertility may be reduced if the uterus, cervix, or fallopian tubes are absent or blocked. Individual anatomy and ovarian

function must be assessed before likely options can be explained.

Does an unusual uterus always need surgery?

No. Many differences cause no symptoms and need no treatment. Surgery may be considered for obstruction, significant pain, a specific pregnancy risk, or a reproductive problem likely to improve with correction.

Will sex be painful?

Not necessarily. A short, narrow, divided, or blocked vagina may cause pain, but treatment and pelvic floor support can help. Sex should not be forced through pain. A specialist can discuss options suited to your goals.

Is this the same as being intersex?

Not always. Some differences occur in people with otherwise typical female chromosomes, hormones, ovaries, and external genitalia. Others may be part of a variation in sex development. Testing depends on the anatomy and other findings.

Could I pass this on to a child?

Most cases have no clear inheritance pattern. Some are linked to genetic conditions and may run in families. Genetic counselling may help when relatives are affected or there are kidney, skeletal, or identified genetic differences.

Sources

- American College of Obstetricians and Gynecologists. Müllerian Agenesis: Diagnosis, Management, and Treatment . Committee Opinion No. 728, 2018.
- American College of Obstetricians and Gynecologists. Management of Acute Obstructive Uterovaginal Anomalies . Committee Opinion No. 779, 2019.
- American College of Obstetricians and Gynecologists. Menstruation in Girls and Adolescents: Using the Menstrual Cycle as a Vital Sign . Committee Opinion No. 651, 2015.
- American Society for Reproductive Medicine. ASRM Müllerian Anomalies Classification 2021 . Fertility and Sterility, 2021.
- European Society of Human Reproduction and Embryology and European Society for Gynaecological Endoscopy. The ESHRE/ESGE Consensus on the Classification of Female Genital Tract Congenital Anomalies . Human Reproduction, 2013.
- National Health Service. Mayer-Rokitansky-Küster-Hauser syndrome . NHS health information.