CONDITION

Dieffenbachia poisoning

Dieffenbachia poisoning is painful irritation caused by chewing the plant or contacting its sap.

WHAT IT IS
Irritation after contact with a Dieffenbachia plant

BODY SYSTEM
Mouth, throat, eyes, skin, and digestive system

MAIN SYMPTOMS
Burning pain, swelling, drooling, trouble swallowing

WHAT CAUSES IT
Needle-shaped calcium oxalate crystals in plant sap

HOW IT IS DIAGNOSED
Exposure history and physical examination

MAIN TREATMENT
Rinsing, cooling measures, pain relief, and airway support

CAN IT BE CURED?
Yes - most effects settle with supportive care

IS IT AN EMERGENCY?
Sometimes - breathing difficulty or rapidly worsening mouth swelling

LONG-TERM OUTLOOK
Full recovery expected after most exposures

Overview

Dieffenbachia poisoning is an irritating injury caused by contact with a Dieffenbachia houseplant. It most often happens when someone bites or chews a leaf or stem. Sap can also irritate the eyes and skin.

The plant contains bundles of microscopic, needle-shaped calcium oxalate crystals. When the plant is damaged, these crystals can be released into nearby tissue. This causes immediate burning, pain, and swelling. Other plant chemicals may add to the irritation.

Most exposures affect only the area that touched the plant. Symptoms are often painful but improve with rinsing, simple comfort measures, and time. This is different from a poison that must be absorbed into the bloodstream before causing symptoms.

Severe swelling of the tongue or throat is uncommon, but it can interfere with breathing. Eye exposure may also damage the surface of the eye if the sap is not washed out promptly.

Symptoms

Symptoms usually begin during or soon after contact with the plant.

After chewing or swallowing part of a Dieffenbachia, symptoms may include:

A person may refuse food or drinks because swallowing hurts. Young children may cry, spit, or touch their mouths rather than explain what happened.

Sap in the eye can cause intense burning, redness, watering, sensitivity to light, or blurred vision. The eyelids may become swollen. Symptoms that continue after thorough rinsing need medical assessment because the clear surface of the eye, called the cornea, may be injured.

Skin contact can cause redness, itching, burning, or a rash-like reaction. Skin symptoms are usually limited to areas that touched the sap.

Breathing difficulty, noisy breathing, or rapidly increasing tongue and throat swelling are emergency symptoms. They may indicate that swelling is narrowing the airway.

- Immediate burning or stinging in the mouth
- Swelling of the lips, tongue, mouth, or throat
- Drooling or difficulty swallowing
- A hoarse voice or trouble speaking clearly
- Nausea, vomiting, or stomach discomfort

Causes and risk factors

All parts of a Dieffenbachia plant can be irritating, including its leaves, stems, and sap. Chewing damages special plant cells and releases calcium oxalate crystals. The crystals penetrate the moist lining of the mouth and trigger pain and inflammation.

This is usually a direct irritant effect rather than a typical food allergy. It can happen the first time someone contacts the plant.

Exposure is more likely when:

The amount of plant material, the area exposed, and how long sap remains on the skin or eye can affect the severity. A small bite that is quickly spat out may still cause marked mouth pain because the crystals act on contact.

Dogs and cats can develop similar mouth irritation after chewing the plant. A veterinarian or animal poison service should advise on pet exposures.

- A young child bites an accessible houseplant
- Leaves or stems are cut, broken, or handled without protection
- Sap is transferred from the hands to the eyes or mouth
- Plant material is mistaken for an edible leaf
- A person has difficulty recognizing or safely handling household plants

Diagnosis

There is no routine blood test that confirms Dieffenbachia poisoning. Diagnosis is usually based on a known or likely plant exposure, the rapid start of symptoms, and an examination of the mouth, throat, skin, or eyes.

A clear photograph of the plant may help identify it. Do not handle broken plant pieces with bare hands to collect a sample. If you bring plant material for identification, place it in a closed container or bag.

During a medical assessment, a clinician may check breathing, oxygen level, voice changes, and the ability to swallow saliva. The mouth and throat may be examined for swelling or tissue injury.

Persistent eye symptoms may require an examination with dye and a special light. This can show scratches or other damage to the cornea.

Further tests are not usually needed after a mild, clearly identified exposure. They may be considered if symptoms are severe, the plant is uncertain, another substance may have been swallowed, or vomiting has caused dehydration.

Treatment and management

Start by stopping further contact. Spit out any plant material and gently remove visible pieces from the mouth. Rinse the mouth with water. Do not forcefully scrape the tongue or gums, as this may worsen irritation.

If the person is fully awake and swallowing normally, cool water, milk, ice chips, or a cold soft food may ease discomfort. Do not give anything by mouth if there is choking, marked drooling, increasing swelling, or difficulty swallowing.

Do not induce vomiting. Vomiting brings irritating material back across the mouth and throat and may increase the risk of choking. Do not try to neutralize the sap with chemicals or household remedies.

Wash exposed skin with soap and plenty of water. Remove clothing or jewellery contaminated with sap.

For eye exposure, begin rinsing immediately with clean, lukewarm running water. Remove contact lenses if this is easy, then continue rinsing for at least 15 to 20 minutes. Avoid rubbing the eye.

Contact a poison information service for advice based on the person's age, symptoms, and type of exposure. In the United States, Poison Control is available at 1-800-222-1222. Elsewhere, use the local poison centre or emergency health service.

Medical treatment is supportive, meaning it controls symptoms while the tissue recovers. It may include pain relief, fluids, treatment for vomiting, and observation for worsening swelling. Clinicians may use additional medicines when inflammation is significant, but no medicine removes the embedded crystals immediately.

Someone with serious mouth or throat swelling may need close airway monitoring. Emergency treatment can include oxygen, breathing support, and placement of a breathing tube if the airway is becoming blocked.

Complications

Most exposures settle without lasting harm. Complications are more likely after substantial chewing, delayed eye rinsing, or severe swelling.

Possible complications include:

Severe airway obstruction is uncommon, but it is the main immediate danger. Eye injuries can also last longer than mouth symptoms and may require follow-up until pain and vision return to normal.

Dieffenbachia does not usually cause the widespread calcium problems associated with swallowing large amounts of soluble oxalate chemicals. Its crystals mainly cause local tissue injury.

- Airway narrowing from tongue or throat swelling
- Dehydration when pain or vomiting prevents drinking
- Breathing plant material or vomit into the lungs

- Corneal scratches or inflammation after eye exposure
- Ongoing mouth or throat pain that limits eating

Prevention

Dieffenbachia poisoning is preventable by avoiding contact with the plant and its sap.

Keep the plant where young children and pets cannot reach it. A high shelf is only useful if leaves cannot fall and nearby furniture cannot be climbed. Consider choosing a different plant if safe separation is not practical.

When trimming or repotting Dieffenbachia, wear gloves and avoid touching your face. Eye protection can reduce the chance of sap splashing into an eye. Wash tools, hands, and work surfaces afterward.

Do not prepare food on a surface where the plant has been cut. Dispose of leaves and stems in a closed rubbish or compost container that children and animals cannot access.

Teach children not to taste houseplants, but do not rely on teaching alone for toddlers. Keep the plant's label or take a clear photograph so it can be identified quickly after an exposure.

Living with it and what to expect

Dieffenbachia poisoning is an acute injury rather than an ongoing illness. The plant material does not remain active in the body for months or years.

After a mild mouth exposure, burning and swelling often begin to ease over several hours. Tenderness or discomfort with eating can continue for a few days. Recovery may take longer if there was extensive swelling or injury to the eye.

Soft, cool foods may be more comfortable while the mouth heals. Avoid very hot, spicy, acidic, or rough foods if they make the pain worse. Normal foods can be resumed as swallowing becomes comfortable.

Routine medical follow-up is not usually needed when symptoms are mild and steadily resolving. Follow-up may be advised after an eye injury, significant throat swelling, or symptoms that interfere with eating and drinking.

Once the affected tissue has healed, lasting effects are not expected after most exposures. Contact with the plant again can cause another reaction, so the plant should be moved, removed, or handled with suitable protection.

When to get medical help

Call emergency services now if there is:

Continue rinsing an exposed eye while arranging help. Sudden vision loss, severe eye pain, or inability to open the eye needs urgent assessment.

Contact a poison centre or seek prompt medical advice if a child has chewed the plant, the plant cannot be identified, or symptoms are more than mild. Medical assessment is also appropriate for repeated vomiting, worsening pain, persistent voice changes, or difficulty drinking enough fluid.

After eye exposure, seek advice if pain, redness, tearing, light sensitivity, or blurred vision continues after thorough rinsing.

A small exposure with mild symptoms that are clearly improving may be managed at home after poison-centre advice. Monitor for increasing swelling or swallowing problems rather than relying only on the amount of plant material involved.

- Trouble breathing or noisy, high-pitched breathing
- Rapidly worsening swelling of the tongue, mouth, or throat
- Inability to swallow saliva or severe choking
- Blue, grey, or unusually pale lips or skin
- Collapse, severe weakness, or reduced responsiveness

Common questions

How poisonous is Dieffenbachia?

Dieffenbachia is strongly irritating, but most human exposures cause local pain and swelling rather than whole-body poisoning. Serious airway swelling is possible but uncommon. The symptoms and ability to breathe or swallow matter more than the size of the bite alone.

What should I do if my child bites a leaf?

Remove any visible plant material, have the child spit, and rinse the mouth with water. Do not induce vomiting. If the child is awake and swallowing normally, a cool drink or ice pop may be soothing. Contact a poison centre for case-specific advice and watch for drooling, increasing swelling, or breathing difficulty.

Can Dieffenbachia poisoning be fatal?

Death is very uncommon. The main life-threatening risk is swelling that blocks the airway.

Breathing difficulty, noisy breathing, or rapidly worsening tongue or throat swelling requires emergency help.

Should I give milk after someone chews the plant?

A small amount of cold milk or water may soothe the mouth if the person can swallow normally. It is a comfort measure, not an antidote. Give nothing by mouth if swallowing is difficult or swelling is increasing.

Can touching the leaves poison me?

Touching an intact leaf is less likely to cause symptoms than contact with sap from a cut or broken plant. Sap can irritate the skin and may cause severe pain if transferred to the eyes. Wash your hands after handling the plant.

How long do symptoms last?

Mild mouth symptoms often improve over several hours and settle within a few days. Eye injuries or substantial mouth and throat swelling may last longer. Symptoms that worsen or fail to improve should be medically assessed.

Sources

- MedlinePlus Medical Encyclopedia: Dieffenbachia poisoning , U.S. National Library of Medicine
- Poison Control: Dieffenbachia and Philodendron-popular but poisonous , National Capital Poison Center
- Gardner DG. "Injury to the oral mucous membranes caused by the common houseplant, Dieffenbachia: a review." Oral Surgery, Oral Medicine, Oral Pathology . 1994;78(5):631-633
- Arditti J, Rodriguez E. "Dieffenbachia: uses, abuses and toxic constituents-a review." Journal of Ethnopharmacology . 1982;5(3):293-302