CONDITION

Disseminated tuberculosis

Disseminated tuberculosis is a tuberculosis infection that has spread through the blood or lymph system to several parts of the body.

WHAT IT IS
Tuberculosis infection spread to multiple nonadjacent body sites

BODY SYSTEM
Multiple organs, often lungs, liver, spleen, bone marrow, or brain

MAIN SYMPTOMS
Fever, night sweats, weight loss, severe tiredness

WHAT CAUSES IT
Mycobacterium tuberculosis infection spreading through the body

HOW IT IS DIAGNOSED
Molecular testing and culture of blood, fluid, sputum, or tissue

MAIN TREATMENT
Several anti-tuberculosis antibiotics taken together for months

CAN IT BE CURED?
Yes - combination antibiotics can eliminate the infection

IS IT AN EMERGENCY?
Sometimes - breathing trouble, confusion, seizures, or severe weakness

LONG-TERM OUTLOOK
Often curable, but delayed treatment increases serious complications

Overview

Disseminated tuberculosis is tuberculosis, or TB, that has spread from its original site to two or more separate parts of the body. The bacteria usually travel through the bloodstream or lymph system.

TB most often begins in the lungs, but disseminated TB may involve the liver, spleen, bone marrow, kidneys, brain, lining around the heart, bones, or other tissues.

Some people develop many tiny spots throughout the lungs or other organs. On an X-ray, these may look like millet seeds, so this pattern is called miliary TB . Miliary TB and disseminated TB overlap, but they do not always mean exactly the same thing.

Disseminated TB can affect several organs at once. Symptoms may be gradual and

nonspecific, which can delay diagnosis. Prompt treatment improves the chance of recovery and reduces the risk of lasting organ damage.

Symptoms

Symptoms may develop over days or weeks. They can progress more quickly in people with a weakened immune system.

Common symptoms include:

Other symptoms depend on the organs involved. TB affecting the brain or its lining, called TB meningitis , may cause persistent headache, vomiting, confusion, neck stiffness, unusual sleepiness, weakness, or seizures.

Bone marrow involvement can reduce blood cell production, causing paleness, breathlessness, frequent infections, or easy bruising. Liver involvement may cause abdominal swelling or yellowing of the skin or eyes. TB around the heart can cause chest discomfort, breathlessness, or leg swelling.

These symptoms have many possible causes and do not necessarily mean that you have disseminated TB.

- Fever, chills, or night sweats
- Unplanned weight loss or reduced appetite
- Marked tiredness or weakness
- Cough, chest discomfort, or shortness of breath
- Swollen lymph nodes
- Abdominal discomfort or swelling

Causes and risk factors

Disseminated TB is caused by Mycobacterium tuberculosis . The bacteria spread through the air when someone with infectious TB of the lungs or throat coughs, speaks, or sings.

After entering the lungs, TB bacteria may be controlled by the immune system without being completely removed. This is called latent TB infection . Latent TB causes no symptoms and is not contagious, but it can later become active. Dissemination can occur during a new infection or when an earlier infection becomes active again.

Anyone can develop disseminated TB, but the risk is higher when the immune system cannot

contain the bacteria effectively. Risk factors include:

Disseminated TB is not inherited. Cases within a family usually reflect shared exposure rather than a genetic condition.

- HIV infection, especially without effective treatment
- Immune-suppressing medicines, including some cancer treatments and biologic medicines
- Organ or stem cell transplantation
- Very young or older age
- Poor nutrition
- Diabetes or advanced kidney disease
- Heavy alcohol use
- Recent TB infection or close contact with infectious TB

Diagnosis

Diagnosis usually requires several tests because symptoms and scans alone cannot confirm TB. Testing may begin urgently if disseminated TB or TB meningitis is suspected.

A chest X-ray may show the tiny, widespread spots associated with miliary TB. CT or MRI scans can provide more detail about the lungs, brain, abdomen, bones, or other organs.

Samples are tested for TB bacteria. Depending on the symptoms, these may include sputum, blood, urine, fluid from around an organ, bone marrow, or a small tissue sample called a biopsy.

A molecular test can quickly detect TB genetic material and may identify resistance to certain antibiotics like amoxicillin or doxycycline . A culture attempts to grow the bacteria, confirming the diagnosis and allowing fuller resistance testing. Culture results may take several weeks.

A negative sample does not always rule out disseminated TB because bacteria may be present in low numbers. More than one sample type may be needed.

A TB skin or blood test can show whether the immune system has encountered TB, but it cannot distinguish latent from active infection. Results may also be negative during severe illness or when the immune system is weakened.

Blood tests assess blood cell counts and liver and kidney function. HIV testing is normally offered because HIV affects treatment planning and the risk of other infections.

Treatment and management

Disseminated TB is treated with several anti-TB antibiotics at the same time. Combination treatment is essential because using a single medicine can allow resistant bacteria to survive.

For TB that is sensitive to standard medicines, treatment commonly lasts at least six months. A longer course may be needed when the brain, spinal cord, bones, or joints are involved, or when the bacteria are drug-resistant. The exact plan depends on resistance tests, affected organs, other health conditions, and treatment tolerance.

Treatment may begin in hospital if you are very unwell, the diagnosis is uncertain, or there is possible brain, heart, or breathing involvement. Care may involve an infectious diseases or TB specialist and the local public health service.

Corticosteroids like prednisone or hydrocortisone may be used to reduce inflammation in certain forms of TB, particularly TB meningitis or TB affecting the lining around the heart. They are not needed in every case.

Monitoring may include blood tests, symptom reviews, eye checks for certain medicines, and repeat imaging. Tell the treatment team promptly about yellow skin or eyes, severe nausea, new vision changes, marked weakness, numbness, rash, or other concerning symptoms. These can be medicine side effects.

Missing doses can make treatment less effective and increase drug resistance. TB services may offer reminders, video appointments, supervised treatment, transport support, or help with medicine costs.

If you have HIV, both TB and HIV need treatment. The timing of HIV medicines is planned carefully because immune recovery can briefly worsen inflammation around TB sites. This is called immune reconstitution inflammatory syndrome , or IRIS.

Complications

Without effective treatment, disseminated TB can damage several organs and become life-threatening. Complications include:

Symptoms can sometimes worsen temporarily after treatment begins because of inflammation, including IRIS. New or worsening symptoms still require assessment and should not be assumed to be part of recovery.

- TB meningitis, with possible stroke, hearing loss, seizures, or lasting neurological problems

- Severe breathing failure
- Fluid or inflammation around the heart
- Liver, kidney, or bone marrow problems
- Adrenal gland damage affecting blood pressure and body salts
- Bone or joint damage
- Infection resistant to standard medicines

Prevention

Disseminated TB cannot always be prevented, particularly when an earlier infection was unknown. Finding and treating TB infection before it becomes active can lower the risk.

People who have had close contact with someone with infectious TB may be offered a skin or blood test and sometimes a chest X-ray. Treatment for latent TB reduces the chance that the bacteria will later become active.

If active TB involves the lungs or throat, measures may be needed to reduce airborne spread. These can include temporarily avoiding work, school, or shared indoor spaces, improving ventilation, and wearing a mask when advised. A TB or public health team determines when precautions are no longer needed.

Disseminated TB does not automatically mean that you are contagious. Infectiousness mainly depends on whether active TB is present in the lungs or throat and whether bacteria are found in respiratory samples.

The BCG vaccine offers some protection against severe TB in young children, but it does not provide complete protection. It is used according to local policies and individual risk.

Living with it and what to expect

Treatment takes months, and improvement is often gradual. Fever and appetite may improve first, while tiredness and weight loss can take longer. Scans may remain abnormal after you feel better because inflammation and tissue changes take time to settle.

Follow-up continues throughout treatment. The care team checks whether symptoms are improving, whether the bacteria are sensitive to the medicines, and whether side effects are developing. Do not stop treatment because you feel better unless the TB team changes the plan.

Most people with drug-sensitive disseminated TB can recover with timely, completed treatment. Recovery may take longer when several organs are involved, diagnosis was delayed, or the immune system is weakened. Brain, bone, or other organ involvement may leave lasting effects requiring rehabilitation or specialist follow-up.

After successful treatment, routine long-term antibiotics are not usually needed. Follow-up depends on the organs affected and the risk of recurrence. A returning fever, night sweats, weight loss, cough, or new neurological symptoms should be assessed.

TB services may also help with work or school documentation, contact testing, travel plans, housing concerns, and arranging medicines around daily life.

When to get medical help

Arrange prompt medical assessment if you have persistent fever, night sweats, unexplained weight loss, or a cough lasting several weeks, especially after close TB exposure or if your immune system is weakened.

If you are being treated for TB, contact the treatment team promptly about worsening symptoms, difficulty taking the medicines, missed doses, or possible side effects. This requires timely follow-up but does not always need emergency care.

Seek emergency help for:

- Severe or rapidly worsening difficulty breathing
- Confusion, unusual drowsiness, fainting, or a seizure
- A severe headache with vomiting, weakness, or neck stiffness
- New weakness on one side, trouble speaking, or loss of consciousness
- Severe chest pain or signs of shock, such as cold skin and extreme weakness
- A severe allergic reaction, including facial swelling or difficulty breathing

Common questions

Is disseminated tuberculosis contagious?

Not always. It is mainly contagious when active TB also affects the lungs or throat and bacteria can enter the air. TB limited to other organs is usually not spread through ordinary contact. Sputum tests and chest imaging help determine whether precautions are needed.

Is disseminated TB the same as miliary TB?

Not exactly. Disseminated TB means infection is present at two or more nonadjacent sites. Miliary TB describes widespread tiny spots caused by blood-borne spread, often seen in the lungs. A person may have both.

Can disseminated tuberculosis be cured?

Yes. Many cases can be cured with a complete course of effective combination antibiotics. Outcomes depend on the organs involved, antibiotic resistance, immune health, and how quickly treatment begins.

How long does treatment take?

Treatment commonly lasts at least six months. It may be longer for TB involving the brain, spinal cord, bones, or joints, or when drug resistance is present. Test results and treatment response guide the final length.

Can I be around my family during treatment?

That depends on whether you have infectious TB in the lungs or throat. A TB or public health team may recommend temporary precautions and testing for close contacts. Normal contact can usually resume after the team confirms that it is safe.

Can disseminated TB come back after treatment?

It can recur, but completing effective treatment lowers the risk. Recurrence may result from remaining bacteria or a new exposure. Seek assessment if TB-type symptoms return.

Sources

- Centers for Disease Control and Prevention. Core Curriculum on Tuberculosis: What the Clinician Should Know . Seventh edition, 2021.
- Centers for Disease Control and Prevention. Clinical Overview of Tuberculosis Disease . CDC Tuberculosis clinical guidance.
- Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis . Clinical Infectious Diseases. 2016;63(7):e147-e195.
- World Health Organization. WHO Consolidated Guidelines on Tuberculosis: Module 4-Treatment: Drug-Susceptible Tuberculosis Treatment . 2022 update.
- National Institute for Health and Care Excellence. Tuberculosis: Prevention, Diagnosis, Management and Service Organisation . NICE guideline NG33.

- Sharma SK, Mohan A, Sharma A, Mitra DK. Miliary Tuberculosis: New Insights Into an Old Disease . The Lancet Infectious Diseases. 2005;5(7):415-430.