CONDITION

Food addiction

Food addiction describes a debated pattern of compulsive eating that continues despite distress or unwanted consequences.

WHAT IT IS
Compulsive, addiction-like eating despite unwanted consequences

BODY SYSTEM
Brain and digestive system

MAIN SYMPTOMS
Cravings, loss of control, repeated overeating, distress

WHAT CAUSES IT
Unknown; brain, stress, and environment may contribute

HOW IT IS DIAGNOSED
Clinical assessment; no accepted confirmatory test

MAIN TREATMENT
Psychological support, regular eating, and related-condition care

CAN IT BE CURED?
Sometimes - symptoms may resolve with sustained support

IS IT AN EMERGENCY?
Sometimes - with suicidal thoughts or severe physical illness

LONG-TERM OUTLOOK
Symptoms can improve, but setbacks may occur

Overview

Food addiction is a term used for eating that feels compulsive or out of control. A person may have strong cravings, repeatedly eat more than intended, or continue despite physical, emotional, or social consequences.

The term remains scientifically debated. Food addiction is not a separate diagnosis in the current major psychiatric classification systems, the DSM-5-TR and ICD-11. No blood test, brain scan, or other test confirms it.

Some researchers think certain eating patterns can resemble substance addiction, with intense urges, difficulty cutting down, and continued behaviour despite harm. Foods high in refined carbohydrates, fat, or salt are often mentioned, but no food or ingredient has been proved to cause addiction in the same way as an addictive drug.

Others argue that the behaviour is better understood through established conditions such as binge-eating disorder, bulimia nervosa, depression, trauma-related difficulties, or restrictive dieting. "Eating addiction" is sometimes preferred because the pattern involves behaviour, emotions, and circumstances rather than one substance.

Whatever label is used, loss of control around eating can cause real distress and deserves respectful assessment rather than blame.

Symptoms

People who identify with food addiction may describe:

Some people eat without physical hunger, hide their eating, or feel unable to stop once they start. Guilt, shame, anxiety, or low mood may follow.

This differs from occasionally overeating or eating more during celebrations. A concerning pattern is repeated, difficult to control, and disruptive to daily life.

Symptoms may overlap with binge-eating disorder. A binge involves eating an unusually large amount within a limited period while feeling unable to control the eating. Binge-eating disorder has specific criteria, including how often episodes happen and the distress they cause.

Bulimia nervosa includes recurrent binges followed by compensating behaviours such as vomiting, fasting, misusing laxatives, or excessive exercise. Restriction can also cause intense hunger and loss-of-control eating without an addiction process.

Body size does not show whether someone has compulsive eating or an eating disorder. These problems affect people at any weight.

- Powerful cravings or preoccupation with food
- Eating more, or for longer, than intended
- Repeated unsuccessful attempts to cut down
- Continuing to eat despite distress or unwanted consequences

Causes and risk factors

There is no single established cause. Research is examining how brain reward pathways, learning, stress, genetics, sleep, food availability, and mental health may interact.

Eating activates brain systems involved in pleasure, motivation, and learning. This is normal and necessary for survival. Rewarding foods may produce stronger urges in some situations,

but brain activation alone does not prove addiction.

Repeated dieting or severe restriction can increase hunger, food preoccupation, and loss-of-control eating. Treating foods as completely forbidden may strengthen a cycle of restriction, craving, overeating, and guilt.

Other possible influences include:

These factors are possible influences, not proof of a cause.

- Chronic stress, loneliness, or difficult emotions
- Trauma or adverse childhood experiences
- Depression, anxiety, or attention-deficit/hyperactivity disorder
- Irregular meals, poor sleep, or limited access to balanced food
- Easy access to heavily marketed foods

Diagnosis

There is no universally accepted medical diagnosis of food addiction and no confirmatory test.

A healthcare professional may ask about cravings, loss of control, restriction, binge episodes, compensating behaviours, mood, medicines, sleep, and effects on daily life. They may check for physical conditions that affect appetite or weight.

The Yale Food Addiction Scale is a research questionnaire based on features of substance-use disorders. It can identify addiction-like eating patterns but does not establish a formal diagnosis. Online versions should not replace a full assessment.

Assessment should consider recognised eating disorders, including binge-eating disorder and bulimia nervosa, as well as depression, anxiety, substance use, and medical causes of appetite changes.

You can have significant distress without meeting criteria for a specific disorder. Support should reflect your symptoms and needs, not only a label.

Treatment and management

Because food addiction is not an established diagnosis, there is no single standard treatment. Care focuses on the eating pattern, its triggers, and any recognised eating disorder or mental health condition.

Regular, adequate meals can reduce extreme hunger and make eating more predictable. An

eating-disorder dietitian can support a flexible pattern that meets nutritional needs. Highly restrictive diets may worsen cravings or binge-restrict cycles.

Psychological treatment may help identify what happens before, during, and after loss-of-control eating. Cognitive behavioural therapy, or CBT, addresses links between thoughts, feelings, and behaviour. Other approaches may focus on emotional regulation, relationships, trauma, or responding differently to urges.

Treatment may also address depression, anxiety, ADHD, sleep problems, or trauma. If binge-eating disorder or bulimia nervosa is diagnosed, care can follow evidence-based treatment for that condition.

No medication is approved specifically for food addiction as a standalone condition. Medicines may be considered for a diagnosed eating disorder, mental health condition, or related physical condition after an individual assessment of benefits and risks.

Peer support can reduce isolation, but programs differ. Be cautious if a program demands extreme restriction, labels many foods as toxic, promises a cure, or encourages stopping prescribed treatment.

Day-to-day strategies include noticing triggers without judgment, planning regular meals, avoiding long gaps without food, and finding non-food ways to manage stress. The goal is safer, more stable eating and less distress-not perfect control.

Complications

Compulsive or chaotic eating can affect emotional health, relationships, work, and finances. Shame may lead to secrecy and delayed support.

Repeated bingeing, restriction, vomiting, laxative misuse, or major weight changes can cause physical problems. Risks depend on the behaviour and may include dental damage, digestive symptoms, dehydration, electrolyte imbalance, and changes in blood sugar, blood pressure, or cholesterol.

Weight cycling-repeated weight loss and regain-can be physically and emotionally difficult. Body weight alone does not provide a complete picture of health.

Severe guilt or hopelessness may contribute to depression, self-harm, or suicidal thoughts. These symptoms need direct attention and are not a failure of willpower.

Prevention

There is no proven way to prevent food addiction because its definition and causes are unsettled.

A stable relationship with food may be supported by regular meals, enough food, adequate sleep, and avoiding unnecessarily strict diets. Children and adults may benefit when food is discussed without moral labels such as "good," "bad," or "guilty."

Early help for bingeing, restriction, emotional distress, or compensating behaviours may prevent greater disruption. This does not mean someone caused the problem by not seeking help sooner.

Food marketing and access, financial pressures, trauma, and mental health all affect eating. Prevention cannot rest only on individual self-control.

Living with it and what to expect

The course differs between people. Symptoms may come and go with stress, sleep, dieting, life changes, or access to support. A setback does not erase progress.

Over months, treatment often focuses on regular meals, reducing binge-restrict cycles, and identifying emotional or environmental triggers. Cravings may become less intense or easier to respond to without acting automatically.

Over years, some people no longer experience significant loss of control. Others have recurring periods of difficulty and benefit from follow-up or renewed treatment. An eating disorder or other mental health condition may also shape recovery.

Follow-up may review eating patterns, mood, physical health, and how well treatment fits your life. Weight change is not the only measure of improvement. Less secrecy, fewer binges, better concentration, improved energy, and greater flexibility around food also matter.

Recovery does not require avoiding all enjoyable food forever. For many people, rigid abstinence is neither practical nor helpful because food is necessary. The best approach depends on the behaviour, nutritional needs, and any diagnosed condition.

When to get medical help

Arrange a routine appointment if eating repeatedly feels out of control, causes distress, or interferes with relationships, work, health, or finances. Seek help if you regularly binge, severely restrict food, vomit, misuse laxatives, or exercise to compensate for eating.

Prompt assessment is important if you are fainting, becoming increasingly weak, vomiting

frequently, or struggling to eat or drink enough. These can indicate a physical complication.

Seek emergency help now for:

These signs are not specific to food addiction. They need urgent assessment regardless of the cause.

- Thoughts of suicide or an immediate risk of self-harm
- Fainting with slow recovery, confusion, or severe weakness
- Chest pain, severe shortness of breath, or a seizure
- Vomiting blood or passing black, tar-like stool
- Severe dehydration, such as very little urine with confusion

Common questions

Is food addiction a real medical diagnosis?

Not as a separate diagnosis in the DSM-5-TR or ICD-11. Addiction-like eating remains an active research area, and the distress can be real even when the label is uncertain.

How is food addiction different from binge-eating disorder?

Binge-eating disorder is a recognised condition with defined criteria. Food addiction is a broader, debated term that may include cravings or compulsive eating without meeting those criteria. They can overlap.

Can sugar be addictive?

Sugar can be rewarding, and restriction may intensify cravings. Human evidence has not established it as addictive in the same way as nicotine or alcohol. Foods usually combine carbohydrates, fats, flavours, and textures, making one ingredient difficult to isolate.

Do I need to avoid trigger foods forever?

Not necessarily. Permanent avoidance may strengthen fear or restriction for some people, while others find temporary structure helpful. A clinician or eating-disorder dietitian can help support adequate nutrition without unnecessary rules.

Can food addiction happen if I am not overweight?

Yes. Loss-of-control eating and eating disorders occur across body sizes. Weight cannot confirm or rule out a problem.

Can food addiction be cured?

Some people reach lasting recovery, while others have symptoms during stressful periods.

Treatment can reduce distress and improve control even if cravings do not disappear completely.

Sources

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR). American Psychiatric Association Publishing; 2022.
- World Health Organization. ICD-11 for Mortality and Morbidity Statistics . Sections on feeding or eating disorders and disorders due to addictive behaviours.
- Gearhardt AN, Corbin WR, Brownell KD. Development of the Yale Food Addiction Scale Version 2.0 . Psychology of Addictive Behaviors. 2016;30(1):113-121.
- Gordon EL, Ariel-Donges AH, Bauman V, Merlo LJ. What Is the Evidence for "Food Addiction?" A Systematic Review . Nutrients. 2018;10(4):477.
- Hebebrand J, Albayrak Ö, Adan R, et al. "Eating addiction," rather than "food addiction," better captures addictive-like eating behavior . Neuroscience & Biobehavioral Reviews. 2014;47:295-306.
- National Institute for Health and Care Excellence. Eating disorders: recognition and treatment, guideline NG69 . Published 2017; updated 2020.