CONDITION

Fuel oil poisoning

Fuel oil poisoning is illness or injury caused by swallowing fuel oil, breathing its fumes, or getting it on the skin or eyes.

WHAT IT IS
Harmful exposure to fuel oil or its fumes

BODY SYSTEM
Lungs, digestive tract, skin, eyes, and nervous system

MAIN SYMPTOMS
Coughing, choking, breathlessness, nausea

WHAT CAUSES IT
Swallowing, inhaling, or skin exposure to fuel oil

HOW IT IS DIAGNOSED
Exposure history, examination, oxygen levels, chest X-ray

MAIN TREATMENT
Supportive care and breathing support

CAN IT BE CURED?
Sometimes - most limited exposures resolve completely

IS IT AN EMERGENCY?
Sometimes - breathing trouble or reduced alertness

LONG-TERM OUTLOOK
Most recover; severe aspiration may cause lasting lung damage

Overview

Fuel oil poisoning happens when fuel oil enters the body or damages tissues. Fuel oils include heating oil, kerosene, diesel fuel, and similar petroleum products. Their exact ingredients differ, so the product label or container can be important when getting advice.

The main danger after swallowing fuel oil is not usually absorption through the stomach. It is aspiration , which means the liquid enters the lungs. This may happen while swallowing, coughing, choking, or vomiting. Even a small amount in the lungs can cause chemical inflammation called chemical pneumonitis .

Breathing concentrated fumes can irritate the airways and affect the brain, particularly in an enclosed or poorly ventilated place. Fuel oil can also irritate the eyes and remove protective oils from the skin.

There is no specific antidote. Care focuses on stopping the exposure, protecting the lungs, and supporting breathing while the body recovers. Many limited exposures do not cause lasting harm, but breathing symptoms need prompt attention because lung problems can develop over several hours.

Symptoms

Symptoms depend on the fuel product, how exposure happened, how much was involved, and whether any entered the lungs.

After swallowing fuel oil, you may have coughing, choking, gagging, nausea, vomiting, or stomach discomfort. A burning feeling may occur around the mouth or throat. Vomiting is important because it raises the chance of fuel entering the lungs.

Fuel in the lungs can cause persistent coughing, wheezing, chest discomfort, rapid breathing, or shortness of breath. Severe exposure may lower the amount of oxygen in the blood. Lung symptoms can begin quickly, but they may also take several hours to become clear.

Breathing concentrated fumes may cause headache, dizziness, nausea, drowsiness, poor coordination, or confusion. Very high exposure can lead to collapse, seizures, or loss of consciousness.

Skin contact may cause redness, dryness, cracking, or a burning sensation. Repeated contact can cause dermatitis, an inflamed and itchy skin rash. Eye exposure can cause pain, redness, watering, or blurred vision.

Causes and risk factors

Fuel oil poisoning is caused by direct exposure rather than infection, inherited traits, or an immune disorder. Exposure may happen through swallowing, breathing fumes, skin contact, or an eye splash.

Accidental swallowing can happen when fuel is stored in a drink bottle or another unlabelled container. Trying to siphon diesel or heating oil by mouth is another important cause. Children may be exposed when fuel products are within reach.

Fume exposure is more concerning in tanks, garages, basements, engine spaces, or other poorly ventilated areas. Entering a space with concentrated fumes can be dangerous, including for someone attempting a rescue without suitable equipment.

The risk of aspiration is higher if a person vomits, has difficulty swallowing, is very drowsy, or

has an impaired cough reflex. Products containing additional solvents, pesticides, or other chemicals may cause effects beyond those expected from fuel oil alone.

Repeated workplace exposure can irritate the skin and airways. This is different from a single episode of acute poisoning, but it still deserves assessment and better exposure controls.

Diagnosis

Fuel oil poisoning is usually diagnosed from what happened, the product involved, and the symptoms. If it is safe to do so, keep the original container or take a photograph of its label. Do not carry a leaking or contaminated container into a vehicle or healthcare setting.

A healthcare professional may check your breathing rate, alertness, heart rate, blood pressure, temperature, and blood oxygen level. They may listen for wheezing or crackling sounds in the lungs.

A chest X-ray may be used when there is coughing, choking, breathlessness, low oxygen, or concern that fuel entered the lungs. Changes from chemical pneumonitis may not appear immediately, so an early normal X-ray does not always exclude lung injury.

Blood tests and an electrocardiogram, or ECG, may be needed after a significant exposure or if there are heart, brain, breathing, or metabolic symptoms. There is no routine blood test that confirms uncomplicated fuel oil poisoning. Testing is chosen to look for injury and guide supportive care.

Treatment and management

Move away from fumes and into fresh air if this can be done without entering an unsafe space. Do not attempt an unprotected rescue in a tank or enclosed area. Emergency services have equipment for hazardous environments.

If fuel oil has been swallowed, do not make yourself or another person vomit. Vomiting can draw fuel into the lungs. Rinse the mouth gently and contact a poison information service for instructions. Do not give food, drink, activated charcoal, or a home remedy unless a poison specialist or healthcare professional advises it.

Remove fuel-soaked clothing while avoiding further skin contact. Wash exposed skin with soap and lukewarm water. If fuel enters an eye, rinse it with clean lukewarm water and remove contact lenses if they come out easily. Persistent pain, redness, or vision changes need medical assessment.

In a healthcare setting, treatment may include observation, oxygen, and repeated checks of breathing and blood oxygen. People with wheezing may receive medicine to open the airways. Severe lung injury may require assisted breathing through a ventilator.

Activated charcoal is not routinely used for fuel oil because it does not remove these products well and may increase vomiting and aspiration. Emptying the stomach is also generally avoided. A toxicology specialist may recommend different care if the product contains another dangerous substance.

Antibiotics like amoxicillin or doxycycline do not treat chemical irritation itself and are not routinely used unless there is evidence of a bacterial infection. Steroid medicines have not shown routine benefit for hydrocarbon-related chemical pneumonitis.

Complications

The most important complication is chemical pneumonitis. Inflamed lungs may become less able to transfer oxygen into the blood. In severe cases, this can progress to respiratory failure or acute respiratory distress syndrome, a serious form of widespread lung injury.

Other possible complications include infection after lung injury, an abnormal heart rhythm, seizures, or reduced consciousness. These are more likely after a large or concentrated exposure.

Prolonged skin contact can cause chemical burns or persistent dermatitis. Eye splashes can injure the surface of the eye, especially if the fuel is not rinsed away promptly.

Most people with limited exposure recover without permanent problems. Lasting breathing difficulty or lung scarring is possible after severe aspiration, although it is not the usual outcome.

Prevention

Fuel oil poisoning can often be prevented. Store heating oil, kerosene, diesel, and related products in their original, clearly labelled containers. Never put fuel into a bottle, cup, or food container.

Keep fuel locked away from children and separate from food, medicines, and household drinks. Do not siphon fuel by mouth. Use pumps or other equipment designed for transferring petroleum products.

Use fuel only in equipment intended for that product. Provide effective ventilation when

engines, heaters, or fuel systems are used or serviced. Never enter a tank or enclosed fuel space unless it has been assessed as safe and you have the required training and protective equipment.

At work, follow instructions for gloves, eye protection, ventilation, spill control, and contaminated clothing. Report repeated skin or fume exposure so that the source can be corrected.

Living with it and what to expect

A brief exposure that causes no symptoms may only require advice and a period of observation. If symptoms do not develop during the recommended observation period, long-term effects are less likely.

Chemical pneumonitis may worsen during the first day or two before it begins to improve. Recovery can take days or weeks, depending on how much fuel entered the lungs and whether breathing support was needed. Cough, tiredness, or reduced exercise tolerance may continue for a time after leaving hospital.

A severe exposure can require intensive care and a longer rehabilitation period. Some people have persistent breathlessness or reduced lung function, but this is not inevitable. Follow-up may include a breathing assessment, oxygen measurement, chest imaging, or lung function tests when symptoms continue.

Repeated exposure at home or work should not be treated as an unavoidable part of recovery. Ongoing cough, rash, headaches, or dizziness may mean the source of exposure has not been controlled. Occupational health services can help with workplace ventilation, protective equipment, and safe return to duties.

When to get medical help

Contact a poison information service promptly after fuel oil is swallowed, even if there are no symptoms. Do not wait for coughing or breathing difficulty to appear. In the United States, Poison Control is available at 1-800-222-1222 . Elsewhere, use your national poison centre or local urgent medical service.

Call emergency services now if there is:

Move into fresh air only if you can do so safely. Do not enter an enclosed contaminated area to rescue someone without specialist training and protective equipment.

Seek urgent assessment for ongoing vomiting, chest pain, low oxygen readings, worsening drowsiness, or eye pain and blurred vision after rinsing. A cough, fever, wheeze, or shortness of breath that appears or worsens in the hours after exposure also needs prompt review.

Routine follow-up is appropriate for a cough, reduced stamina, skin rash, or other symptoms that continue after initial treatment but are not rapidly worsening. New breathing difficulty is not a routine follow-up issue and should be assessed urgently.

- Trouble breathing, severe wheezing, or rapidly worsening coughing
- Blue, grey, or unusually pale lips or skin
- Collapse, a seizure, confusion, or difficulty waking
- Persistent choking after swallowing fuel
- Exposure to concentrated fumes in an enclosed space
- A large exposure or uncertainty about what chemicals were involved

Common questions

What should I do if I swallowed fuel oil?

Do not make yourself vomit. Rinse your mouth and contact a poison information service immediately for instructions based on the exact product and exposure. Call emergency services if coughing, choking, drowsiness, or breathing trouble develops.

How quickly do fuel oil poisoning symptoms start?

Coughing, choking, nausea, dizziness, and eye or skin irritation may start immediately. Lung inflammation can become clearer over several hours. Feeling well at first does not always rule out aspiration, especially if you coughed or choked while swallowing.

Can the smell of heating oil make you sick?

Yes. Strong fumes can cause headache, nausea, dizziness, throat irritation, or drowsiness, particularly in an enclosed space. Move to fresh air if it is safe. Persistent symptoms or any breathing difficulty need medical advice.

Is fuel oil poisoning the same as gasoline poisoning?

Both involve petroleum hydrocarbons and can damage the lungs if aspirated. However, gasoline evaporates more readily, and individual products contain different chemicals. Tell the poison service or healthcare team the exact product whenever possible.

Will fuel oil show up on a chest X-ray?

Fuel oil itself may not be visible. A chest X-ray can show inflammation caused by fuel entering the lungs, but these changes may be absent early in the exposure. The decision to observe or treat you is based on the full situation, not the X-ray alone.

Can you recover completely from fuel oil poisoning?

Many people recover fully after a limited exposure. Recovery is less predictable when a significant amount enters the lungs, oxygen levels fall, or intensive breathing support is required. Continuing symptoms may need follow-up to check lung recovery.

Sources

- Agency for Toxic Substances and Disease Registry: Medical Management Guidelines for Fuel Oils , U.S. Centers for Disease Control and Prevention
- MedlinePlus Medical Encyclopedia: Fuel oil poisoning , U.S. National Library of Medicine
- Merck Manual Professional Edition: Hydrocarbon Poisoning
- Royal Children's Hospital Melbourne: Clinical Practice Guideline-Hydrocarbon poisoning
- Tormoehlen LM, Tekulve KJ, Nañagas KA. "Hydrocarbon toxicity: A review." Clinical Toxicology . 2014;52(5):479-489. doi:10.3109/15563650.2014.923904