CONDITION

Grass and weed killer poisoning

Grass and weed killer poisoning is illness caused by swallowing, breathing in, or contacting a harmful amount of herbicide.

WHAT IT IS
Harmful exposure to chemicals designed to kill unwanted plants

BODY SYSTEM
Digestive, breathing, nervous, kidney, liver, and circulation systems

MAIN SYMPTOMS
Nausea, vomiting, mouth irritation, breathing difficulty

WHAT CAUSES IT
Swallowing, inhaling, or skin and eye exposure

HOW IT IS DIAGNOSED
Exposure history, examination, and targeted blood or urine tests

MAIN TREATMENT
Decontamination, monitoring, and supportive hospital care

CAN IT BE CURED?
Sometimes - mild exposures recover; severe injury may be irreversible

IS IT AN EMERGENCY?
Sometimes - swallowing or serious breathing or neurological symptoms

LONG-TERM OUTLOOK
Mild cases recover; severe poisoning can cause lasting organ damage

Overview

Grass and weed killers are herbicides, which are chemicals used to control unwanted plants. Poisoning happens when enough of a product enters the body to cause harm. This may occur by swallowing it, breathing in spray or mist, or getting concentrated product on the skin or in the eyes.

Herbicides are not one chemical group. Products may contain glyphosate, paraquat, diquat, chlorophenoxy herbicides such as 2,4-D, or other active ingredients. They may also contain solvents and surfactants, which help the chemical spread or mix. These added ingredients can contribute to irritation or poisoning.

The risk is shaped by the exact product, its concentration, the amount involved, and how exposure happened. A small splash of a diluted household product is different from swallowing

an agricultural concentrate. Paraquat is especially dangerous if swallowed because a small amount of concentrated product can cause severe lung and organ damage.

Do not wait for symptoms after swallowing a herbicide. Contact a poison centre immediately. The product name and label are important because treatment is not the same for every herbicide.

Symptoms

Symptoms may begin within minutes or hours. Some serious effects, particularly after paraquat exposure, can develop over several days.

Contact with the eyes may cause burning, redness, watering, blurred vision, or injury to the eye surface. Skin exposure can cause redness, irritation, or chemical burns. Some herbicides are absorbed through damaged skin more easily than through intact skin.

Swallowing a product commonly causes:

More severe poisoning can affect several organs. Warning signs include difficulty breathing, low blood pressure, an abnormal heartbeat, seizures, reduced urine, or loss of consciousness. Blood tests may show acid levels outside the normal range, kidney or liver injury, or muscle breakdown.

Paraquat can burn the mouth, throat, and digestive tract. Kidney and liver injury may follow. Lung inflammation and scarring can then develop, causing increasing breathlessness even when early symptoms seemed limited.

Chlorophenoxy herbicides can affect the brain, muscles, heart, and kidneys after substantial exposure. Diquat can cause severe digestive irritation and may injure the kidneys, liver, or nervous system.

- Burning or soreness in the mouth and throat
- Nausea, vomiting, stomach pain, or diarrhoea
- Coughing, choking, drooling, or trouble swallowing
- Dizziness, drowsiness, confusion, or weakness

Causes and risk factors

Poisoning may follow an accidental splash, a leaking container, drift from spraying, or swallowing a product stored in a drink bottle. It can also occur during mixing or application

when the label's protective measures are not followed.

Risk is higher with concentrated agricultural products than with many ready-to-use garden sprays. However, no product should be assumed harmless without checking its ingredients and label.

Factors that can increase harm include:

Children may swallow products left within reach. People who mix or spray herbicides for work may have repeated opportunities for exposure, although routine contact is not the same as acute poisoning.

An intentional ingestion is a medical emergency. It also signals a need for compassionate mental health support once immediate physical safety has been addressed.

- Swallowing the product, especially a concentrate
- Delayed removal of contaminated clothing
- Exposure in a poorly ventilated enclosed space
- Prolonged contact with wet clothing or damaged skin
- Mixing products or using them outside label directions
- Delayed access to poison advice or medical care

Diagnosis

There is no single test for all grass and weed killer poisoning. Diagnosis usually begins with the exposure history, symptoms, examination, and the product label.

Clinicians may need the product name, active ingredients, concentration, estimated amount, time of exposure, and route of contact. Bring the container or a photograph of the full label if this can be done safely. Do not carry a leaking or contaminated container into an enclosed vehicle or healthcare setting.

Tests may include blood pressure, heart monitoring, oxygen measurement, and blood tests for kidney function, liver function, salts, acid balance, and muscle injury. Urine output may also be monitored. A chest X-ray or other imaging may be used when breathing problems or aspiration into the lungs is suspected.

Specialist laboratories can measure some herbicides. A urine dithionite test may help identify significant paraquat exposure, and blood paraquat levels can help estimate risk when available. These tests are time-sensitive and are not offered everywhere. Most glyphosate exposures are assessed from the history and clinical findings rather than a routine confirmatory test.

A poison centre or medical toxicologist may help interpret the label and decide what monitoring is needed.

Treatment and management

Immediate treatment starts by stopping further exposure. Move away from spray or fumes into fresh air without entering an unsafe area. Remove contaminated clothing and place it in a bag, avoiding contact with the chemical.

Wash exposed skin thoroughly with running water and soap. Rinse an affected eye continuously with clean, lukewarm water for at least 15 minutes, removing contact lenses if they come out easily.

If the product was swallowed, do not make yourself or another person vomit. Vomiting can cause more injury or allow liquid to enter the lungs. Do not give food, drinks, charcoal, or home remedies unless a poison specialist or healthcare professional advises it.

There is no general antidote that works for all herbicides. Hospital care may include:

Activated charcoal or another binding material may be considered soon after certain ingestions, but it must be given under professional direction. The possible benefit must be balanced against choking or aspiration.

Paraquat poisoning needs urgent specialist care. There is no proven antidote that reliably reverses established injury. Treatment focuses on limiting absorption, supporting organs, and avoiding unnecessary high oxygen levels because oxygen may worsen paraquat-related lung injury. Oxygen is still used when it is needed to treat dangerously low levels.

After a minor skin or eye exposure, poison-centre advice and thorough washing may be all that is required. Ongoing symptoms, concentrated exposures, and all swallowed exposures need professional assessment.

- Support for breathing and blood pressure
- Intravenous fluids and medicines for symptoms
- Heart, oxygen, kidney, and liver monitoring
- Treatment for seizures, burns, or chemical eye injury
- Carefully selected gut decontamination after some recent ingestions
- Kidney support or dialysis when severe complications develop

Complications

Many limited exposures cause temporary irritation and settle without lasting harm. Severe poisoning can damage the mouth, digestive tract, lungs, kidneys, liver, heart, muscles, or brain.

Vomiting can lead to dehydration or aspiration, meaning stomach contents enter the lungs. Chemical burns may cause swelling or later narrowing of the throat or oesophagus, the tube carrying food to the stomach.

Kidney injury can reduce the body's ability to remove waste and balance fluids. Muscle breakdown can place further strain on the kidneys. Severe changes in blood pressure, heart rhythm, or acid balance can be life-threatening.

Paraquat may cause progressive scarring of the lungs, called pulmonary fibrosis. This can lead to worsening breathlessness and low oxygen levels days or weeks after ingestion. Severe paraquat poisoning can be fatal despite intensive treatment.

Eye exposure can injure the clear surface at the front of the eye. Prompt rinsing lowers the amount of chemical in contact with the tissue, but persistent pain or altered vision needs urgent assessment.

Prevention

Most accidental herbicide poisonings can be prevented, although it is not possible to remove every risk when these products are used.

Keep herbicides in their original, labelled containers. Never transfer them into water, soft drink, or food containers. Store them locked away from children, food, medicines, and animal feed.

Read the complete label before mixing or spraying. Use the gloves, eye protection, clothing, or breathing protection named on the label. More protective equipment is not a substitute for correct handling and ventilation.

Avoid spraying into the wind or near other people. Do not eat, drink, or smoke while handling a product. Wash your hands and exposed skin after use, and change contaminated clothing promptly.

Do not mix herbicides unless the label specifically permits it. Mixing may change how the product behaves and can make exposure harder to assess.

Dispose of containers and unused product according to local hazardous-waste and label instructions. Do not reuse empty containers.

Living with it and what to expect

Grass and weed killer poisoning is usually an acute event rather than a lifelong condition. After a mild exposure, irritation and digestive symptoms often improve over hours or days. Recovery time cannot be predicted from the product category alone.

Follow-up may be needed after a significant ingestion or abnormal blood tests. Kidney, liver, lung, or muscle tests may be repeated to make sure an injury is improving. Eye injuries may require review until pain, redness, and vision have recovered.

People with serious lung or organ damage may need rehabilitation, breathing support, kidney treatment, or specialist follow-up over months. Some injuries can leave lasting limits on breathing or organ function. Paraquat-related lung injury may continue to progress after the first symptoms.

After discharge, keep the product details and hospital information available. Attend planned blood tests and appointments even if you feel better, because some complications can appear later.

An accidental poisoning can leave you feeling worried about delayed effects or using chemicals again. Clear advice from the poison centre or treating team can help you understand what follow-up is needed. If exposure was intentional, continued mental health care and practical support are important parts of recovery.

When to get medical help

Contact a poison centre immediately after any swallowed herbicide exposure, even if there are no symptoms. Do the same after a concentrated eye exposure, a large skin exposure, or significant inhalation of spray.

Call emergency services now if there is:

In the United States, call 911 for emergency symptoms and Poison Control at 1-800-222-1222 for immediate poison advice. Elsewhere, contact your local emergency service or poison centre.

For skin contact without symptoms, remove contaminated clothing, wash thoroughly, and contact a poison centre for product-specific advice. Persistent rash, pain, eye redness, blurred vision, cough, vomiting, or dizziness needs same-day medical assessment.

Routine follow-up is different from emergency care. Keep scheduled appointments after hospital treatment, and report new breathlessness, reduced urine, ongoing swallowing difficulty, or symptoms that return after initially improving.

- Trouble breathing, blue or grey lips, or severe coughing
- Collapse, loss of consciousness, confusion, or a seizure
- Severe mouth or throat burns, drooling, or inability to swallow
- Repeated vomiting, vomiting blood, or severe abdominal pain
- Very little urine, extreme weakness, or rapidly worsening symptoms
- A known or possible paraquat ingestion
- An intentional ingestion

Common questions

What should I do if I swallowed a small amount of weed killer?

Rinse your mouth gently and contact a poison centre immediately. Do not cause vomiting or wait for symptoms. "A small amount" can mean very different levels of risk depending on the product and concentration.

Is glyphosate poisoning always serious?

No. Brief contact with a diluted product often causes only irritation, but swallowing concentrated formulations can cause severe digestive symptoms, breathing problems, low blood pressure, kidney injury, or abnormal heart rhythms. Product-specific poison advice is still needed.

Can weed killer soak through my skin?

Some herbicides can be absorbed through skin, especially concentrated products and prolonged contact on damaged skin. Remove contaminated clothing and wash with soap and running water. Seek advice for a large exposure, burns, or ongoing symptoms.

How long after exposure can symptoms start?

Irritation, nausea, and vomiting may start within minutes or hours. Kidney, liver, or lung injury may take longer to appear. Paraquat-related breathing problems can worsen over several days, so follow-up should not be skipped.

Is there an antidote for weed killer poisoning?

There is no single antidote for all herbicides, and many have no specific antidote. Treatment limits absorption when possible, manages symptoms, and supports affected organs. Early identification of the exact product helps guide care.

Can I recover fully from herbicide poisoning?

Full recovery is common after many mild exposures. The outlook after severe poisoning is shaped by the chemical, concentration, amount absorbed, and organs affected. Severe paraquat poisoning and major organ injury may cause permanent damage or death.

Sources

- US Environmental Protection Agency. Recognition and Management of Pesticide Poisonings . 6th edition. Chapter 12: Herbicides. Office of Pesticide Programs; 2013.
- National Pesticide Information Center. Glyphosate General Fact Sheet . Oregon State University and US Environmental Protection Agency.
- National Institute for Occupational Safety and Health. Paraquat Dichloride: Systemic Agent . Emergency Response Safety and Health Database. US Centers for Disease Control and Prevention.
- Gawarammana IB, Buckley NA. Medical management of paraquat ingestion. British Journal of Clinical Pharmacology . 2011;72(5):745-757.
- Bradberry SM, Proudfoot AT, Vale JA. Glyphosate poisoning. Toxicological Reviews . 2004;23(3):159-167.