CONDITION

Halushi-behcet's syndrome

A long-term inflammatory condition that causes recurring ulcers and can affect the eyes, skin, joints, blood vessels, and nervous system.

WHAT IT IS
Immune-mediated inflammation of blood vessels throughout the body

BODY SYSTEM
Mouth, skin, eyes, joints, blood vessels, and nervous system

MAIN SYMPTOMS
Mouth ulcers, genital ulcers, eye inflammation, skin changes

WHAT CAUSES IT
Immune overactivity influenced by genes and environment

HOW IT IS DIAGNOSED
Clinical assessment and tests excluding similar conditions

MAIN TREATMENT
Anti-inflammatory and immune-suppressing medicines

CAN IT BE CURED?
No - treatment can control inflammation

IS IT AN EMERGENCY?
Sometimes - sudden vision loss, chest pain, or neurological symptoms

LONG-TERM OUTLOOK
Flare-ups may ease, but organ damage can persist

Overview

"Halushi-Behcet's syndrome" is not the standard medical spelling. It usually refers to Behçet's syndrome , named after the Turkish dermatologist Hulusi Behçet. It is also sometimes called Behçet syndrome or Behçet's disease.

Behçet's syndrome is a long-term condition involving inflammation of blood vessels. Blood-vessel inflammation is called vasculitis . It can affect arteries and veins of different sizes, so symptoms may occur in several parts of the body.

Recurring mouth ulcers are the most common feature. Genital ulcers, skin changes, painful joints, and eye inflammation may also occur. Less commonly, the condition affects major blood vessels, the brain and spinal cord, or the digestive system.

Symptoms often come in flare-ups. A flare is a period when inflammation becomes more active. Between flares, symptoms may partly or completely settle.

The condition matters because inflammation in the eyes, nervous system, or major blood vessels can cause lasting harm if it is not treated promptly. Many other symptoms, such as mouth ulcers and joint pain, are uncomfortable but do not usually cause permanent damage.

Symptoms

The pattern differs from person to person. Symptoms may appear months or years apart, and not everyone develops every feature.

Common symptoms include:

Mouth ulcers usually look like common canker sores. They often heal within one to three weeks, but new ulcers may return. Mouth ulcers alone are common and do not mean that someone has Behçet's syndrome.

Genital ulcers can occur on the vulva, vagina, scrotum, or penis. They are not caused by a sexually transmitted infection, although testing may be needed to rule out infections with similar symptoms.

Eye inflammation may involve the middle and back parts of the eye. This is called uveitis . It can cause redness, pain, floating spots, blurred vision, or reduced vision. Eye symptoms need prompt assessment because repeated inflammation can damage sight.

Inflamed joints can be painful and swollen. Joint symptoms often come and go and usually do not cause the progressive joint destruction seen in some forms of arthritis.

Blood-vessel inflammation can cause a painful swollen limb from a deep vein blood clot. It can also weaken an artery, creating a bulge called an aneurysm . Lung artery involvement is uncommon but serious.

Nervous-system involvement may cause severe headache, weakness, difficulty speaking, poor balance, confusion, or changes in sensation. Digestive involvement can cause abdominal pain, diarrhoea, or bleeding.

- Repeated, painful mouth ulcers
- Genital ulcers, sometimes followed by scarring
- Red, tender, acne-like, or raised skin spots
- Painful or swollen joints, especially knees, ankles, or wrists
- Eye pain, redness, blurred vision, or sensitivity to light

- Marked tiredness during flares

Causes and risk factors

The exact cause is not known. Behçet's syndrome appears to involve an immune system that becomes overactive and attacks healthy blood vessels and tissues.

Genes influence susceptibility. One genetic marker, HLA-B51 , is more common among people with the condition. However, most people with this marker never develop Behçet's syndrome, and a negative test does not rule it out. It is not usually inherited in a simple parent-to-child pattern.

Environmental factors may help trigger inflammation in someone who is already susceptible. Researchers have studied infections and changes in the microorganisms that normally live in the mouth and gut, but no single infection has been established as the cause.

The condition is more common along the historic Silk Road, including parts of Turkey, the Middle East, and East Asia. It can occur in any ethnic group and in any part of the world.

Behçet's syndrome often begins in early or middle adulthood, although children and older adults can develop it. Sex does not determine who gets it. In some populations, men have a greater risk of severe eye, blood-vessel, or nervous-system involvement.

It is not contagious and cannot be passed through kissing, sex, shared food, or everyday contact.

Diagnosis

There is no single blood test, scan, or biopsy that confirms Behçet's syndrome. Diagnosis is based on the pattern of symptoms, examination findings, and the exclusion of other possible causes.

A clinician will usually ask how often ulcers occur, how long they last, and whether there have been eye, skin, joint, blood-vessel, digestive, or neurological symptoms. Photographs taken during a flare can be useful if sores or skin changes have healed before an appointment.

Tests may include:

A pathergy test may support the diagnosis. The skin is pricked with a sterile needle and checked later for an unusually strong inflammatory reaction. A positive result can occur in Behçet's syndrome, but many affected people have a negative result.

Published diagnostic or classification criteria use combinations of mouth ulcers, genital ulcers,

eye disease, skin findings, blood-vessel problems, and pathergy. These criteria support a careful clinical assessment; they do not replace it.

Other conditions can cause similar symptoms. These include inflammatory bowel conditions, lupus, reactive arthritis, recurrent aphthous ulcers, infections, and some blood disorders. Diagnosis may take time when symptoms appear separately.

- Blood tests for inflammation and conditions with similar features
- Swabs or blood tests to check for infection
- A detailed eye examination
- Scans of blood vessels, the brain, or other organs when indicated
- A skin biopsy in selected cases

Treatment and management

Treatment aims to settle active inflammation, reduce future flares, relieve symptoms, and protect organs from permanent damage. The approach is based on which parts of the body are affected and how severe the inflammation is.

Milder mouth, genital, and skin symptoms may be managed with topical treatments applied directly to the affected area. These can include corticosteroid like prednisone or hydrocortisone preparations, which reduce inflammation, and pain-relieving mouth care.

Medicines that act throughout the body may be considered when ulcers, skin symptoms, or joint inflammation are frequent. Colchicine is one medicine used for some skin, genital, and joint symptoms.

Eye, nervous-system, major blood-vessel, or severe digestive involvement usually requires stronger treatment. Options include corticosteroids and medicines that suppress specific parts of the immune response. These may include conventional immune-suppressing medicines or biologic medicines such as tumour necrosis factor inhibitors.

These treatments require monitoring because they can affect infection risk, blood cells, the liver, or other organs. The benefits and risks differ between medicines.

Blood clots associated with Behçet's syndrome are linked to inflammation inside the blood vessel. Controlling that inflammation is an important part of treatment. Decisions about blood-thinning medicine require specialist assessment, particularly because an artery aneurysm can increase bleeding risk.

Surgery or a procedure may occasionally be needed for a damaged or dangerously enlarged

blood vessel. Surgery does not treat the underlying immune inflammation, so medical treatment is often needed as well.

Eye care, dental care, and support with pain, fatigue, sexual health, or emotional wellbeing may form part of ongoing management. Treatment plans should be reviewed before pregnancy or when pregnancy is possible because some medicines need to be changed.

Complications

Many ulcers and skin symptoms heal without permanent damage. Genital ulcers may leave scars, and repeated painful ulcers can interfere with eating, sleep, and sexual wellbeing.

Repeated uveitis can lead to cataracts, glaucoma, retinal damage, or permanent loss of vision. Early recognition and specialist treatment reduce this risk.

Inflammation can cause clots in deep veins, narrowing of blood vessels, or aneurysms. An aneurysm can rupture and bleed. Lung artery aneurysms may cause coughing up blood.

Inflammation of the brain or spinal cord, sometimes called neuro-Behçet's , can cause lasting problems with movement, speech, balance, memory, or behaviour. Complications depend on the area affected and how quickly inflammation is controlled.

Digestive ulcers can sometimes cause significant bleeding or a hole in the bowel wall. This is uncommon but requires urgent care.

The condition and its treatment can also affect mood, work, relationships, and energy levels. These effects are important even when tests show little permanent organ damage.

Prevention

There is no proven way to prevent Behçet's syndrome. Genetic susceptibility cannot be changed, and no particular infection or behaviour has been shown to cause the condition.

Once the condition is present, taking prescribed treatment and attending monitoring appointments can help reduce uncontrolled inflammation. Regular eye reviews are particularly important for anyone with previous eye involvement.

There is no specific diet proven to prevent flares. A food diary may help identify foods that irritate mouth ulcers, but avoiding a food does not treat blood-vessel inflammation. Good mouth care and avoiding tobacco may support general health, although they cannot cure the condition.

Living with it and what to expect

Behçet's syndrome usually has a relapsing and remitting course. This means symptoms become active for a time and then improve. The timing, length, and severity of flares cannot be predicted reliably.

For some people, the main long-term problem is recurring ulcers, skin symptoms, or joint pain. Others need continued specialist care for eye, blood-vessel, neurological, or digestive inflammation.

Disease activity often becomes less intense over many years, but this is not guaranteed. Damage already caused by severe inflammation, such as reduced vision or neurological disability, may not fully recover even after the inflammation settles.

Most people can continue many usual activities, although work, study, exercise, and social plans may need adjustment during flares. Pacing activities can help when fatigue is prominent. Gentle movement may reduce stiffness, while painful or swollen joints may need temporary rest.

Keeping a record of symptoms can show patterns over time. Include ulcers, eye symptoms, skin changes, joint swelling, headaches, digestive symptoms, and medicine changes. Photographs can help document short-lived skin or ulcer flares.

Routine follow-up may involve rheumatology, dermatology, ophthalmology, neurology, vascular, or digestive specialists. The team involved reflects the organs affected rather than the number of symptoms.

When to get medical help

Arrange a routine appointment if you have repeated mouth or genital ulcers, unexplained skin changes, recurring joint swelling, or symptoms that continue to return. These findings have many possible causes and need assessment rather than self-diagnosis.

If you already have Behçet's syndrome, contact your usual care team when symptoms become more frequent, treatment is not controlling them, or medicine side effects develop.

Seek urgent same-day medical assessment for:

Call emergency services for sudden loss of vision, chest pain, difficulty breathing, coughing up blood, fainting, new weakness on one side, difficulty speaking, a seizure, or sudden confusion. These symptoms can have several causes, but they require immediate assessment.

- A painful, red eye or new blurred vision

- A newly swollen, warm, or painful leg or arm
- Persistent abdominal pain, bloody diarrhoea, or black stools
- A new severe headache, especially with other neurological symptoms

Common questions

Is Halushi-Behcet's syndrome the same as Behçet's syndrome?

The term most likely refers to Behçet's syndrome and reflects a misspelling of Hulusi Behçet, the doctor after whom the condition was named. A medical record using an unfamiliar name should be clarified with the service that wrote it.

Is Behçet's syndrome contagious?

No. It is an inflammatory immune condition, not an infection. It cannot be caught through touch, kissing, sex, shared utensils, or living with someone.

Can Behçet's syndrome be cured?

There is currently no cure that permanently removes the condition. Treatment can control inflammation, shorten or prevent some flares, and lower the risk of organ damage. Some people have long periods with few or no symptoms.

Is Behçet's syndrome inherited?

It is not usually inherited in a simple way. Certain genes can raise susceptibility, but genes alone do not determine whether someone develops it. Most children of an affected parent do not develop the condition.

What foods should I avoid with Behçet's syndrome?

No standard diet has been shown to control the underlying condition. Spicy, acidic, salty, or hard foods may make active mouth ulcers more painful. Individual triggers can be discussed without unnecessarily restricting a balanced diet.

Does Behçet's syndrome affect life expectancy?

Many people have a normal lifespan, especially when inflammation is limited to ulcers, skin, or joints. Major blood-vessel, neurological, or severe eye involvement carries greater risk and needs close specialist treatment. Individual outlook is shaped mainly by the organs affected and response to treatment.

Sources



- National Institute of Arthritis and Musculoskeletal and Skin Diseases. Behçet's Disease . US National Institutes of Health.
- NHS. Behçet's disease . Overview of symptoms, diagnosis, treatment, and urgent complications.
- Hatemi G, Christensen R, Bang D, et al. "2018 update of the EULAR recommendations for the management of Behçet's syndrome." Annals of the Rheumatic Diseases . 2018;77(6):808-818.
- International Team for the Revision of the International Criteria for Behçet's Disease. "The International Criteria for Behçet's Disease: a collaborative study of 27 countries." Journal of the European Academy of Dermatology and Venereology . 2014;28(3):338-347.
- Yazici H, Seyahi E, Hatemi G, Yazici Y. "Behçet syndrome: a contemporary view." Nature Reviews Rheumatology . 2018;14(2):107-119.
- Nair JR, Moots RJ. "Behçet's disease." Clinical Medicine . 2017;17(1):71-77.