CONDITION

Hereditary urea cycle abnormality

An inherited condition that reduces the body's ability to remove ammonia made during protein breakdown.

WHAT IT IS
Inherited failure to remove ammonia from protein breakdown

BODY SYSTEM
Liver metabolism and nervous system

MAIN SYMPTOMS
Vomiting, unusual drowsiness, confusion, seizures

WHAT CAUSES IT
Inherited changes in urea-cycle genes

HOW IT IS DIAGNOSED
Blood ammonia, amino acids, urine tests, genetic confirmation

MAIN TREATMENT
Specialist diet, ammonia-lowering medicines, emergency metabolic care

CAN IT BE CURED?
Sometimes - liver transplant corrects urea-cycle function

IS IT AN EMERGENCY?
Sometimes - confusion or unusual drowsiness may mean high ammonia

LONG-TERM OUTLOOK
Good control is possible, but episodes may cause brain injury

Overview

A hereditary urea cycle abnormality is an inherited condition affecting how the body handles nitrogen from protein. It is also called a urea cycle disorder.

When you eat protein, your body breaks it into amino acids. This process produces nitrogen, which can form ammonia. Ammonia is toxic at high levels, especially to the brain. The liver normally changes ammonia into urea through a series of chemical steps called the urea cycle. Urea then leaves the body in urine.

In this condition, one of the enzymes or transport proteins needed for that cycle does not work properly. Ammonia may build up in the blood. This is called hyperammonaemia or hyperammonemia .

Some abnormalities cause severe symptoms soon after birth. Others leave part of the cycle working, so symptoms may not appear until childhood or adulthood. A person can seem well between episodes.

High ammonia needs prompt treatment because it can affect brain function. However, ongoing specialist care, nutrition planning, medicines and an emergency plan can reduce the risk of episodes.

Symptoms

Symptoms differ according to which part of the urea cycle is affected and how much function remains. They can also change quickly when ammonia rises.

A newborn with a severe abnormality may initially appear healthy. Within the first few days, feeding may become difficult. The baby may vomit, become unusually sleepy, breathe rapidly or have poor muscle tone. These symptoms can look like a serious infection, so urgent testing is important.

Later-onset symptoms may be less obvious. They can include:

Children may have feeding difficulties, slow growth or developmental concerns. School performance, attention or planning skills may be affected, particularly after episodes of high ammonia.

Some people first develop symptoms during a physical stress such as infection, prolonged fasting, surgery, childbirth or a sudden large increase in protein intake. Others have milder ongoing symptoms before the condition is recognised.

Arginase deficiency, one specific urea cycle abnormality, often has a different pattern. Progressive stiffness in the legs, walking difficulties and developmental problems may be more prominent than sudden severe ammonia episodes.

Symptoms alone cannot confirm that ammonia is high. Similar symptoms have many possible causes.

- Repeated vomiting, especially during illness
- Unusual sleepiness, confusion or changes in behaviour
- Headaches, poor concentration or problems with coordination
- Poor appetite or a strong dislike of protein-rich foods
- Seizures or reduced consciousness during a severe episode

Causes and risk factors

Hereditary urea cycle abnormalities are caused by changes in genes that provide instructions for urea-cycle enzymes or transport proteins. The specific condition is often named after the affected protein, such as ornithine transcarbamylase deficiency, citrullinaemia type 1 or argininosuccinic aciduria.

Most types follow an autosomal recessive inheritance pattern. This usually means a person has inherited one altered copy of the gene from each parent. The parents are often healthy carriers.

Ornithine transcarbamylase deficiency usually follows an X-linked pattern because its gene is on the X chromosome. It may be severe in males, who usually have one X chromosome. Females with an altered gene can also develop symptoms, ranging from mild to severe. This is partly because cells do not all use the same X chromosome.

A new genetic change can occasionally occur in someone without a known family history. Therefore, the absence of affected relatives does not rule out the condition.

Factors that may trigger high ammonia in someone with a urea cycle abnormality include:

Do not stop a prescribed medicine without medical advice. A metabolic specialist can help review medicines before surgery, pregnancy or other periods of increased risk.

- Fever, infection or another major illness
- Vomiting, fasting or not taking in enough energy
- Surgery, injury or childbirth
- Rapid tissue breakdown or intense physical stress
- A sudden high protein intake or concentrated protein supplements
- Certain medicines, including valproate, in susceptible people

Diagnosis

If high ammonia is suspected, a blood ammonia test is needed promptly. The sample must be collected and handled carefully because delays or contamination can produce an inaccurate result. An abnormal result is usually checked while treatment and further testing begin.

Blood tests measure amino acids, blood sugar, salts, liver function and the body's acid balance. A urine test for orotic acid , a substance related to the urea cycle, can help distinguish between different abnormalities.

Genetic testing may identify the altered gene and confirm the specific type. It can also help with testing relatives and understanding inheritance. If genetic testing is unclear, specialised enzyme testing or other biochemical tests may sometimes be needed.

Newborn screening can identify certain urea cycle abnormalities, including some cases of citrullinaemia and argininosuccinic aciduria. Screening programmes differ by location, and not every type is detectable. A normal newborn screening result does not rule out all urea cycle conditions.

Testing is usually coordinated by a metabolic specialist. If a genetic cause is confirmed, close relatives may be offered targeted testing even if they feel well.

Treatment and management

Treatment has two goals: lowering ammonia during an acute episode and preventing future rises.

A suspected ammonia crisis is treated in hospital. Care may include giving glucose and sometimes fats to provide energy and reduce the breakdown of body protein. Protein may be paused briefly under specialist supervision, then restarted carefully because the body still needs it.

Medicines called nitrogen scavengers provide other ways for the body to remove nitrogen. Arginine, citrulline or other supplements may be used for particular urea cycle abnormalities. Treatment is chosen according to the affected step, laboratory results and clinical situation.

If ammonia is dangerously high or is not falling quickly enough, dialysis may be needed. Dialysis filters ammonia directly from the blood. The team also treats any infection or other trigger.

Long-term care may involve:

A low-protein plan does not mean avoiding all protein. Too little protein can cause poor growth, muscle loss and other nutritional problems. Dietary changes should be planned with a metabolic dietitian.

Liver transplantation can provide working urea-cycle enzymes and prevent further metabolic crises. It is major surgery and requires lifelong anti-rejection medicine. It cannot reverse brain injury that occurred before transplantation, so its possible benefits and risks need careful specialist assessment.

- A personalised protein intake that supports growth and nutrition
- Special medical foods or essential amino acid mixtures when needed

- Regular ammonia-lowering medicines
- Vitamin, mineral and amino acid monitoring
- A written emergency plan for illness, vomiting or reduced food intake
- Follow-up with metabolic, dietetic and other specialist services

Complications

The main complication is injury to the brain from high ammonia. A severe rise can cause brain swelling, seizures, coma and, in some cases, death. Fast recognition and treatment reduce this risk but cannot guarantee that injury will be avoided.

Repeated or prolonged episodes may affect memory, attention, coordination, learning or behaviour. Effects can occur even when a person appears to recover fully after an episode.

Some specific abnormalities have additional complications. Argininosuccinic aciduria can be associated with liver problems and brittle hair. Arginase deficiency can cause increasing muscle stiffness and difficulty walking.

Long-term dietary restriction can lead to poor growth or nutrient deficiencies if it is not monitored carefully. Medicines can also cause side effects. Regular blood tests and diet reviews are therefore part of routine follow-up.

Prevention

The inherited genetic change itself cannot be prevented.

Once the condition is known, many ammonia episodes may be preventable or made less severe. This usually involves following the agreed nutrition and medicine plan, avoiding prolonged fasting, and using a specialist emergency plan during illness. Families and caregivers may need copies of that plan for school, childcare or travel.

Before planned surgery, pregnancy or childbirth, the metabolic team can arrange extra monitoring and a plan to prevent the body from breaking down its own protein.

Genetic counselling can explain how the specific abnormality is inherited. It can also discuss carrier testing, testing during pregnancy and reproductive options. These are personal choices rather than requirements.

Living with it and what to expect

The course over months and years is strongly influenced by the specific genetic abnormality, how much enzyme function remains and whether previous ammonia episodes caused injury.

Some people with a partial abnormality remain well for long periods and continue education, work, relationships and family life. They may still need daily treatment and extra planning around illness, travel, exercise or medical procedures.

Children with severe early-onset conditions may need ongoing help with feeding, development, movement or learning. Their care may involve metabolic specialists, dietitians, neurologists, therapists and school support.

Needs often change with growth, puberty, pregnancy, ageing or other health conditions. Protein and medicine plans may need adjustment over time. Routine follow-up usually includes reviewing growth or weight, nutrition, ammonia control, medicines and cognitive or developmental concerns.

Knowing early symptoms and having an emergency letter can make day-to-day life more manageable. It is also important not to treat every headache or stomach upset as an ammonia crisis. The agreed plan should explain which symptoms require testing or emergency care.

When to get medical help

Seek emergency medical care if someone with a known or suspected urea cycle abnormality develops:

Tell emergency staff about the risk of hyperammonemia and show them the metabolic emergency plan if one is available. A person with altered consciousness should not be given food, drink or oral medicine because of the risk of choking.

If you have a diagnosed condition, contact your metabolic team promptly during fever, vomiting, poor intake or before a planned operation. Follow the personalised illness plan rather than making major protein or medicine changes yourself.

Routine appointments are appropriate for stable concerns such as diet questions, medicine side effects, growth changes, school difficulties or planning for pregnancy and travel.

- New confusion, extreme sleepiness or unusual behaviour
- Repeated vomiting with worsening alertness
- A seizure, collapse or reduced consciousness
- Trouble breathing or severe loss of coordination

Common questions

Can a urea cycle abnormality first appear in an adult?

Yes. The genetic change is present from birth, but partial enzyme function may prevent symptoms for years. Illness, fasting, surgery, childbirth or another physical stress can reveal the condition later.

Can I eat protein if I have this condition?

Usually, yes. The body needs protein for muscles, growth and tissue repair. The safe amount and type must be personalised. A zero-protein diet is not a safe long-term treatment.

Does newborn screening find every urea cycle abnormality?

No. Screening detects some types more reliably than others, and programmes differ between regions. Ornithine transcarbamylase deficiency is not reliably found through standard newborn screening.

Can ammonia be high when someone looks well?

It can be mildly raised with few obvious symptoms. However, ammonia levels can also increase quickly. Regular monitoring and an illness plan are important for people with a confirmed condition.

Does a liver transplant cure it?

A successful transplant can correct the liver's inability to run the urea cycle and prevent new ammonia crises. It does not undo established brain injury and brings major surgical and lifelong treatment risks.

Could my siblings or children have it?

They may be at risk, but the chance depends on the specific gene and inheritance pattern. Genetic testing and counselling can clarify which relatives may benefit from testing.

Sources

- Ah Mew N, Simpson KL, Gropman AL, Lanpher BC, Chapman KA, Summar ML. Urea Cycle Disorders Overview . GeneReviews®, NCBI Bookshelf
- Häberle J, Burlina A, Chakrapani A, et al. Suggested guidelines for the diagnosis and management of urea cycle disorders: First revision . Journal of Inherited Metabolic Disease. 2019;42:1192-1230



- Lichter-Konecki U, Caldovic L, Morizono H, Simpson K. Ornithine Transcarbamylase Deficiency . GeneReviews®, NCBI Bookshelf
- British Inherited Metabolic Disease Group. Urea cycle disorders: emergency management guidance
- US Food and Drug Administration. Ravicti (glycerol phenylbutyrate) prescribing information