CONDITION

Hypervitaminosis A

Hypervitaminosis A is a harmful buildup of preformed vitamin A in the body.

WHAT IT IS
Harmful buildup of preformed vitamin A

BODY SYSTEM
Liver, skin, bones, brain, and nervous system

MAIN SYMPTOMS
Headache, nausea, dry skin, bone pain

WHAT CAUSES IT
Excess vitamin A from supplements, medicines, or food

HOW IT IS DIAGNOSED
Exposure history, examination, and blood tests

MAIN TREATMENT
Stopping excess vitamin A and supportive care

CAN IT BE CURED?
Yes - often resolves after excess stops

IS IT AN EMERGENCY?
Sometimes - severe headache, confusion, or repeated vomiting

LONG-TERM OUTLOOK
Usually improves; liver or bone damage may persist

Overview

Hypervitaminosis A means there is enough vitamin A in the body to cause harm. It can happen after one very large exposure, called acute toxicity, or after taking too much over weeks, months, or years, called chronic toxicity.

Vitamin A supports vision, immune function, growth, and healthy skin. The body stores it mainly in the liver. This storage is useful when intake is low, but it also means excessive amounts can build up.

Hypervitaminosis A is mainly caused by preformed vitamin A , such as retinol and retinyl esters. These forms are found in some supplements, prescription retinoid medicines, liver, and fish liver oils.

Beta-carotene is different. The body converts it to vitamin A only as needed, so beta-carotene

from food does not usually cause hypervitaminosis A. Large amounts can turn the skin yellow-orange, but this is a separate condition called carotenemia.

Most people improve after the excess source is removed. Recovery time depends on how much was taken, how long exposure lasted, and whether the liver, bones, or other organs were affected.

Symptoms

Symptoms differ between acute and chronic toxicity. They can also resemble many other conditions, so symptoms alone cannot confirm hypervitaminosis A.

Acute toxicity may cause symptoms within hours or days of a very large exposure:

Chronic toxicity tends to develop gradually. Possible symptoms include:

Liver injury may cause pain under the right ribs, yellow skin or eyes, dark urine, or swelling. Increased pressure around the brain can cause a persistent headache, vomiting, vision changes, or a pulsing sound in the ears.

In infants, signs may include irritability, poor feeding, vomiting, or a bulging soft spot on the head. These symptoms need prompt medical assessment.

- Severe headache
- Nausea or repeated vomiting
- Dizziness, blurred vision, or unusual sleepiness
- Irritability or confusion
- Poor coordination
- Skin peeling, sometimes appearing later
- Dry, rough, itchy, or peeling skin
- Cracked lips
- Hair thinning or hair loss
- Ongoing headache
- Tiredness or weakness
- Reduced appetite or unintended weight loss
- Bone, muscle, or joint pain
- Nausea or abdominal discomfort

Causes and risk factors

The usual cause is taking more preformed vitamin A than the body can safely handle. This may come from:

Labels may list vitamin A in different units and forms. A multivitamin, a separate vitamin A product, and another supplement marketed for skin, vision, or immunity may all contain it. This "stacking" can make the total difficult to recognise.

Prescription retinoids are related to vitamin A and can cause similar harmful effects. Their risks differ from those of ordinary supplements. Follow the monitoring and pregnancy-prevention instructions provided with the medicine.

People with liver or kidney problems may be more vulnerable to harmful effects. Regular heavy alcohol use can also add to liver risk. Children can become unwell after smaller absolute amounts than adults because of their size.

During pregnancy, excessive preformed vitamin A and certain retinoid medicines can harm fetal development. This risk is important even when the pregnant person does not feel unwell.

- High-strength vitamin A supplements
- Several supplements containing vitamin A taken together
- Prescription retinoid medicines, such as oral acne or psoriasis treatments
- Frequent large servings of liver
- Fish liver oil or other concentrated animal-liver products
- Accidental swallowing of supplements, especially by a child

Diagnosis

There is no single blood result that always confirms hypervitaminosis A. Diagnosis usually combines your symptoms, an examination, and a detailed history of supplements, medicines, and foods.

Bring product containers or clear photographs of the labels to an appointment. The exact form of vitamin A, strength, serving size, and length of use can be important.

Blood tests may check:

A normal blood retinol level does not always rule out toxicity. The body regulates this level closely, and it may not reflect how much vitamin A is stored in the liver. Other vitamin A measurements can sometimes support the diagnosis, but they must be interpreted alongside

the exposure history.

Further tests depend on the symptoms. An eye examination may look for swelling of the optic nerve caused by pressure around the brain. Bone-density testing or imaging may be considered after prolonged exposure or fractures. Pregnancy testing may be relevant when a retinoid medicine or high-dose exposure is involved.

- Vitamin A levels
- Liver function
- Calcium levels
- Kidney function
- Cholesterol and triglycerides

Treatment and management

The main treatment is removing the excess source of vitamin A. This often means stopping non-prescription vitamin A products and avoiding other concentrated sources while the situation is assessed.

If the suspected source is a prescription retinoid, contact the prescriber promptly rather than making a long-term medication change without advice. Severe symptoms require urgent care instead of waiting for a routine appointment.

Treatment may also include:

Severe acute toxicity, significant liver injury, or neurological symptoms may require hospital care. There is no routine antidote that immediately removes stored vitamin A. The body needs time to process and clear the excess.

Do not try to "flush out" vitamin A with extreme diets or other supplements. Some products may add further liver stress or cause new nutrient problems.

- Fluids and medicines for nausea or pain
- Blood-test monitoring
- Care for liver injury or high calcium
- Treatment for increased pressure around the brain
- Nutrition advice to maintain adequate, but not excessive, vitamin A intake

Complications

Long-lasting or severe excess can affect several parts of the body.

Possible complications include liver inflammation, scarring, and reduced liver function. Chronic toxicity may weaken bones and increase fracture risk. High calcium levels can affect the kidneys, digestive system, and alertness.

Pressure around the brain, sometimes called intracranial hypertension, can threaten vision if it is not treated. Very large acute exposures can cause serious neurological or liver problems, although this is not the usual course.

Exposure during pregnancy can cause major birth defects. Some prescription retinoids remain in the body after treatment ends, so their specific pregnancy-prevention instructions must be followed.

Prevention

Hypervitaminosis A is usually preventable.

Check the labels of all supplements rather than considering each product separately. Look for retinol, retinyl acetate, retinyl palmitate, fish liver oil, and other forms of preformed vitamin A. Avoid combining vitamin A products unless a qualified health professional has recommended and reviewed the total amount.

Use supplements only as directed. More vitamin A does not improve vision, skin, or immunity once the body has enough. Keep concentrated supplements away from children.

Liver is highly nutritious but can contain large amounts of preformed vitamin A. Regular large servings may create a risk, particularly when combined with supplements.

If you are pregnant, could become pregnant, or are breastfeeding, check supplements and retinoid medicines with a pharmacist or relevant clinician. Do not assume that a product is safe during pregnancy because it is labelled "natural."

You still need some vitamin A. Prevention does not mean avoiding vegetables containing beta-carotene or removing all vitamin A from your diet.

Living with it and what to expect

After the excess source is stopped, nausea and headache may improve within days. Skin, hair, and energy changes may take weeks or months. Vitamin A stored in the liver can take time to fall.

Follow-up may include repeat liver, calcium, or vitamin A testing. People with prolonged exposure may need bone assessment or longer monitoring. If a medicine caused the problem, future treatment choices may need to account for that reaction.

Many people recover without lasting effects, especially when the condition is recognised early. Liver scarring, weakened bones, or vision damage may not fully reverse if they were advanced before treatment.

Once recovery is complete, most people can eat a balanced diet containing ordinary food sources of vitamin A. Ongoing severe restriction is not usually needed unless a clinician identifies another reason for it.

When to get medical help

Seek emergency help for:

Contact a poison information service or emergency service after a suspected large accidental exposure. Do not induce vomiting unless specifically instructed.

Get urgent medical advice for persistent vomiting, jaundice, dark urine, significant abdominal pain, or a new persistent headache while taking a retinoid medicine. Also seek prompt advice if excessive vitamin A or a retinoid may have been used during pregnancy.

A routine appointment is appropriate for gradual symptoms such as dry skin, hair loss, fatigue, or bone pain when you regularly use vitamin A supplements. Bring all supplement and medicine labels for review.

- A severe or rapidly worsening headache with vomiting or vision changes
- Confusion, collapse, a seizure, or difficulty waking
- Suspected swallowing of a large amount by a child
- A very large exposure followed by significant symptoms

Common questions

Can you get hypervitaminosis A from food?

Yes, but it is uncommon with a typical varied diet. Repeated large servings of animal liver can provide very high amounts of preformed vitamin A. Vegetables rich in beta-carotene do not usually cause vitamin A toxicity.

How much vitamin A is too much?

The safe upper amount differs by age, pregnancy status, and the form of vitamin A. It is not a precise toxicity threshold, and some prescription retinoids are assessed differently. Check the total from every supplement rather than relying on one product's serving information.

Does beta-carotene cause vitamin A toxicity?

Beta-carotene from food does not usually cause hypervitaminosis A because the body limits how much it converts. Large amounts may cause harmless yellow-orange skin colouring. High-dose beta-carotene supplements have other health considerations and are not suitable for everyone.

How long does vitamin A toxicity last?

Acute symptoms may improve over days after the exposure ends. Chronic symptoms can take weeks or months because vitamin A is stored in the liver. Recovery may be longer when the liver, bones, or nervous system have been affected.

Should I stop a vitamin A supplement if I have symptoms?

Do not take further non-prescription high-dose vitamin A until you have obtained medical advice. If vitamin A or a retinoid was prescribed, contact the prescriber promptly. Use urgent or emergency care for severe headache, repeated vomiting, confusion, or vision changes.

Is hypervitaminosis A fatal?

Most recognised cases improve after the excess source is removed. However, an extremely large exposure or severe liver or brain involvement can be life-threatening. A suspected large ingestion should be discussed immediately with a poison information service or emergency service.

Sources

- National Institutes of Health, Office of Dietary Supplements. Vitamin A and Carotenoids: Fact Sheet for Health Professionals
- National Institute of Diabetes and Digestive and Kidney Diseases. LiverTox: Vitamin A
- Merck Manual Professional Edition. Vitamin A Toxicity
- Penniston KL, Tanumihardjo SA. "The Acute and Chronic Toxic Effects of Vitamin A." American Journal of Clinical Nutrition . 2006;83(2):191-201
- European Food Safety Authority Panel on Nutrition, Novel Foods and Food Allergens. "Scientific Opinion on the Tolerable Upper Intake Level for Preformed Vitamin A and -Carotene." EFSA Journal . 2024