CONDITION

Hypomelanosis of ito

Hypomelanosis of Ito is a mosaic condition that causes pale streaks or whorls on the skin and may affect other parts of the body.

WHAT IT IS
Mosaic skin condition causing pale streaks and whorls

BODY SYSTEM
Skin; sometimes nervous system, bones, muscles, eyes, and teeth

MAIN SYMPTOMS
Pale skin patterns, seizures, developmental differences, body asymmetry

WHAT CAUSES IT
Genetic mosaicism, often with no identified change

HOW IT IS DIAGNOSED
Skin examination, sometimes Wood's lamp and genetic testing

MAIN TREATMENT
Skin protection, camouflage, and treatment of associated problems

CAN IT BE CURED?
No - no established skin cure

IS IT AN EMERGENCY?
Sometimes - a first or prolonged seizure

LONG-TERM OUTLOOK
Usually stable skin; outlook reflects associated health problems

Overview

Hypomelanosis of Ito is a condition that causes areas of skin to contain less pigment, or melanin, than the surrounding skin. Melanin gives skin much of its colour. The lighter areas usually form streaks, swirls, or wave-like patterns across the body.

These patterns follow Blaschko lines . These are invisible pathways formed as skin cells spread during development before birth. They are not related to nerves, blood vessels, or anything that has touched the skin.

Hypomelanosis of Ito is better understood as a description of a pigment pattern than as one single disease. It is often grouped under the broader term pigmentary mosaicism . Mosaicism means that some cells have a genetic difference while other cells in the same person do not.

For some people, the skin pattern is the only feature. Others have differences affecting development, the nervous system, bones, muscles, eyes, or teeth. This is why a medical assessment may look beyond the skin, especially when the pattern is first recognised in a baby or child.

The condition is not contagious. It is not caused by poor hygiene, sun exposure, or anything a parent did during pregnancy.

Symptoms

The main sign is a flat area of skin that is lighter than the surrounding skin. The lighter colour may be present at birth or become noticeable during infancy or early childhood. It can be harder to see on very light skin and may become clearer after surrounding skin tans.

The pattern may:

The affected skin is usually not itchy, painful, inflamed, or scaly. Its texture generally feels the same as nearby skin.

Some people have features outside the skin. These are not present in everyone and can range from mild to significant. Possible findings include:

A pale skin pattern does not show how likely someone is to have these additional features. A person can have extensive skin markings without major health problems, while a smaller skin area does not rule them out.

- Form streaks, whorls, waves, or patches
- Follow the length of an arm or leg
- Cross several areas of the trunk or limbs
- Appear on one side or both sides of the body
- Have an uneven boundary rather than a simple round shape
- Delayed speech, movement, learning, or other developmental skills
- Seizures
- Differences in muscle tone, coordination, or strength
- Intellectual disability
- Differences in limb length or body symmetry
- Curvature of the spine, called scoliosis

- Eye, vision, dental, hair, or nail differences

Causes and risk factors

Hypomelanosis of Ito usually results from genetic mosaicism. A genetic change happens in one cell after an embryo has started developing. As that cell divides, its descendants carry the change while other cells do not. The affected skin cells then spread along Blaschko lines.

Many different chromosome or gene changes can produce this pattern. In some people, testing finds a specific change. In others, no cause is identified. A negative blood test does not always exclude mosaicism because the change may be present only in the affected skin or another tissue.

Hypomelanosis of Ito is usually sporadic, meaning it occurs without a family history. It is not generally considered one inherited disorder with a single inheritance pattern. However, an identified genetic or chromosome finding may have implications for relatives or future pregnancies.

There are no known lifestyle factors that cause this condition. Skin colour, diet, medicines, and routine activities do not create the mosaic pattern.

Diagnosis

Diagnosis begins with an examination of the skin. A dermatologist may recognise the typical pale streaks or whorls following Blaschko lines. The clinician will also ask when the pattern appeared and whether it was preceded by redness, blisters, scaling, or another rash.

A Wood's lamp may be used. This handheld device shines ultraviolet light on the skin and can make differences in pigmentation easier to see. It does not hurt and does not involve taking a sample.

The assessment may include checks of development, movement, muscle tone, spine alignment, limb symmetry, vision, hearing, teeth, and neurological function. Which checks are needed is guided by age, symptoms, and examination findings.

Genetic testing may be considered when there are developmental, neurological, skeletal, or other associated features. Testing can include a chromosome analysis or chromosomal microarray, which looks for missing or extra pieces of genetic material. Sometimes both a blood sample and a sample of affected skin are examined because mosaic changes can be missed in blood.

A skin biopsy is not always needed. If one is performed, a small sample is studied under a microscope. Findings may show reduced pigment, but they are not usually unique to hypomelanosis of Ito.

Other conditions can cause pale skin areas. These include vitiligo, a stable pale birthmark called nevus depigmentosus, tuberous sclerosis, and later-stage incontinentia pigmenti. The pattern, medical history, examination, and sometimes testing help distinguish them.

Treatment and management

The pale skin itself does not usually require medical treatment. There is no established treatment that reliably removes the mosaic pattern or restores lasting, even pigmentation.

Sun protection can reduce the contrast between affected and surrounding skin. Broad-spectrum sunscreen, protective clothing, and shade also protect against ordinary sun damage. The lighter areas may have less melanin, but the surrounding skin needs protection as well.

Cosmetic camouflage products or carefully chosen self-tanning products may make the colour difference less noticeable. A dermatologist or skin-camouflage service can help with colour matching. Some light-based or laser treatments have been described, but evidence is limited and results may not last.

Management mainly focuses on any associated health or developmental needs. This might involve:

Follow-up should be based on the features present rather than the skin pattern alone. A child with normal development and no other findings may need only routine observation. Someone with seizures, movement difficulties, or skeletal differences may need ongoing specialist care.

- Developmental, physical, occupational, or speech therapy
- Specialist treatment for seizures
- Eye and vision care
- Monitoring or treatment for scoliosis and limb differences
- Dental care for abnormal tooth development
- Educational support for learning differences

Complications

The skin pattern itself usually does not damage internal organs or develop into skin cancer. Its visibility can still affect confidence, body image, or social experiences, especially when it involves the face or large areas.

The more important possible complications come from associated neurological or structural differences. Seizures can affect safety and learning. Muscle, joint, limb, or spine differences may affect movement or cause discomfort. Developmental or learning differences may influence school support and future independence.

Not everyone develops these complications. Their presence and severity reflect the underlying mosaic change, not the amount of pale skin alone.

Prevention

Hypomelanosis of Ito cannot currently be prevented. The mosaic change usually happens by chance during early development and is not caused by an action taken before or during pregnancy.

There is no diet, supplement, skin product, or behaviour known to prevent the condition. Sun protection can reduce tanning contrast and sun damage, but it cannot prevent or remove the underlying pattern.

If genetic testing identifies a specific change, genetic counselling can explain what it may mean for the individual and their family. Without an identified change, inheritance and recurrence cannot be predicted from the skin pattern alone.

Living with it and what to expect

The skin markings are usually long-lasting, but their appearance can change. They may become less obvious as a child grows or as skin colour changes with age. Seasonal tanning can make the contrast stronger. This does not necessarily mean that the condition is spreading or worsening.

The pale areas generally do not become painful or inflamed. New redness, scaling, itching, or rapid colour loss may have another cause and should not automatically be attributed to hypomelanosis of Ito.

When the condition is limited to the skin, it usually does not restrict everyday activities. People can attend school, work, exercise, swim, and take part in other usual activities. Cosmetic camouflage is a personal choice, not a medical requirement.

Associated neurological or developmental features may become clearer as a child grows and more complex skills are expected. Early support can help with communication, movement, learning, and daily activities. In adulthood, care often shifts toward managing established needs rather than monitoring the skin.

Long-term outlook is closely linked to any associated brain, developmental, or skeletal differences. Some people need little or no ongoing care. Others need continuing support for seizures, mobility, learning, or independent living.

When to get medical help

Arrange a routine medical appointment if you or your child has an unexplained pale streaked or whorled skin pattern. An assessment is particularly useful when the markings appeared in infancy or are accompanied by developmental, movement, vision, dental, or skeletal differences.

Seek prompt medical advice for:

A first suspected seizure needs urgent medical assessment. Call emergency services if a seizure lasts five minutes or longer, another seizure begins before recovery, breathing is difficult, a serious injury occurs, or the person does not return to their usual level of awareness.

Sudden weakness, severe confusion, loss of consciousness, or breathing difficulty also needs emergency care. These symptoms should not be assumed to be part of the skin condition.

- Loss of previously gained developmental skills
- New episodes of staring, jerking, collapse, or altered awareness
- Increasing weakness, balance problems, or difficulty walking
- A noticeable change in spine shape or limb symmetry
- New pain, inflammation, scaling, or sores in the pale skin

Common questions

Is hypomelanosis of Ito the same as vitiligo?

No. Vitiligo is an acquired condition in which pigment-producing cells are lost, often creating sharply defined white patches that may enlarge over time. Hypomelanosis of Ito usually begins at birth or in early childhood and follows Blaschko lines. A skin examination can help tell them apart.

Will the pale patches spread?

The underlying mosaic pattern is usually established before birth and does not spread like an infection. It may appear to change as the body grows or as surrounding skin tans. Rapidly developing or expanding pale areas should be assessed because another pigment condition may be responsible.

Does everyone with hypomelanosis of Ito have brain problems?

No. Some people have only the skin pattern. Others have seizures, developmental differences, or other neurological findings. The size or brightness of the pale area cannot reliably predict whether the nervous system is involved.

Can the normal skin colour come back?

There is no established treatment that reliably creates permanent, even repigmentation. The contrast may become less noticeable with age or changes in surrounding skin colour. Camouflage products can help if the appearance is a concern.

Is hypomelanosis of Ito inherited?

It is usually caused by mosaicism that develops after conception rather than by a genetic change inherited in every cell. However, some identified chromosome or gene findings may affect genetic counselling. Testing results and family history are needed to discuss inheritance accurately.

Can someone with hypomelanosis of Ito have a normal life?

Many people whose condition affects only the skin take part in usual activities without medical restrictions. When seizures, developmental differences, or skeletal problems are present, support needs can be greater. The skin pattern alone does not determine health, abilities, or life expectancy.

Sources

- Kromann AB, Ousager LB, Ali IKM, Aydemir N, Bygum A. "Pigmentary mosaicism: a review of original literature and recommendations for future handling." Orphanet Journal of Rare Diseases . 2018;13:39.
- Taibjee SM, Bennett DC, Moss C. "Abnormal pigmentation in hypomelanosis of Ito and pigmentary mosaicism: the role of pigmentary genes." British Journal of Dermatology . 2004;151(2):269-282.
- Sybert VP. "Hypomelanosis of Ito: a description, not a diagnosis." Journal of Investigative Dermatology . 1994;103(5 Supplement):141S-143S.



- Ruiz-Maldonado R, Toussaint S, Tamayo L, Laterza A, del Castillo V. "Hypomelanosis of Ito: diagnostic criteria and report of 41 cases." Pediatric Dermatology . 1992;9(1):1-10.
- Orphanet. "Hypomelanosis of Ito." ORPHA:3447.
- DermNet New Zealand. "Hypomelanosis of Ito." Dermatology reference article.