CONDITION

Immune hemolytic anemia

Immune hemolytic anemia occurs when the immune system destroys red blood cells faster than the body can replace them.

WHAT IT IS
Antibody-driven destruction of red blood cells

BODY SYSTEM
Blood and immune system

MAIN SYMPTOMS
Fatigue, breathlessness, jaundice, dark urine

WHAT CAUSES IT
Immune misdirection, sometimes linked to illness or medicines

HOW IT IS DIAGNOSED
Blood counts, hemolysis tests, direct antiglobulin test

MAIN TREATMENT
Cause-specific immune treatment, supportive care, sometimes transfusion

CAN IT BE CURED?
Sometimes - trigger removal or lasting remission

IS IT AN EMERGENCY?
Sometimes - chest pain, fainting, or breathlessness at rest

LONG-TERM OUTLOOK
Often manageable, but relapses and chronic forms occur

Overview

Immune hemolytic anemia is a condition in which the immune system destroys red blood cells too early. This process is called hemolysis . Anemia develops when the bone marrow cannot make replacement cells quickly enough.

Red blood cells carry oxygen around the body. Losing too many can cause tiredness, weakness, breathlessness, or a rapid heartbeat. Their breakdown can also release pigments that cause yellow skin or eyes, known as jaundice, and dark urine.

In many cases, the immune system makes proteins called antibodies that attach to red blood cells by mistake. The spleen, liver, or bloodstream then removes or destroys the affected cells.

The most common autoimmune forms are classified by how the antibodies behave:

Immune hemolysis can also follow certain medicines, a blood transfusion, or transfer of antibodies during pregnancy. These situations have different causes and treatments.

The condition ranges from mild and stable to severe and fast-moving. Its importance depends on how quickly red blood cells are being lost, the hemoglobin level, your symptoms, and whether another illness is involved.

- Warm autoimmune hemolytic anemia involves antibodies that react most strongly near normal body temperature
- Cold autoimmune hemolytic anemia involves antibodies that become more active at lower temperatures
- Mixed autoimmune hemolytic anemia has features of both

Symptoms

Symptoms may appear slowly or develop over several days. Mild anemia may be found during a blood test before you notice anything unusual.

Possible symptoms include:

Cold-antibody forms can also cause painful, numb, or blue fingers, toes, ears, or nose after cold exposure. This happens when blood flow through small vessels is disturbed. Symptoms may become worse during cold weather, fever, or infection.

These symptoms are not specific to immune hemolytic anemia. Other types of anemia, liver conditions, bleeding, and several common illnesses can cause similar problems.

- Unusual tiredness or weakness
- Shortness of breath, especially during activity
- Dizziness or headache
- A fast or pounding heartbeat
- Pale skin
- Yellow skin or whites of the eyes
- Tea-colored, brown, or red urine
- Discomfort or fullness under the left ribs from an enlarged spleen

Causes and risk factors

Autoimmune hemolytic anemia happens when the immune system mistakes red blood cells for a target. Sometimes no underlying cause is found. This is called primary autoimmune hemolytic anemia.

In other cases, it is associated with another condition, such as:

Some medicines can cause immune hemolysis. The reaction may involve antibodies against the medicine, the red blood cell, or both. Do not stop a prescribed medicine based only on symptoms or a test result. A clinician should assess whether the medicine is responsible and choose a safe alternative if needed.

Cold autoimmune hemolysis may follow an infection and then improve as the infection resolves. Chronic cold agglutinin disease is often linked to an abnormal group of antibody-producing cells in the bone marrow, even when there is no obvious lymphoma.

Alloimmune hemolytic anemia is different from an autoimmune form. It occurs when antibodies react to red blood cells from another person, such as after an incompatible transfusion or during pregnancy. Blood services reduce this risk by testing and matching donated blood.

- An autoimmune condition, including lupus
- A blood or lymph system condition, such as chronic lymphocytic leukemia or lymphoma
- Certain infections
- An immune system disorder

Diagnosis

Diagnosis requires evidence that red blood cells are being destroyed, along with testing for an immune cause. A single abnormal result does not always confirm the condition.

A complete blood count measures hemoglobin and the number and size of blood cells. A reticulocyte count measures young red blood cells. It is often raised because the bone marrow is trying to replace those being lost, although this response can sometimes be delayed or inadequate.

Other blood tests look for signs of hemolysis:

The direct antiglobulin test , also called the direct Coombs test, checks whether antibodies or immune proteins called complement are attached to red blood cells. The result can help distinguish warm and cold forms. A positive result alone does not prove active hemolysis, and a small number of immune cases have a negative result.

Further tests may look for cold-reacting antibodies, medicine-related antibodies, infections,

autoimmune conditions, or blood and lymph disorders. Bone marrow testing is not needed for everyone, but it may be used when blood production is unexpectedly low or an underlying bone marrow condition is suspected.

- Increased bilirubin, a pigment released when red blood cells break down
- Increased lactate dehydrogenase, or LDH, released from damaged cells
- Reduced haptoglobin, a protein that binds free hemoglobin
- Changes in red blood cell shape on a blood film

Treatment and management

Treatment is based on the type of antibody, the severity of anemia, your symptoms, and any underlying cause. Mild, stable cases may initially need monitoring rather than immediate treatment.

For warm autoimmune hemolytic anemia, corticosteroids like prednisone or hydrocortisone are commonly used to reduce immune destruction. If the condition does not respond, returns, or requires ongoing steroid treatment, other options may include antibody-based treatment such as rituximab, additional immune-suppressing medicines, or removal of the spleen. Each option has different infection and treatment risks.

Cold autoimmune hemolytic anemia is managed differently. Protecting the whole body from cold can reduce symptoms, but it does not remove the underlying antibody. Treatment may target the abnormal immune cells making the antibody or block the complement pathway that damages red blood cells. Corticosteroids and spleen removal are generally less effective for primary cold agglutinin disease than for warm autoimmune hemolysis.

When a medicine is confirmed as the trigger, stopping it under medical supervision is usually central to management. An infection or related autoimmune, blood, or lymph condition may also need treatment.

Supportive care can include folic acid when red blood cell production is increased. A red blood cell transfusion may be needed for severe or symptomatic anemia. Finding perfectly matched blood can take longer because the antibodies may interfere with compatibility testing, but urgent transfusion can still be given when medically necessary.

Treatment aims to control hemolysis, improve hemoglobin, reduce symptoms, and limit treatment side effects. A normal antibody test is not always required for someone to be in remission.

Complications

Rapid or severe anemia can reduce the oxygen reaching the heart, brain, and other organs. This may cause fainting, chest pain, marked breathlessness, or heart strain, especially in people who already have heart or lung conditions.

Other possible complications include:

Treatments can also have risks. Immune-suppressing medicines may increase infection risk. Long-term corticosteroid use can affect bones, blood sugar, blood pressure, eyes, and mood. Spleen removal increases the risk of certain serious infections, so vaccination and infection precautions are important.

- Blood clots in the legs, lungs, or other blood vessels
- Gallstones caused by long-term red blood cell breakdown
- Kidney injury during severe destruction within the bloodstream
- An enlarged spleen
- Complications from an underlying infection or blood condition

Prevention

Most autoimmune hemolytic anemia cannot be prevented. It is not usually caused by diet, exercise, or something you did.

If a medicine is confirmed as the cause, avoiding that medicine in the future can prevent another reaction. Keep the medicine's name in your health records and tell clinicians and pharmacists about the reaction.

If you have cold-antibody disease, warm clothing and avoiding sudden or prolonged cold exposure may reduce symptoms. This manages triggers but does not prevent the condition itself.

Careful blood typing, antibody screening, and matching reduce the risk of transfusion-related immune hemolysis. People with known red blood cell antibodies should make sure this information remains in their medical record.

Living with it and what to expect

The course differs among immune hemolytic conditions. Warm autoimmune hemolytic anemia

may settle with treatment, return after a remission, or require longer-term management. Some people have only one episode, while others need repeat treatment.

Cold hemolysis following an infection may improve as the infection clears. Primary cold agglutinin disease is usually a long-term condition. Symptoms and anemia can become more noticeable during winter, infections, surgery, or other physical stress.

Follow-up often includes a complete blood count and tests for hemolysis. The timing is based on how active the condition is and whether you are receiving treatment. Report new symptoms rather than waiting only for the next scheduled test.

Activity may need to be reduced when anemia is active. As hemoglobin improves, many people can gradually return to usual work, exercise, and social activities. Cold-water swimming and other intense cold exposure may not be suitable for people with cold-antibody disease.

If your spleen has been removed, follow the vaccination and infection plan provided by your healthcare team. Fever after spleen removal needs prompt medical assessment because some infections can progress quickly.

When to get medical help

Arrange a routine or prompt appointment if you develop ongoing tiredness, mild breathlessness, yellowing of the eyes, dark urine, or reduced exercise tolerance. Contact your care team if known symptoms return after improving or if treatment side effects are affecting daily life.

Seek same-day medical advice for rapidly worsening weakness, a new fast heartbeat, much darker urine, increasing jaundice, or breathlessness with light activity. These changes may mean hemolysis or anemia is becoming more active.

Seek emergency care for:

Do not drive yourself if you feel faint, confused, or severely short of breath.

- Chest pain
- Fainting or being difficult to wake
- Severe breathlessness or breathlessness at rest
- Confusion
- Blue or grey lips
- Signs of a serious transfusion reaction, such as fever, chills, back pain, breathing difficulty, or dark urine during or soon after a transfusion

Common questions

Is immune hemolytic anemia the same as iron-deficiency anemia?

No. Iron-deficiency anemia usually happens when the body lacks enough iron to make red blood cells. Immune hemolytic anemia happens because red blood cells are destroyed too early. Iron supplements do not treat immune destruction unless iron deficiency is also present.

Can immune hemolytic anemia go away?

Sometimes. A medicine-related or infection-related episode may resolve after the trigger is removed. Autoimmune forms can enter remission but may return. Chronic cold agglutinin disease often needs long-term monitoring or treatment.

Does a positive Coombs test mean I have this condition?

Not by itself. A direct Coombs test shows that antibodies or complement are attached to red blood cells. Diagnosis also requires evidence of red blood cell destruction and consideration of symptoms, medicines, transfusions, pregnancy, and other conditions.

Is immune hemolytic anemia cancer?

No. It is an immune blood condition, not cancer. However, it can sometimes occur alongside a blood or lymph cancer. Finding an associated condition affects treatment, but many people with immune hemolysis do not have cancer.

Can I exercise with immune hemolytic anemia?

Light activity may be possible when symptoms are stable. More demanding exercise may worsen breathlessness, dizziness, or a racing heartbeat when anemia is active. Build activity gradually and stop if you develop chest pain, faintness, or unusual breathlessness.

Is it contagious?

No. Immune hemolytic anemia cannot spread from one person to another. Some infections that trigger hemolysis may be contagious, but the immune anemia itself is not.

Sources

- Hill QA, Stamps R, Massey E, Grainger JD, Provan D, Hill A. The diagnosis and management of primary autoimmune haemolytic anaemia . British Journal of Haematology . 2017;176(3):395-411.



- Jäger U, Barcellini W, Broome CM, et al. Diagnosis and treatment of autoimmune hemolytic anemia in adults: Recommendations from the First International Consensus Meeting . Blood Reviews . 2020;41:100648.
- Kuter DJ. Warm autoimmune hemolytic anemia and the best treatment strategies . Hematology: American Society of Hematology Education Program . 2022;2022(1):105-113.
- Berentsen S. How I treat cold agglutinin disease . Blood . 2021;137(10):1295-1303.
- National Library of Medicine. Hemolytic anemia . MedlinePlus Medical Encyclopedia . Accessed August 10, 2026.