CONDITION

Insecticide poisoning

Insecticide poisoning happens when exposure to an insect-killing chemical disrupts normal body functions.

WHAT IT IS
Harmful exposure to an insect-killing chemical

BODY SYSTEM
Nervous system, lungs, heart, skin, and digestive system

MAIN SYMPTOMS
Nausea, sweating, breathing trouble, confusion

WHAT CAUSES IT
Swallowing, breathing, or absorbing insecticide

HOW IT IS DIAGNOSED
Exposure history, examination, and sometimes blood tests

MAIN TREATMENT
Decontamination, supportive care, and specific antidotes

CAN IT BE CURED?
Sometimes - most recover after exposure stops

IS IT AN EMERGENCY?
Sometimes - breathing trouble, seizures, or severe confusion

LONG-TERM OUTLOOK
Usually good after mild exposure; severe poisoning may cause disability

Overview

Insecticide poisoning happens when an insect-killing chemical enters the body in a harmful amount. Exposure may occur by swallowing the product, breathing in spray or fumes, or getting it on the skin or in the eyes.

Insecticides include several chemical groups. Organophosphates and carbamates interfere with an enzyme needed for normal nerve function. Pyrethrins and pyrethroids can irritate the skin, eyes, and airways and may affect the nervous system in larger exposures. Other products have different effects.

Symptoms can begin within minutes, but some take hours to appear. The risk depends on the chemical, its concentration, the amount involved, the route of exposure, and how quickly it is removed from the body. A small skin splash is different from swallowing a concentrated

agricultural product.

Severe poisoning can disrupt breathing, heart function, or consciousness. Prompt advice from a poison information service can help determine whether monitoring at home is reasonable or emergency assessment is needed.

Symptoms

Symptoms differ between insecticides. The product label or container can provide important clues, so keep it available if this can be done safely.

Mild exposure may cause:

Organophosphate and carbamate poisoning can cause a group of symptoms called a cholinergic syndrome . This means certain nerves are overstimulated. Signs may include very small pupils, blurred vision, heavy sweating, drooling, vomiting, diarrhea, and loss of bladder control.

Breathing may become difficult because the airways narrow, fluid collects in them, or the muscles needed for breathing become weak. Muscle twitching, weakness, confusion, unusual sleepiness, seizures, or loss of consciousness can occur in severe cases.

Pyrethrin or pyrethroid exposure often causes irritation. Skin exposure may produce temporary tingling or burning, especially on the face. Breathing in a spray can trigger coughing or wheezing. Serious nervous system effects, including seizures, are uncommon but possible after substantial exposure.

Some insecticide products also contain solvents or petroleum-based liquids. These ingredients can cause additional problems, particularly if vomit enters the lungs.

- Eye, nose, throat, or skin irritation
- Headache, dizziness, nausea, or vomiting
- Coughing or chest discomfort
- Tingling, burning, or numbness of exposed skin

Causes and risk factors

Poisoning may follow an accidental spill, incorrect mixing, use in an enclosed space, or failure of protective equipment. It can also occur when a product is transferred into a drink bottle or another unlabelled container.

Agricultural workers, pesticide applicators, pest-control workers, and people handling concentrated products may have a higher exposure risk. Children are vulnerable because of their smaller body size and tendency to put objects or liquids in their mouths.

Risk may also be increased by:

Intentional swallowing is a medical emergency. It also requires compassionate mental health support once immediate physical safety has been addressed.

- Using more product than the label directs
- Mixing products without approved instructions
- Spraying without adequate ventilation
- Eating, drinking, or smoking while handling insecticides
- Wearing contaminated clothing after a spill
- Re-entering a treated area too soon
- Storing products within children's reach

Diagnosis

Diagnosis begins with what happened: the product involved, the amount, when exposure occurred, and whether it was swallowed, inhaled, or spilled on the body. Clinicians also consider symptoms and changes in breathing, pulse, blood pressure, pupil size, sweating, and muscle strength.

Bring the container, label, or a clear photograph to the hospital if it is safe to do so. Do not carry a leaking or contaminated container into a vehicle or treatment area.

Blood tests may measure electrolytes, kidney and liver function, blood gases, and other signs of organ stress. An electrocardiogram, or ECG, records the heart's electrical activity.

For suspected organophosphate or carbamate poisoning, blood cholinesterase tests may support the diagnosis. Cholinesterase is an enzyme involved in nerve signalling. A low level can suggest exposure, but results may not be available quickly and do not always match symptom severity. Treatment should not be delayed when the clinical signs are clear.

Chest imaging may be needed if breathing problems develop or if vomit may have entered the lungs. There is no single test that confirms every type of insecticide poisoning.

Treatment and management

First aid starts by stopping exposure without putting anyone else at risk. Move away from spray or fumes and into fresh air. Avoid entering a contaminated enclosed space unless trained and properly equipped.

Remove contaminated clothing and place it in a sealed bag. Wash exposed skin and hair thoroughly with soap and water. If the eyes are affected, rinse them gently with clean, lukewarm water. Contact lenses should be removed if they come out easily.

Do not make the person vomit unless a poison information service specifically directs it. Do not give food, drink, oils, milk, or other home remedies to "neutralise" the chemical. These may worsen the situation or interfere with treatment.

Hospital care focuses on protecting breathing and supporting circulation. Treatment may include oxygen, airway suction, intravenous fluids, heart monitoring, and mechanical ventilation if a breathing machine is needed. Medicines can be used to control seizures.

Some organophosphate and carbamate poisonings are treated with atropine. This antidote reduces dangerous secretions and helps open the airways. Pralidoxime may be used in certain organophosphate poisonings to restore enzyme function, especially when there is muscle weakness. The choice and timing depend on the product and clinical findings.

There is no single antidote for all insecticides. Pyrethrin and pyrethroid poisoning is generally managed by removing the chemical and treating symptoms. Antibiotics like amoxicillin or doxycycline do not treat chemical poisoning unless a separate infection is present.

People with mild irritation may improve after washing and a period of observation. Significant symptoms often require hospital monitoring because breathing, heart rhythm, or muscle strength can change over time.

Complications

The most serious immediate complication is respiratory failure, meaning the body cannot breathe effectively enough. This may result from airway secretions, narrowing of the airways, reduced consciousness, or weak breathing muscles.

Other possible complications include abnormal heart rhythms, seizures, low blood pressure, kidney injury, and brain injury caused by a lack of oxygen. Vomiting can lead to aspiration, which means stomach contents enter the lungs and cause inflammation or breathing difficulty.

After organophosphate poisoning, some people develop delayed weakness over the following days. This can affect the neck, arms, legs, or breathing muscles. A less common delayed nerve problem may appear weeks later, causing pain, tingling, or weakness in the limbs.

Repeated occupational exposure may cause ongoing symptoms, but headache, fatigue, or concentration problems have many possible causes. These symptoms need a medical assessment rather than being assumed to come from an insecticide.

Prevention

Most accidental insecticide poisonings can be prevented. The safest approach is to use the least hazardous suitable method and follow the product label exactly.

Keep insecticides in their original, labelled containers. Never transfer them into food or drink containers. Store them locked away from children and pets and separate from food, medicines, and animal feed.

When using a product:

Wash contaminated work clothing separately. Workers should receive training and have access to the protective equipment required for the specific chemical.

For household pests, physical barriers, traps, cleaning, and removal of food or standing water may reduce the amount of insecticide needed.

- Follow mixing, ventilation, protective clothing, and re-entry instructions
- Do not combine products unless the label specifically allows it
- Avoid spraying into the wind or near other people
- Wash hands before eating, drinking, smoking, or touching the face
- Remove work clothing before entering living areas
- Clean equipment and dispose of containers as the label directs

Living with it and what to expect

Many people with mild skin, eye, or airway exposure recover after the chemical is removed and irritation settles. This may happen within hours or days, although skin tingling from some pyrethroids can last longer.

Recovery after severe poisoning is less predictable. A hospital stay may be needed until breathing, alertness, muscle strength, and heart function are stable. Fatigue and weakness can continue after discharge, particularly after organophosphate poisoning.

Routine follow-up may include checking breathing, strength, coordination, mood, and the ability to return safely to work. Rehabilitation can help if nerve or muscle damage affects

walking or daily activities. Workplace safety review may be useful before further pesticide handling.

Most people do not remain "poisoned" once the chemical has been cleared. However, lasting problems are possible after severe oxygen loss, prolonged seizures, major lung injury, or delayed nerve damage.

Emotional recovery also matters, especially after intentional poisoning or a frightening workplace incident. Follow-up mental health care can support safety and recovery without judgment.

When to get medical help

Contact a poison information service promptly after a significant or uncertain exposure, even if symptoms are not yet present. Have the product name, label, estimated amount, time of exposure, and current symptoms available.

Seek urgent medical assessment for repeated vomiting, worsening dizziness, marked sweating, very small pupils, muscle twitching, increasing weakness, wheezing, or persistent eye pain.

Call emergency services now if there is:

Do not drive someone who is unconscious, having seizures, or struggling to breathe. Keep yourself away from contaminated clothing, liquid, or fumes while waiting for help.

After recovery, arrange routine follow-up if weakness, numbness, breathing symptoms, memory problems, or mood changes continue or appear later.

- Trouble breathing, blue or grey lips, or choking secretions
- A seizure, collapse, or loss of consciousness
- Severe confusion or inability to stay awake
- Rapidly worsening muscle weakness
- Intentional swallowing or suspected self-harm

Common questions

How quickly do symptoms of insecticide poisoning start?

Some symptoms begin within minutes, especially after inhaling spray or swallowing a concentrated product. Others may take several hours. Delayed muscle weakness can occur

days after certain organophosphate poisonings.

Can I be poisoned just by getting insecticide on my skin?

Yes, because some insecticides can pass through the skin. Risk is higher with concentrated products, large spills, damaged skin, or prolonged contact. Remove contaminated clothing and wash promptly with soap and water.

Should I make myself vomit after swallowing insecticide?

No. Vomiting can draw the product into the lungs, especially when it contains petroleum-based solvents. Contact a poison information service or emergency service for product-specific instructions.

Does insecticide poisoning have an antidote?

Some types do. Atropine and sometimes pralidoxime are used for certain organophosphate or carbamate poisonings. Many other insecticides have no specific antidote and are treated with decontamination, monitoring, and supportive care.

Can insecticide poisoning cause permanent damage?

It can, particularly after severe breathing failure, lack of oxygen, prolonged seizures, or delayed nerve injury. Many people with mild or promptly treated exposure recover without lasting effects.

Is the smell of insecticide enough to prove poisoning?

No. Odour does not reliably show how much chemical entered the body, and some hazardous products have little smell. Symptoms, exposure details, examination, and sometimes tests are more useful.

Sources

- World Health Organization. Clinical Management of Acute Pesticide Intoxication: Prevention of Suicidal Behaviours . 2008.
- US Environmental Protection Agency. Recognition and Management of Pesticide Poisonings , sixth edition. 2013.
- National Institute for Occupational Safety and Health. Organophosphorus Pesticides: Emergency Response Safety and Health Database .
- Eddleston M, Buckley NA, Eyer P, Dawson AH. Management of acute organophosphorus pesticide poisoning. The Lancet . 2008;371:597-607. 61202-1)

- Agency for Toxic Substances and Disease Registry. Medical Management Guidelines for Pyrethrins and Pyrethroids .
- National Pesticide Information Center. First Aid in Case of Pesticide Exposure .