CONDITION

Misophonia

Misophonia causes intense emotional and physical reactions to specific everyday sounds or related sights.

WHAT IT IS
Strong, involuntary reactions to specific everyday sounds

BODY SYSTEM
Nervous system and emotional processing

MAIN SYMPTOMS
Anger, disgust, distress, physical arousal

WHAT CAUSES IT
Unknown, likely altered processing of trigger sounds

HOW IT IS DIAGNOSED
Clinical assessment; no confirmatory test

MAIN TREATMENT
Adapted talking therapy, coping tools, practical accommodations

CAN IT BE CURED?
No, no established cure yet

IS IT AN EMERGENCY?
Sometimes, with immediate risk of self-harm or harming others

LONG-TERM OUTLOOK
Often persistent, but distress and disruption can improve

Overview

Misophonia is a condition in which certain everyday sounds cause an unusually strong emotional and physical reaction. The word means "hatred of sound," but the experience is more complex than simply disliking noise.

Common triggers include chewing, breathing, sniffing, throat-clearing, keyboard tapping, and pen clicking. A trigger is usually linked to a specific pattern or meaning rather than its volume. A quiet chewing sound may cause intense distress, while a louder neutral sound may not.

The reaction can feel automatic. Anger, disgust, panic, or an urgent need to escape may appear before you have time to think. Some people also react to related movements, such as seeing someone chew or tap a foot.

Misophonia matters because it can affect meals, relationships, education, work, and social life. It is not a sign of poor character or a deliberate attempt to control other people. However, it can create serious conflict if the condition and its effects are not understood.

There is no single agreed medical test or universally accepted cure. Research is still developing, but support and treatment can help reduce distress and make triggers easier to manage.

Symptoms

Misophonia reactions are usually tied to particular sounds. The same sound may be less difficult when made by an animal, a machine, or the person with misophonia. Who makes the sound, whether it can be controlled, and the setting may all affect the response.

Emotional symptoms may include:

The body may enter a "fight, flight, or freeze" response. This is the nervous system's automatic reaction to something it reads as threatening. You might notice a racing heart, muscle tension, sweating, faster breathing, or pressure in the chest.

Some people repeatedly scan their surroundings for possible triggers. This is called hypervigilance. Anticipating a sound can become nearly as distressing as hearing it.

Misophonia is different from hyperacusis, in which ordinary sounds seem painfully or unbearably loud. It also differs from hearing damage, although these conditions can occur together. Misophonia is mainly about the specific pattern and personal meaning of a trigger.

Symptoms range from manageable irritation to severe disruption. A strong reaction does not mean that someone will act aggressively, but anger and loss of control should be taken seriously.

- Sudden anger, disgust, anxiety, irritation, or panic
- Feeling trapped, overwhelmed, or unable to concentrate
- An urgent need to stop the sound or leave
- Shame or guilt after a strong reaction

Causes and risk factors

The cause of misophonia is not known. Research suggests that the brain may give certain sounds unusually high emotional importance. Connections between sound processing,

attention, memory, and the nervous system's threat response may all be involved.

Triggers can become linked with previous experiences or with repeated anticipation of distress. This does not mean the reaction is imagined. The physical and emotional response can be immediate and involuntary.

Misophonia often begins during childhood or adolescence, although it may be recognised later. Some people report relatives with similar sound sensitivities, but no single inherited cause has been established.

It may occur alongside anxiety, obsessive-compulsive symptoms, attention-deficit hyperactivity disorder, autism, tinnitus, or other forms of sensory sensitivity. These conditions do not automatically cause misophonia, and you do not need to have another condition to experience it.

Stress, tiredness, illness, and feeling unable to leave a situation may make reactions stronger.

Diagnosis

There is no blood test, brain scan, or hearing test that confirms misophonia. Diagnosis is based on a detailed clinical assessment of your symptoms and how they affect daily life.

A clinician may ask about:

Questionnaires may help describe severity, but they do not confirm the condition by themselves. Diagnostic definitions and measurement tools are still being refined.

A hearing assessment may be useful if sounds seem painful, unusually loud, distorted, or linked with tinnitus. A clinician may also consider conditions with overlapping symptoms, such as hyperacusis, migraine-related sound sensitivity, post-traumatic stress, obsessive-compulsive disorder, anxiety conditions, or broader sensory processing differences.

Misophonia is not currently defined consistently across all major diagnostic systems. This can make access to knowledgeable care difficult. Lack of a standard test does not make the symptoms unreal.

- Which sounds or sights trigger reactions
- The emotions and physical sensations involved
- When the problem began and how often it happens
- Avoidance, family conflict, or effects on work and education
- Hearing symptoms and other physical or mental health concerns

Treatment and management

Treatment aims to reduce distress, improve control over responses, and protect daily activities and relationships. No single approach works for everyone, and research remains limited.

Adapted cognitive behavioural therapy, or CBT, has the strongest clinical trial evidence so far. CBT is a talking therapy that examines links between thoughts, feelings, body reactions, and behaviour. For misophonia, it may include learning about automatic arousal, practising attention skills, changing unhelpful expectations around triggers, and developing more flexible ways to respond.

Any work involving trigger sounds should be planned carefully and collaboratively. Simply forcing yourself to tolerate sounds, or being repeatedly exposed without support, is not an established cure and may increase distress.

Practical strategies may include:

Earplugs, headphones, or noise masking can help in selected situations. Wearing them constantly may increase isolation or make it harder to function when protection is unavailable. Keep headphone volume at a safe level.

Family or relationship support can help others understand that the reaction is involuntary while still respecting everyone's needs. Schools and workplaces may be able to offer reasonable adjustments, such as a quieter workspace, permission to use headphones, or flexible seating.

No medicine has been shown to treat misophonia itself. Medication may sometimes be used for a separate condition, such as anxiety or depression, after an appropriate assessment.

- Using background sound during meals or focused work
- Taking planned breaks before distress becomes overwhelming
- Choosing seating that allows space or an easy exit
- Agreeing on calm signals and boundaries with household members
- Using relaxation or grounding skills when physical arousal starts

Complications

Without effective support, misophonia can contribute to social withdrawal, missed meals, concentration problems, family conflict, and difficulty attending school or work. Avoiding more places and people may provide short-term relief but gradually narrow daily life.

Repeated anger and distress can also lead to exhaustion, guilt, hopelessness, or low mood.

These effects deserve care even when the trigger sensitivity itself has not changed.

Misophonia does not mean that someone is violent. However, thoughts of self-harm, harming another person, or losing control require prompt help.

Prevention

There is no proven way to prevent misophonia because its cause is not understood. It is not known to result from poor parenting, lack of discipline, or ordinary noise exposure.

Early recognition may help limit conflict and excessive avoidance. Supportive communication, practical adjustments, and treatment for related anxiety or low mood may reduce its effect on everyday life, but they cannot guarantee that triggers will disappear.

Living with it and what to expect

Misophonia is often long-lasting. Triggers may remain stable, become more numerous, or change over time. Reactions can also become stronger or weaker depending on stress, tiredness, relationships, and how much control you have over the situation.

Some people improve enough to take part in meals, work, travel, and relationships with fewer restrictions. Improvement may mean recovering more quickly, feeling less physical arousal, or having more choices about how to respond. It does not always mean that trigger sounds become pleasant or neutral.

Complete avoidance is rarely practical and can make life smaller over time. A more sustainable plan usually combines selected accommodations with skills for handling situations that cannot be controlled.

People close to you may need clear explanations. It can help to describe the reaction without blaming the person making the sound. Shared plans, such as background music at meals or agreed break times, can reduce tension.

Follow-up is usually routine when symptoms are stable. Review may be useful if avoidance is increasing, coping methods no longer work, or misophonia is affecting nutrition, sleep, education, work, or relationships.

When to get medical help

Arrange a routine appointment if sound-triggered reactions regularly interfere with daily life,

cause significant conflict, or lead you to avoid important activities. You can also seek help if you have painful hearing, tinnitus, unusual sensitivity to loudness, anxiety, low mood, or symptoms of another condition.

A primary care clinician may help assess your general and mental health. Depending on your symptoms and local services, assessment may involve a mental health professional, audiologist, or ear, nose, and throat specialist.

Seek urgent help if you feel unable to control aggressive impulses, are thinking about self-harm or suicide, or cannot safely eat, sleep, attend school, or care for yourself because of distress.

Call emergency services or go to an emergency department if there is an immediate danger that you may harm yourself or someone else. Move away from triggers and weapons or other means of harm if you can do so safely, and involve a trusted person.

Common questions

Is misophonia real?

Yes. Misophonia describes a consistent pattern of strong, involuntary reactions to specific triggers. Researchers have not yet agreed on every diagnostic detail, but the distress and daily impairment can be substantial.

Is misophonia just being annoyed by chewing?

No. Ordinary annoyance is usually mild and passes quickly. Misophonia can cause immediate anger, disgust, panic, physical arousal, and a powerful need to escape. The main difference is the intensity and effect on daily life.

Is misophonia a type of autism, OCD, or anxiety?

Misophonia is not the same as autism, obsessive-compulsive disorder, or an anxiety condition. It can occur alongside any of them, and some symptoms may overlap. A careful assessment can help separate the different concerns.

Why are chewing and breathing common triggers?

The reason is not fully understood. These sounds are repetitive, close to the body, and often made by another person. Their social meaning and predictability may help explain why the brain gives them unusual emotional importance.

Can misophonia be cured?

There is no established cure. Treatment may still reduce the intensity of reactions, improve recovery after triggers, and make school, work, meals, and relationships easier to manage.

Do trigger sounds damage my hearing?

Misophonia triggers do not usually damage hearing when they are at ordinary volume. Painful sounds, sudden hearing changes, or severe loudness sensitivity may point to another hearing problem and should be assessed.

Sources

- Swedo SE, Baguley DM, Denys D, and colleagues. "Consensus Definition of Misophonia: A Delphi Study." Frontiers in Neuroscience . 2022;16:841816.
- Brout JJ, Edelstein M, Erfanian M, and colleagues. "Investigating Misophonia: A Review of the Empirical Literature, Clinical Implications, and a Research Agenda." Frontiers in Neuroscience . 2018;12:36.
- Jager IJ, de Koning P, Bost T, Denys D, and Vulink N. "Misophonia: Phenomenology, Comorbidity and Demographics in a Large Sample." PLOS ONE . 2020;15(4):e0231390.
- Jager IJ, Vulink NCC, Bergfeld IO, van Loon A, and Denys DAJP. "Cognitive Behavioral Therapy for Misophonia: A Randomized Clinical Trial." Depression and Anxiety . 2021;38(7):708-718.
- Schröder A, Vulink N, van Loon A, and Denys D. "Cognitive Behavioral Therapy Is Effective in Misophonia: An Open Trial." Journal of Affective Disorders . 2017;217:289-294.