CONDITION

Narcolepsy

Narcolepsy is a long-term condition that disrupts control of sleep and wakefulness.

WHAT IT IS
Chronic sleep-wake control condition

BODY SYSTEM
Nervous system and sleep regulation

MAIN SYMPTOMS
Daytime sleepiness, sleep attacks, cataplexy, sleep paralysis

WHAT CAUSES IT
Immune-related orexin loss or an unknown cause

HOW IT IS DIAGNOSED
Overnight sleep study and multiple sleep latency test

MAIN TREATMENT
Sleep routines, planned naps, and symptom-targeted medicines

CAN IT BE CURED?
No - symptoms can be managed long term

IS IT AN EMERGENCY?
Sometimes - after a crash or serious injury

LONG-TERM OUTLOOK
Usually lifelong, often manageable with treatment and support

Overview

Narcolepsy is a long-term condition that affects how the brain controls sleep and wakefulness. It can cause overwhelming daytime sleepiness even after what seems like enough sleep at night. Some people fall asleep unexpectedly during ordinary activities.

Sleep can also intrude into wakefulness. This may cause dream-like experiences, temporary inability to move, or sudden loss of muscle strength. At night, sleep may be broken and restless rather than unusually long.

There are two main types:

Orexin, also called hypocretin, helps keep you awake and stabilizes transitions between sleep and wakefulness.

Narcolepsy can affect driving, education, work, relationships, and mental health. It is often

mistaken for insufficient sleep, depression, epilepsy, or another sleep condition. A careful sleep assessment is important because several conditions and medicines can cause similar symptoms.

- Type 1 narcolepsy involves cataplexy, which is sudden muscle weakness while awake, or very low levels of the brain chemical orexin
- Type 2 narcolepsy causes daytime sleepiness without cataplexy and without known low orexin levels

Symptoms

The central symptom is excessive daytime sleepiness . This is more than ordinary tiredness. You may struggle to stay awake while reading, watching television, attending a meeting, travelling as a passenger, or doing repetitive work. Brief naps may be refreshing, but sleepiness usually returns.

A "sleep attack" means falling asleep with little warning. Not everyone with narcolepsy has sudden attacks; some experience a steady and powerful pressure to sleep instead.

Other possible symptoms include:

Cataplexy can affect the jaw, face, neck, knees, or the whole body. Speech may become unclear, the head may drop, or the knees may buckle. Consciousness is usually preserved, and an episode often lasts seconds to a few minutes.

Sleep paralysis and sleep-related hallucinations can also occur in people without narcolepsy. Their presence alone does not confirm the condition.

Symptoms commonly begin during adolescence or early adulthood, but they can start at other ages. Cataplexy may begin later than daytime sleepiness.

- Cataplexy: sudden muscle weakness, often triggered by laughter, surprise, excitement, anger, or stress
- Sleep paralysis: temporary inability to move or speak when falling asleep or waking
- Sleep-related hallucinations: vivid sights, sounds, or sensations during the transition into or out of sleep
- Broken night-time sleep: frequent waking, vivid dreams, or difficulty staying asleep
- Automatic behaviour: continuing a familiar task without full awareness, then having little memory of it

Causes and risk factors

In type 1 narcolepsy, most people have lost many of the brain cells that make orexin. Evidence suggests that the immune system mistakenly targets these cells in someone who is genetically susceptible. The exact trigger is usually not known.

Certain versions of an immune-system gene are strongly associated with type 1 narcolepsy. However, these gene versions are common in people who never develop the condition. Genetic testing therefore cannot confirm narcolepsy by itself.

Type 2 narcolepsy is less well understood. Orexin levels are often normal, and the reason for the disrupted sleep-wake control may remain unknown.

Narcolepsy occasionally develops after an injury, tumour, inflammation, or another condition damages the part of the brain that regulates sleep. This is sometimes called secondary narcolepsy.

Having a close family member with narcolepsy raises the chance of developing it, but most people with narcolepsy do not have an affected relative. It is not caused by laziness, poor motivation, or simply choosing to stay up late.

Diagnosis

Diagnosis usually begins with a detailed history of your sleep, daily routine, medicines, and symptoms. A clinician may ask about cataplexy and whether anyone has seen you fall asleep unexpectedly. A sleep diary or wrist activity monitor may be used to check your sleep schedule before testing.

Other possible causes of sleepiness need to be considered. These include ongoing sleep loss, shift work, obstructive sleep apnoea, restless legs syndrome, depression, thyroid problems, and sedating medicines.

The main tests are usually performed at a sleep centre:

The overnight test is normally done first. It checks that you had enough sleep and looks for other sleep conditions that could affect the daytime test.

For the multiple sleep latency test, an average time to fall asleep of eight minutes or less, together with early REM sleep during at least two sleep periods, supports narcolepsy in the right clinical setting. An early REM period from the previous night may count toward this total.

Test results can be affected by insufficient sleep, irregular schedules, other sleep conditions, and some medicines. Do not stop a medicine before testing unless the clinician arranging the

test gives you a safe plan.

In selected cases, a lumbar puncture may be used to measure orexin in the fluid around the brain and spinal cord. A very low level supports type 1 narcolepsy. There is no routine blood test that confirms the condition.

- Overnight polysomnography: records breathing, oxygen levels, heart rhythm, movement, and brain activity while you sleep
- Multiple sleep latency test: measures how quickly you fall asleep during several scheduled daytime naps and whether rapid eye movement, or REM, sleep starts unusually early

Treatment and management

There is no cure at present, but treatment can reduce symptoms and improve safety. Management is usually tailored to the symptoms that affect you most.

Regular sleep habits can make symptoms easier to manage. A consistent bedtime and wake time, enough night-time sleep, and one or more short planned naps may improve alertness. Planned naps are particularly useful before driving or other activities that require sustained attention, but they do not replace medical treatment when symptoms remain significant.

Medicines may include:

Choice of medicine depends on your symptoms, other health conditions, pregnancy plans, possible interactions, side effects, and what is approved where you live. Some treatments can affect blood pressure, heart rate, mood, or hormonal contraception. Oxybate medicines can cause dangerous sedation when combined with alcohol or other sedating drugs, so their safety instructions are especially important.

Treatment often needs adjustment over time. Follow-up may include reviewing sleepiness, cataplexy, night-time sleep, side effects, blood pressure, and whether symptoms affect driving or work.

Practical support can also help. Schools or workplaces may be able to provide scheduled nap breaks, flexible start times, recorded teaching, extra examination time, or adjustments to repetitive tasks. Local disability and driving rules differ.

Medicines commonly used for this include amphetamine and dextroamphetamine . Which one suits you, if any, is a decision for you and your doctor.

- Wake-promoting medicines to reduce daytime sleepiness
- Stimulant medicines when other treatments are not suitable or effective

- Oxybate medicines for daytime sleepiness, cataplexy, and disrupted night-time sleep
- Medicines that reduce cataplexy , including pitolisant or, in some cases, certain antidepressants like sertraline or fluoxetine

Complications

Uncontrolled sleepiness can increase the risk of road crashes, workplace injuries, burns, and falls. Cataplexy may also cause injury if it happens while standing, using stairs, cooking, swimming, or operating equipment.

Narcolepsy can interfere with concentration and memory because staying awake requires so much effort. School or work performance may be affected even when ability and motivation are unchanged.

Some people experience low mood, anxiety, social isolation, or stigma. Others gain weight, particularly around the time symptoms begin. These concerns deserve attention as part of narcolepsy care rather than being treated as personal failings.

Sleep paralysis and hallucinations can be frightening, but they are usually brief and do not by themselves mean that you have a psychiatric condition. New experiences that happen while fully awake should still be assessed.

Prevention

Primary narcolepsy cannot currently be prevented. There is no proven diet, supplement, vaccine avoidance strategy, or lifestyle change that prevents the immune-related loss of orexin cells.

You can reduce the chance of symptom-related injury. Avoid driving when sleepy, take planned breaks, and follow local driving restrictions. Use extra care around heights, open water, heat, sharp tools, and machinery if sleep attacks or cataplexy are not well controlled.

Regular sleep does not prevent narcolepsy, but it can reduce additional sleep deprivation that makes daytime symptoms worse.

Living with it and what to expect

Narcolepsy is usually lifelong. Daytime sleepiness often continues, although its severity can change. Cataplexy may become less frequent for some people over the years, while disrupted

night-time sleep may remain.

Finding an effective treatment plan can take time. Symptoms may change with work schedules, pregnancy, other health conditions, ageing, or new medicines. Periodic routine follow-up allows treatment and practical support to be reviewed.

Many people continue with education, employment, relationships, exercise, and travel. This may require planned naps, medication, schedule changes, and honest discussions with people who need to understand the condition. Jobs involving driving, heights, weapons, or heavy machinery may need particular assessment.

Driving safety requires ongoing attention, even when treatment helps. Never rely on loud music, fresh air, caffeine, or an open window to overcome severe sleepiness. Follow the medical and licensing requirements where you live.

If you may become pregnant, discuss treatment in advance. The risks of untreated symptoms and the safety of each medicine need individual review. Do not stop prescribed treatment suddenly without medical advice.

When to get medical help

Arrange a routine medical appointment if you regularly struggle to stay awake during the day despite allowing enough time for sleep. Also seek assessment for episodes of emotion-triggered weakness, repeated sleep paralysis, vivid experiences around sleep, or sleepiness that affects work, school, or relationships.

Seek prompt medical advice if sleep attacks occur while driving, cooking, swimming, working at height, or operating machinery. Stop the unsafe activity until the risk has been assessed.

Get emergency help after a road crash, serious fall, significant head injury, or near-drowning. New one-sided weakness, trouble speaking, chest pain, severe breathing difficulty, or loss of consciousness should also be treated as an emergency rather than assumed to be narcolepsy or cataplexy.

A typical cataplexy episode preserves awareness and resolves quickly. A first episode, an unusual prolonged episode, or weakness without an emotional trigger should be medically assessed because other conditions can look similar.

Common questions

Can you have narcolepsy without suddenly falling asleep?

Yes. Some people experience constant severe sleepiness rather than dramatic sleep attacks. Symptoms and sleep-test findings are considered together.

Is narcolepsy the same as being tired all the time?

No. Tiredness can mean low energy or exhaustion. Narcolepsy causes an abnormal tendency to fall asleep, although a person may describe both feelings.

Does everyone with narcolepsy have cataplexy?

No. Cataplexy is associated with type 1 narcolepsy. People with type 2 narcolepsy do not have cataplexy, although the diagnosis may be reviewed if it develops later.

Can I drive if I have narcolepsy?

Some people can drive when symptoms are well controlled, while others cannot do so safely. Licensing and reporting rules differ by location. Driving should stop if you cannot reliably remain alert.

Can narcolepsy go away?

It usually does not go away. Symptoms may become more or less noticeable, and treatment can provide meaningful control, but ongoing management is generally needed.

Is narcolepsy a mental health condition?

No. It is a neurological sleep condition involving the brain's control of sleep and wakefulness. However, its effects can contribute to depression, anxiety, or social stress.

Sources

- American Academy of Sleep Medicine: Treatment of central disorders of hypersomnolence - clinical practice guideline , Journal of Clinical Sleep Medicine , 2021
- National Institute of Neurological Disorders and Stroke: Narcolepsy , condition information
- NHS: Narcolepsy , symptoms, diagnosis, and treatment information
- European Academy of Neurology, European Sleep Research Society, and European Narcolepsy Network: European guideline and expert statements on narcolepsy management , European Journal of Neurology , 2021
- Bassetti CLA and colleagues: Narcolepsy - clinical spectrum, aetiopathophysiology, diagnosis and treatment , Nature Reviews Neurology , 2019

- American Academy of Sleep Medicine: Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in adults , Journal of Clinical Sleep Medicine , 2021