CONDITION

Paint, lacquer, and varnish remover poisoning

Paint, lacquer, and varnish remover poisoning is harmful exposure to the solvents or caustic chemicals in stripping products.

WHAT IT IS
Toxic injury from paint-removing chemicals

BODY SYSTEM
Lungs, nervous system, heart, skin, and digestive system

MAIN SYMPTOMS
Headache, dizziness, breathing trouble, skin or eye burns

WHAT CAUSES IT
Exposure to toxic solvents or caustic chemicals

HOW IT IS DIAGNOSED
Exposure history, examination, blood tests, and heart monitoring

MAIN TREATMENT
Decontamination, oxygen, monitoring, and supportive hospital care

CAN IT BE CURED?
Sometimes - prompt care can reverse effects

IS IT AN EMERGENCY?
Yes - after swallowing or serious fume exposure

LONG-TERM OUTLOOK
Often full recovery; severe exposure can cause lasting injury

Overview

Paint, lacquer, and varnish removers are products that soften or dissolve old finishes. Poisoning happens when their chemicals are swallowed, inhaled, splashed into the eyes, or left on the skin.

These products do not all contain the same ingredients. Some contain solvents such as methylene chloride, also called dichloromethane, or chemicals such as methanol, acetone, toluene, or N-methyl-2-pyrrolidone. Others use strong acids or alkalis, which can burn tissue. A product described as "safer" or "low odor" can still be harmful.

Fumes can build up quickly, especially in a bathroom, basement, workshop, tank, or other enclosed space. Some solvents affect the brain and cause sleepiness or loss of coordination. Methylene chloride can also be changed by the body into carbon monoxide, reducing the

blood's ability to carry oxygen.

The outcome depends on the ingredient, amount, route of exposure, ventilation, and how quickly help begins. A brief, minor skin exposure is very different from swallowing remover or breathing concentrated fumes. The product container or safety data sheet can help poison specialists identify the risks.

Symptoms

Symptoms may begin during exposure or within minutes to hours. Some lung and carbon monoxide effects can become clearer after the person has left the area.

Breathing fumes may cause:

Swallowing remover may cause burning in the mouth or throat, drooling, vomiting, stomach pain, or difficulty swallowing. Solvents can enter the lungs during swallowing or vomiting. This is called aspiration and can lead to serious lung inflammation.

Skin contact can cause redness, pain, blistering, numbness, or a chemical burn. Solvents may remove the skin's protective oils, leading to cracking and inflammation. Some chemicals can also pass through the skin.

Eye exposure may cause intense pain, watering, redness, blurred vision, or damage to the surface of the eye.

Carbon monoxide produced from methylene chloride exposure may cause headache, nausea, confusion, chest pain, or weakness. A standard fingertip oxygen monitor can look normal even when carbon monoxide is affecting the blood.

- Headache, dizziness, nausea, or unusual tiredness
- Confusion, poor coordination, drowsiness, or fainting
- Cough, throat irritation, chest tightness, or shortness of breath
- A fast or irregular heartbeat
- Seizures or coma after severe exposure

Causes and risk factors

Poisoning is caused by direct exposure to the product's chemicals. Common situations include stripping furniture indoors, working without enough ventilation, spilling remover on the body, transferring it to an unlabelled container, or accidental swallowing by a child.

Risk is higher when:

Children are at particular risk from small amounts because of their size. An intentional exposure requires emergency care, regardless of the amount reported.

A cloth face covering, surgical mask, or ordinary dust mask does not protect against solvent vapors. Even some gloves allow solvents to pass through, so protection must match the product label or safety data sheet.

- The product is used in an enclosed or poorly ventilated space
- Exposure lasts a long time or involves a large surface
- Respiratory or skin protection is unsuitable for the chemical
- The person has heart or lung disease
- Alcohol, sedatives, or other substances add to drowsiness
- The product contains methylene chloride, methanol, or a caustic chemical

Diagnosis

Diagnosis starts with finding out what happened. Clinicians may ask for the product name, ingredients, route of exposure, approximate amount, time involved, symptoms, and whether the work took place in an enclosed space. Bring the container or a photograph of its label if this can be done safely. Do not bring a leaking container into a vehicle or medical building.

There is no single test for every remover. The examination focuses on breathing, alertness, heart function, skin, eyes, mouth, and signs of chemical burns.

Testing may include:

Blood levels for certain solvents may be available, but treatment usually begins from the exposure history and symptoms rather than waiting for a specialized result.

- Blood tests for oxygen balance, acid levels, kidney function, and liver injury
- A carboxyhemoglobin test for carbon monoxide exposure
- An electrocardiogram, or ECG, to check the heart rhythm
- Chest imaging when aspiration or lung injury is suspected
- Eye examination after a chemical splash

Treatment and management

The first step is to stop the exposure without putting anyone else at risk. After breathing fumes, move into fresh air only if it is safe to do so. Do not enter a contaminated enclosed space to rescue someone without trained emergency personnel and proper breathing equipment.

Remove contaminated clothing and jewelry. Brush off any dry chemical before rinsing. Wash exposed skin with plenty of lukewarm running water. If the eyes were splashed, begin continuous rinsing with clean lukewarm water and remove contact lenses if they come out easily. Do not delay rinsing while looking for a specific solution.

Do not make someone vomit after swallowing remover. Do not give food, drink, activated charcoal, or a chemical intended to "neutralize" the product unless a poison specialist specifically directs it. Vomiting can pull solvents into the lungs, while attempts to neutralize caustic chemicals can create heat and further injury.

Hospital treatment is based on the ingredient and effects. It may include oxygen, airway support, heart monitoring, intravenous fluids, medicines for seizures or breathing problems, and care for skin or eye burns.

People exposed to methylene chloride may receive high-concentration oxygen because the body can convert it into carbon monoxide. Hyperbaric oxygen, delivered under pressure in a special chamber, may be considered for selected severe cases. It is not needed after every exposure.

There is no single antidote for all paint removers. Observation may be needed because breathing problems, abnormal heart rhythms, or carbon monoxide effects can continue or appear after the initial exposure.

Complications

Many limited exposures improve after decontamination and fresh air. More serious poisoning can cause:

Severe methylene chloride exposure can cause carbon monoxide poisoning. People with heart disease may be especially vulnerable because their heart already has a higher need for oxygen.

Repeated workplace exposure may contribute to ongoing headaches, slowed thinking, skin inflammation, or other organ effects, depending on the chemical. These concerns require occupational health assessment rather than simply treating each episode separately.

- Chemical burns and scarring of the skin, eyes, mouth, or digestive tract
- Aspiration pneumonitis, meaning lung inflammation after solvent enters the airway

- Fluid buildup or other delayed injury in the lungs
- Abnormal heart rhythms, low blood pressure, or heart strain
- Seizures, prolonged unconsciousness, or brain injury from low oxygen
- Kidney or liver injury
- Lasting vision problems after a severe eye burn

Prevention

Accidental paint-remover poisoning is often preventable. However, no setup makes every stripping chemical risk-free.

Read the entire label before opening the product. Check the safety data sheet when using it at work or when the label does not clearly describe ventilation and protective equipment. Consider a less hazardous removal method when one is suitable.

Use the product outdoors when the label allows. If indoor use is permitted, provide effective ventilation and avoid small enclosed spaces. A fan or open window may not provide enough protection from a highly volatile solvent.

Wear the eye protection, clothing, gloves, and respiratory protection specified for that chemical. Do not assume household gloves or a dust mask are suitable.

Keep removers in their original, closed containers. Never transfer them into a cup, drink bottle, or food container. Store them away from children, pets, heat, flames, and food. Do not mix a remover with another cleaning or stripping product.

Follow local rules for disposal. Do not pour leftover product into an unlabelled container or leave open waste materials where vapors can build up.

Living with it and what to expect

The course depends on the chemical and the severity of exposure. Mild headache or dizziness from a brief inhalation may improve over hours after fresh air and medical assessment. Minor skin irritation may settle over several days, although dryness or inflammation can last longer.

Chemical burns take longer to heal and may need wound, eye, or digestive-system follow-up. Lung irritation after aspiration can worsen during the first several hours and may require hospital observation. Recovery from severe lung or brain injury can take weeks or months.

After carbon monoxide exposure from methylene chloride, most improvement may occur

once the exposure stops and oxygen treatment begins. Some people develop later problems with memory, concentration, mood, balance, or headaches. New neurological symptoms after discharge should be assessed promptly.

People who recover from a limited exposure often return to their usual activities without lasting effects. Severe exposure can leave breathing, vision, heart, or neurological problems. Follow-up should focus on the organs affected and on preventing another exposure.

If the poisoning happened at work, record the product, task, ventilation, protective equipment, and length of exposure. Occupational health services can assess whether the work process needs to change.

When to get medical help

Call emergency services now if someone has:

Do not enter an enclosed contaminated area yourself. Wait for trained rescuers.

Contact a poison centre immediately after any swallowing, significant fume exposure, or chemical splash to the eyes. Also call for persistent symptoms after skin contact. In the United States, Poison Help is available at 1-800-222-1222 . Elsewhere, use your local poison information service or emergency number. Have the product label nearby if possible.

After initial treatment, arrange prompt follow-up for continuing cough, shortness of breath, swallowing difficulty, skin wounds, eye pain, blurred vision, headaches, memory problems, or unusual fatigue. Routine follow-up may also be advised after a serious exposure even if you feel well.

- Trouble breathing, severe chest pain, or blue or gray lips
- Collapsed, had a seizure, or cannot be awakened
- Severe confusion, loss of coordination, or worsening drowsiness
- Swallowed remover intentionally or swallowed a substantial amount
- Been overcome by fumes in an enclosed space

Common questions

What should I do if I breathed in paint remover fumes?

Leave the area and get into fresh air without re-entering an unsafe space. Contact a poison centre for product-specific advice. Call emergency services for breathing difficulty, chest pain,

confusion, fainting, seizures, or severe drowsiness.

Can paint remover poisoning happen through the skin?

Yes. Some ingredients cause chemical burns, while certain solvents can pass through the skin. Remove contaminated clothing and rinse the skin thoroughly. Seek urgent advice for pain, blistering, numbness, a large exposure, or symptoms such as dizziness.

How long do paint remover fumes stay in your system?

There is no single time because products contain different chemicals. Some solvents are cleared over hours, while methylene chloride can keep producing carbon monoxide after exposure has ended. Symptoms and blood tests guide how long monitoring is needed.

Can paint remover fumes cause carbon monoxide poisoning?

Methylene chloride can. The body converts some of it into carbon monoxide. This can cause headache, nausea, confusion, chest pain, or loss of consciousness. A normal reading on a standard fingertip oxygen monitor does not rule it out.

Is paint remover poisoning fatal?

Severe exposure can be fatal, particularly after swallowing, concentrated inhalation, loss of consciousness, or exposure in an enclosed space. Prompt decontamination, poison-centre advice, and emergency treatment improve the chance of recovery.

Should I make someone vomit after they swallow paint remover?

No. Vomiting can cause solvent to enter the lungs and produce serious inflammation. Call a poison centre or emergency service immediately and follow its instructions.

Sources

- U.S. National Library of Medicine. MedlinePlus Medical Encyclopedia: Paint, lacquer, and varnish remover poisoning
- Agency for Toxic Substances and Disease Registry. Medical Management Guidelines for Methylene Chloride
- Agency for Toxic Substances and Disease Registry. Toxicological Profile for Methylene Chloride
- National Institute for Occupational Safety and Health. NIOSH Pocket Guide to Chemical Hazards: Methylene Chloride
- America's Poison Centers. Poison First Aid

- U.S. Environmental Protection Agency. Risk Management for Methylene Chloride Under the Toxic Substances Control Act