CONDITION

Piroxicam overdose

Piroxicam overdose is the harmful effect of taking more piroxicam than the body can safely handle.

WHAT IT IS
Toxic effects from taking too much piroxicam

BODY SYSTEM
Digestive, kidneys, nervous system, heart and circulation

MAIN SYMPTOMS
Nausea, vomiting, stomach pain, drowsiness

WHAT CAUSES IT
Taking excess piroxicam, accidentally or intentionally

HOW IT IS DIAGNOSED
Exposure history, examination, and blood and urine tests

MAIN TREATMENT
Supportive hospital care and treatment of complications

CAN IT BE CURED?
Yes - effects usually resolve with supportive care

IS IT AN EMERGENCY?
Yes - suspected overdose needs immediate poison advice

LONG-TERM OUTLOOK
Usually full recovery; severe organ injury may persist

Overview

Piroxicam is a prescription nonsteroidal anti-inflammatory drug like ibuprofen or naproxen , or NSAID. It reduces pain and inflammation. A piroxicam overdose happens when someone takes more than their body can safely handle.

An overdose may be accidental, such as taking a dose twice or confusing medicines. It may also be intentional. Taking piroxicam together with other NSAIDs can increase harmful effects, even when the products have different names.

Piroxicam can irritate the stomach and intestines, affect kidney function, and disturb the nervous and circulatory systems. Serious poisoning is not inevitable, but it is possible. The risk cannot be judged safely from symptoms alone.

Piroxicam remains in the body for a relatively long time compared with many other NSAIDs. Someone who feels well soon after taking it may still need observation or medical advice.

Any suspected overdose needs immediate guidance from a poison centre or emergency service. Do not wait for symptoms to develop.

Symptoms

The first symptoms often involve the digestive system or nervous system. Possible symptoms include:

Bleeding from the stomach or intestines may cause vomit containing blood or material that looks like coffee grounds. Stools may become black, sticky, or visibly bloody.

Severe poisoning can cause very low blood pressure, seizures, loss of consciousness, kidney failure, or abnormal acid levels in the blood. Breathing may become slow or ineffective. These effects are uncommon but require emergency treatment.

Symptoms may be affected by the amount taken, the person's age and health, and whether alcohol or other medicines were also used. Kidney disease, dehydration, stomach ulcers, and medicines that affect bleeding or kidney function can increase concern.

A lack of symptoms does not confirm that the exposure is safe.

- Nausea, vomiting, or stomach pain
- Heartburn or irritation in the upper abdomen
- Drowsiness, dizziness, headache, or confusion
- Ringing in the ears
- Reduced urine output
- Fast or difficult breathing

Causes and risk factors

Piroxicam overdose is caused by taking too much piroxicam at one time or over a period of time. Common situations include taking a second dose after forgetting an earlier dose, misunderstanding instructions, or using another person's medicine.

Risk may also increase when piroxicam is taken with another NSAID. Examples include ibuprofen, naproxen, diclofenac, aspirin used for pain, and some combination cold or pain products. Different brand names can make accidental duplication difficult to notice.

More serious effects may be more likely in people who have:

Children can be harmed by medicines left within reach. An intentional overdose may occur during severe emotional distress or a mental health crisis and needs both medical care and compassionate psychological support.

There is no single amount that can reliably predict what will happen to every person. Tablet strength, timing, body size, health conditions, and other substances all matter. A poison centre can assess these details without asking you to make that judgment yourself.

- Kidney, heart, or liver problems
- A history of stomach ulcers or digestive bleeding
- Dehydration from vomiting, diarrhoea, illness, or poor fluid intake
- Older age or frailty
- Medicines that affect blood clotting, blood pressure, or kidney function
- Exposure to several medicines at the same time

Diagnosis

Piroxicam overdose is usually identified from the medication history rather than one confirmatory test. Clinicians will ask what was taken, approximately how much, when it was taken, and whether any other medicines, drugs, or alcohol were involved.

Bring the medicine packet, bottle, or a photograph of the label if this can be done safely. Do not delay emergency care to search for packaging.

An examination may include checks of alertness, breathing, heart rate, blood pressure, temperature, hydration, and abdominal tenderness. Urine output may be monitored.

Blood tests can check kidney function, electrolytes such as sodium and potassium, blood cell counts, liver function, and the body's acid balance. Urine testing may provide further information about kidney function and hydration. An electrocardiogram, or ECG, may be used to check the heart rhythm, especially if the exact exposure is uncertain.

A piroxicam blood level is not routinely available and generally does not guide treatment. If the overdose was intentional or the substances taken are unclear, tests for other common poisons may be needed.

Treatment and management

There is no specific antidote that reverses piroxicam. Treatment supports the body while the medicine is processed and addresses any complications.

The first step is immediate advice from a poison centre or emergency service. Depending on the amount, timing, symptoms, and medical history, the person may be monitored at home under poison-centre guidance or assessed in an emergency department.

Do not make the person vomit. Do not give food, drink, antacids, or another medicine unless a poison specialist or clinician recommends it.

Activated charcoal can bind some medicine in the digestive tract. It may be considered by healthcare professionals after certain recent or substantial overdoses. It is not appropriate in every case and can enter the lungs if given to someone who is drowsy or vomiting. It should not be used at home unless specifically directed by a poison specialist.

Hospital treatment may include:

Piroxicam attaches strongly to proteins in the blood. Because of this, dialysis does not remove much of the drug. Dialysis may still be needed to support the body if severe kidney failure or dangerous blood chemistry develops.

People who took piroxicam intentionally should also receive a mental health assessment and a safe follow-up plan.

- Observation and repeated checks of vital signs
- Intravenous fluids when needed for dehydration or low blood pressure
- Medicines for nausea or digestive irritation
- Oxygen or breathing support
- Treatment for seizures
- Treatment of stomach or intestinal bleeding
- Monitoring and support for kidney failure or abnormal blood chemistry

Complications

The digestive tract and kidneys are the main areas of concern.

Piroxicam can damage the protective lining of the stomach or intestines. This may lead to inflammation, an ulcer, bleeding, or, rarely, a hole in the digestive tract called a perforation. Significant bleeding can cause weakness, faintness, or anaemia, which means too few healthy red blood cells.

Reduced blood flow through the kidneys can cause acute kidney injury. This may be temporary,

especially when treated promptly, but severe injury can require prolonged monitoring or kidney support. Dehydration can make this complication more likely.

Other possible complications include:

Complications may be caused partly by another medicine taken at the same time. This is one reason it is important to report all possible exposures, even if the details are uncertain.

- Severe drowsiness, confusion, seizures, or coma
- Slow or ineffective breathing
- Low blood pressure and poor circulation
- Excess acid in the blood
- Liver injury
- Worsening of an existing heart or kidney condition

Prevention

Piroxicam overdose can often be prevented, although mistakes can still happen.

Keep the medicine in its original labelled container and follow the instructions provided with it. A medication list, diary, timed reminder, or suitable pill organiser may reduce repeated doses. If you are unsure whether a dose was already taken, seek advice rather than taking another automatically.

Check the active ingredients in pain, cold, and flu products. Do not combine piroxicam with another NSAID unless a clinician has specifically advised this. Pharmacists can help identify products from different brands that contain similar medicines.

Store piroxicam in a closed container, out of sight and reach of children. Do not share prescription medicine.

If there has been an intentional overdose, prevention also involves addressing the distress behind it. A safety plan, mental health care, and arranging safer medicine storage can reduce the chance of another crisis.

Living with it and what to expect

Piroxicam overdose is usually a single acute event rather than an ongoing condition. Many people with mild poisoning recover with observation and supportive care. However, the course cannot be predicted from the number of tablets alone.

Because piroxicam stays in the body for a long time, symptoms and test abnormalities may continue beyond the first several hours. Medical teams may repeat kidney tests or blood counts before deciding that it is safe to leave hospital.

After discharge, tiredness or digestive discomfort may take time to settle. Follow any instructions about repeat blood tests, fluid intake, and when to restart regular medicines. Do not restart piroxicam or substitute another NSAID unless this has been discussed with the clinician responsible for follow-up.

Kidney injury or significant digestive bleeding can take weeks or longer to recover. A small number of people may have lasting organ damage after severe poisoning.

After an intentional overdose, emotional recovery matters as much as physical recovery. Follow-up may involve primary care, mental health services, crisis support, and trusted family or friends. This support is part of treatment, not a punishment.

When to get medical help

Contact a poison centre or urgent medical service immediately after any suspected piroxicam overdose , even if there are no symptoms. In the United States, Poison Help is available at 1-800-222-1222 . Elsewhere, use your local poison information service or emergency number.

Call emergency services now if the person:

Do not leave someone alone after an intentional overdose. If it is safe, move remaining medicines away and call emergency services.

Routine follow-up is different from emergency care. After the immediate event, arrange the recommended follow-up for kidney tests, digestive symptoms, or mental health support. Seek earlier review if vomiting, abdominal pain, reduced urine, swelling, unusual tiredness, or dark stools appear or worsen.

- Has collapsed, had a seizure, or cannot be awakened
- Is confused, extremely drowsy, or behaving unusually
- Has slow, difficult, or stopped breathing
- Vomits blood or passes black or bloody stool
- Has severe or worsening abdominal pain
- Is faint, pale, clammy, or very weak
- Produces very little or no urine

- Took the medicine intentionally or may take more

Common questions

How much piroxicam counts as an overdose?

There is no single amount that is safe to use as a cutoff. Risk is influenced by tablet strength, body size, age, health conditions, timing, and other substances taken. Contact a poison centre with the product and exposure details.

I accidentally took an extra dose. What should I do?

Call a poison centre or urgent medical advice service promptly. Do not take more piroxicam, another NSAID, or a home remedy while waiting for instructions. Do not make yourself vomit.

How long does a piroxicam overdose last?

Piroxicam leaves the body slowly, so monitoring or symptoms may continue for more than a few hours. Mild digestive effects may settle sooner, while kidney injury or bleeding can require care over days or longer.

Will I need to have my stomach pumped?

Stomach pumping is not routine for NSAID overdose. A medical team may consider activated charcoal in selected circumstances, particularly soon after a substantial exposure. The decision is based on timing, symptoms, and whether the airway can be protected.

Can dialysis remove piroxicam?

Dialysis removes little piroxicam because the drug binds strongly to blood proteins. It may still be used to manage severe kidney failure or dangerous changes in blood chemistry.

Can a piroxicam overdose cause permanent damage?

Most people recover, but severe kidney injury, major digestive bleeding, or complications from low oxygen or poor circulation can have lasting effects. Prompt assessment and monitoring help identify these problems early.

Sources

- DailyMed: Piroxicam capsules - prescribing information, including the Overdosage section
- MedlinePlus Medical Encyclopedia: Piroxicam overdose
- America's Poison Centers: Poison Help and emergency poison advice

- Royal Children's Hospital Melbourne: Clinical practice guideline for nonsteroidal anti-inflammatory drug poisoning
- Hunter LJ, Wood DM, Dargan PI. The patterns of toxicity and management of acute nonsteroidal anti-inflammatory drug overdose. Open Access Emergency Medicine . 2011;3:39-48.
- Chyka PA, Seger D, Krenzelok EP, Vale JA. Position paper: Single-dose activated charcoal. Clinical Toxicology . 2005;43(2):61-87.