CONDITION

Reactive attachment disorder

Reactive attachment disorder is a childhood condition in which severe early neglect disrupts comfort-seeking and emotional connection with caregivers.

WHAT IT IS
Childhood disorder affecting attachment to caregivers

BODY SYSTEM
Emotional development and relationships

MAIN SYMPTOMS
Limited comfort-seeking, little response to comfort, low positive emotion

WHAT CAUSES IT
Severe early neglect or repeated caregiver disruption

HOW IT IS DIAGNOSED
Developmental history and observed caregiver interactions

MAIN TREATMENT
Stable caregiving and relationship-based therapy

CAN IT BE CURED?
Sometimes - stable care and therapy bring improvement

IS IT AN EMERGENCY?
No - unless a child faces immediate danger

LONG-TERM OUTLOOK
Improvement is possible, but emotional difficulties may remain

Overview

Reactive attachment disorder, often shortened to RAD, is a condition affecting young children who have not had enough stable, responsive care. A child with RAD rarely seeks comfort from a familiar caregiver when distressed and may not respond much when comfort is offered.

The condition begins in early childhood. It is linked to severe social neglect, repeated changes of primary caregiver, or care in settings where there were too few caregivers to meet each child's emotional needs.

RAD is more than shyness, difficult behaviour, or taking time to settle after adoption or foster placement. Many children who experience stress or changes in care do not develop it. A secure attachment may also take time to grow without the child having RAD.

Attachment is the emotional bond that helps a child turn to a familiar adult for safety and comfort. When early care is severely limited or unpredictable, this development can be disrupted. RAD matters because it may affect emotional regulation, relationships, learning, and behaviour.

RAD is different from disinhibited social engagement disorder. Children with that condition may approach unfamiliar adults too freely. A child with RAD is instead emotionally withdrawn from caregivers. The two conditions should not be treated as interchangeable.

Symptoms

The central feature is an emotionally withdrawn pattern with familiar caregivers. When upset, the child rarely or only minimally seeks comfort. The child also shows little response when a caregiver tries to provide comfort.

Other possible features include:

Symptoms must be considered in relation to the child's age and developmental level. Babies and toddlers communicate through facial expressions, movement, crying, play, and turning toward or away from adults. A developmental delay can change how these behaviours appear.

RAD does not mean that a child has no feelings or cannot form relationships. Some children show warmth in certain situations while still having major difficulty using a caregiver as a source of comfort.

Aggression, lying, stealing, controlling behaviour, and cruelty are sometimes described online as defining signs of RAD. They are not the core diagnostic features. These behaviours can have many causes, including trauma, anxiety, developmental conditions, or stress in the child's environment.

- Showing little positive emotion during everyday interaction
- Seeming emotionally distant, watchful, or hard to soothe
- Episodes of sadness, fear, or irritability without an obvious threat
- Limited back-and-forth social engagement with caregivers

Causes and risk factors

RAD is associated with extremely insufficient early care. This means a child's basic emotional needs for comfort, affection, stimulation, and reliable adult attention were repeatedly not met.

Circumstances associated with the condition include:

A history of one of these experiences does not prove that a child has RAD. Most children exposed to adversity do not develop the condition, and children can respond differently to similar experiences.

RAD is not known to be inherited. It is not caused by a child being stubborn or by an ordinary parenting mistake. Adoption and foster care do not themselves cause RAD. Rather, some children entering care have already experienced the severe neglect or disruption linked to it.

Autism, intellectual disability, language difficulties, depression, anxiety, and post-traumatic stress can affect social behaviour. They are not causes of RAD, but some may occur alongside it or resemble parts of it.

- Severe and persistent neglect
- Frequent changes in foster carers or other primary caregivers
- Repeated separations that prevent a stable bond from forming
- Institutional care with very high numbers of children per caregiver

Diagnosis

There is no blood test, brain scan, questionnaire, or single observation that confirms RAD. Diagnosis requires a detailed assessment by a professional with experience in child development, attachment, and trauma.

The assessment usually considers the child's early care, developmental history, current behaviour, and relationships with caregivers. Observing the child with familiar caregivers can show whether the child seeks and accepts comfort when distressed.

Information may also come from foster carers, adoptive parents, birth family members, nursery staff, teachers, social workers, and medical records. The aim is to understand the pattern across time and settings rather than judge one difficult interaction.

Under standard diagnostic criteria, symptoms must begin before age 5, although the diagnosis may be considered later when there is clear evidence that the pattern began early. The child must have reached a developmental age at which selective attachment is possible, usually at least 9 months.

The assessment also checks for other explanations. In particular, RAD is not diagnosed when autism better explains the social difficulties. Hearing or language problems, developmental delay, trauma-related conditions, anxiety, depression, and current instability may also need evaluation.

Be cautious about services that diagnose RAD from a checklist alone or use the term for almost any serious behaviour problem. A careful diagnosis connects the child's specific attachment behaviour with a verified history of severely insufficient care.

Treatment and management

The foundation of treatment is a safe, stable, emotionally available caregiving relationship. Whenever possible, unnecessary changes in primary caregiver should be reduced. Predictable routines and calm, consistent responses can help the child learn that adults are available and reliable.

Treatment often works with the caregiver and child together. Relationship-based therapy may help the caregiver notice the child's signals, respond sensitively, and build positive shared experiences. Caregiver coaching or parenting support can also help adults understand trauma-related behaviour without labelling the child as manipulative.

The exact approach should match the child's age, development, history, and current needs. Older children may also benefit from treatment addressing trauma, anxiety, grief, or emotional regulation. Nursery or school support can provide consistent expectations and a trusted adult during the day.

There is no medicine specifically approved to treat the attachment pattern itself. Medication may sometimes be considered for a separate condition, such as attention-deficit/hyperactivity disorder, anxiety, or depression, after an appropriate assessment.

Coercive treatments are not recommended. These include forced holding, restraint intended to provoke attachment, "rebirthing," withholding food or water, or requiring a child to earn basic needs. Such practices can cause physical and emotional harm. Treatment should protect the child's dignity, safety, and right to basic care.

Complications

RAD may make it harder for a child to accept comfort, manage strong emotions, and build trusting relationships. This can place strain on family life, especially when caregivers interpret withdrawal as rejection.

Some children also have developmental delays, learning difficulties, trauma symptoms, sleep problems, or other mental health conditions. These may reflect the same early adversity rather than RAD alone.

Without stable care and suitable support, social and emotional difficulties may continue.

However, long-term research is limited, and one diagnosis cannot predict an individual child's future. Early improvement may be gradual and uneven rather than a sudden change.

Placement breakdown is an important risk when caregivers lack support or receive unrealistic promises about treatment. Practical respite, informed professional help, and coordinated school and social care can help protect both the child and the caregiving relationship.

Prevention

RAD can often be prevented by ensuring that babies and young children receive consistent, responsive care. Their needs for food, safety, comfort, play, affection, and interaction should be met by a small number of familiar adults.

When a child cannot remain safely with their original caregivers, timely decisions and stable placements may reduce further disruption. Support for struggling families may also help prevent severe neglect.

Not every separation or placement change can be avoided. A necessary move to protect a child is not a failure, and temporary separation does not automatically cause RAD. The goal is to restore safe, predictable care as soon as possible.

RAD is not an inherited condition, so genetic testing or measures used to prevent inherited disorders do not apply.

Living with it and what to expect

Progress often centres on the child's growing ability to use a caregiver for comfort. Early signs may be small, such as staying nearby when upset, accepting brief reassurance, sharing enjoyment, or recovering more quickly after distress.

Change may take months or years. A child can improve while still having difficulties during transitions, conflict, illness, anniversaries, or other stressful periods. Developmental stages may also bring new challenges as relationships become more complex.

Caregivers may find the child's limited response painful. It can help to remember that emotional withdrawal developed as an adaptation to early care; it is not a deliberate verdict on the current family. Warmth, boundaries, and predictable routines remain valuable even when the child does not respond as expected.

Some children make substantial gains after entering stable, sensitive care. Others continue to need support with trust, emotions, behaviour, or relationships. Evidence about RAD into

adulthood is limited, and adult relationship problems should not automatically be labelled as childhood RAD.

Regular follow-up may review development, school progress, caregiver-child relationships, and other mental health needs. The plan may change as the child grows. Caregivers may also need their own emotional and practical support.

When to get medical help

Arrange a routine assessment if a young child consistently does not seek or accept comfort from familiar caregivers, especially when there is a history of severe neglect or repeated changes in care. You can also seek help if relationship or emotional difficulties are affecting home life, nursery, school, or placement stability.

Prompt professional review is appropriate if behaviour changes suddenly, the child stops engaging in usual activities, or there are concerns about depression, trauma, developmental delay, or another condition. A sudden change is not typical evidence of RAD by itself and may have a different cause.

Seek urgent help through local child-protection or emergency services if:

Call emergency services for serious injury, loss of consciousness, severe breathing difficulty, poisoning, or any immediate life-threatening situation.

- A child may be experiencing abuse, neglect, exploitation, or unsafe restraint
- The child has been abandoned or does not have a safe caregiver
- There is an immediate risk of serious harm to the child or another person
- The child expresses an intention to die or seriously injure themselves

Common questions

Can a child with reactive attachment disorder love someone?

RAD does not mean a child is unable to love. It affects how the child seeks safety and comfort from caregivers. Feelings may not appear through the closeness or reassurance-seeking that adults expect. With stable, sensitive care, the child's ability to engage and trust may improve.

Is reactive attachment disorder the same as autism?

No. Both can involve differences in social interaction, but they have different defining features and histories. RAD requires severely insufficient early care and a particular lack of

comfort-seeking toward caregivers. Autism involves broader developmental differences and is not caused by neglect. A specialist assessment helps distinguish them.

Does every adopted or foster child have RAD?

No. Adoption and foster care do not define the condition. Most adopted and foster children do not have RAD. Some may need time and support to build trust after loss, trauma, or changes in care without meeting the diagnostic criteria.

Can reactive attachment disorder be cured?

Some children improve greatly when they have stable care and appropriate relationship-based treatment. Others continue to have emotional or social difficulties. There is no quick cure, and progress cannot be guaranteed, but the child's current environment and support can make an important difference.

Can adults have reactive attachment disorder?

RAD is defined as a childhood condition that begins before age 5. Research on how its specific features continue into adulthood is limited. Adults may have relationship, trauma, anxiety, or mood difficulties, but these need their own assessment rather than assuming they are adult RAD.

Is "holding therapy" an effective treatment?

No reliable evidence supports coercive holding, forced restraint, or rebirthing as treatment for RAD. These practices can be dangerous. Recommended care focuses on safety, caregiver sensitivity, stable relationships, and non-coercive psychological support.

Sources

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR). American Psychiatric Association Publishing; 2022.
- World Health Organization. ICD-11 for Mortality and Morbidity Statistics: 6B44 Reactive attachment disorder. ICD-11.
- Zeanah CH, Chesher T, Boris NW. Practice parameter for the assessment and treatment of children and adolescents with reactive attachment disorder and disinhibited social engagement disorder. Journal of the American Academy of Child & Adolescent Psychiatry. 2016;55(11):990-1003.



- National Institute for Health and Care Excellence. Children's attachment: attachment in children and young people who are adopted from care, in care or at high risk of going into care. NICE guideline NG26; 2015.
- Zeanah CH, Gleason MM. Annual research review: Attachment disorders in early childhood-clinical presentation, causes, correlates, and treatment. Journal of Child Psychology and Psychiatry. 2015;56(3):207-222.