CONDITION

Retrograde ejaculation

Retrograde ejaculation is when semen travels into the bladder instead of leaving the penis during orgasm.

WHAT IT IS
Semen flows into the bladder during orgasm

BODY SYSTEM
Reproductive and urinary systems

MAIN SYMPTOMS
Dry orgasm, little semen, cloudy urine after orgasm, infertility

WHAT CAUSES IT
Bladder-neck dysfunction from surgery, nerve damage, or medicines

HOW IT IS DIAGNOSED
Urine test after orgasm showing sperm

MAIN TREATMENT
Cause review, bladder-neck medicines, or fertility treatment

CAN IT BE CURED?
Sometimes - reversible causes can be treated

IS IT AN EMERGENCY?
No - new pain or bleeding needs prompt assessment

LONG-TERM OUTLOOK
Usually harmless, though fertility may be reduced

Overview

Retrograde ejaculation happens when semen travels backward into the bladder during orgasm instead of leaving through the penis. It may cause a "dry orgasm," meaning that little or no semen comes out.

Normally, a ring of muscle at the opening of the bladder closes during ejaculation. This is called the bladder neck. It directs semen forward through the urethra, the tube that carries urine and semen out of the penis. If the bladder neck does not close properly, semen can enter the bladder.

You can still have sexual desire, an erection, and the sensation of orgasm. Retrograde ejaculation does not usually cause physical harm. Semen that enters the bladder mixes with urine and leaves the body the next time you urinate.

The condition matters mainly because it can reduce fertility. It may also be upsetting if ejaculation changes unexpectedly. Sometimes it points to an underlying issue, such as nerve damage, a medicine effect, or a previous operation.

Retrograde ejaculation is not the same as anejaculation. In anejaculation, semen is not released forward or into the bladder.

Symptoms

The main sign is a noticeable reduction in the amount of semen released during orgasm. There may be no visible semen at all.

Common features include:

Retrograde ejaculation usually does not cause pain. It also does not directly prevent erections or reduce sexual desire. However, the condition causing it, such as diabetes-related nerve damage, may affect other parts of sexual function.

A small amount of semen does not always mean retrograde ejaculation. Low semen production, dehydration, frequent ejaculation, blockage, hormone problems, and other ejaculation disorders can cause similar changes.

- Little or no semen coming from the penis during orgasm
- Cloudy urine after sex or masturbation
- Difficulty causing a pregnancy
- An orgasm that feels different, although sensation may be unchanged

Causes and risk factors

Retrograde ejaculation develops when the bladder neck cannot close firmly during ejaculation. This can result from surgery, nerve damage, or certain medicines.

Operations involving the bladder neck or prostate are common causes. For example, procedures that remove or reshape prostate tissue to improve urine flow may affect the muscle at the bladder opening. The exact risk differs between procedures.

Nerves control the muscles involved in ejaculation. Conditions that may damage these nerves include:

Certain medicines can disrupt forward ejaculation. These include some medicines used for an enlarged prostate, high blood pressure, and mental health conditions. Not every change in

ejaculation caused by a medicine is true retrograde ejaculation; some medicines reduce or stop semen release in other ways.

Rarely, a person is born with a structural or nerve problem that affects the bladder neck.

Having one of these risk factors does not mean you will develop retrograde ejaculation. When symptoms begin soon after a new medicine or procedure, that timing can help identify the cause.

- Diabetes, especially when it has caused long-term nerve damage
- Multiple sclerosis
- Spinal cord injury
- Some operations involving the pelvis or lower abdomen

Diagnosis

A healthcare professional will ask about the amount of semen released, whether orgasm still occurs, and when the change began. They may review your operations, medical conditions, and medicines. Do not stop a prescribed medicine without discussing it with the person who prescribed it.

The main test is a urine test performed shortly after orgasm. You may be asked to empty your bladder, provide a semen sample through masturbation, and then provide a urine sample. The laboratory checks the urine for sperm.

Finding a significant amount of sperm in the urine, together with little or no semen released, supports the diagnosis. Small numbers of sperm can sometimes appear in urine without retrograde ejaculation, so the result is considered alongside your symptoms.

If fertility is a concern, testing may also include a semen analysis. This measures semen volume and looks at sperm number, movement, and shape. Blood tests or other examinations may be used when there are signs of hormone, nerve, or urinary problems, but these do not by themselves confirm retrograde ejaculation.

If no sperm are found in either semen or post-orgasm urine, another condition may be responsible.

Treatment and management

Treatment is not always needed. Retrograde ejaculation is generally harmless, so you may

choose no treatment if it is not distressing and you are not trying to cause a pregnancy.

Management starts with looking for the cause. If symptoms began after starting a medicine, the prescriber may consider changing the medicine or adjusting treatment. This is not always possible, and ejaculation may not return immediately. Never change or stop medicine on your own.

Some medicines can help the bladder neck stay closed during ejaculation. They work by increasing the activity of nerves or muscles at the bladder opening. They are more likely to help when the muscle and nerve pathways are still intact, such as in some medicine-related or diabetes-related cases. They are less likely to work when surgery has permanently altered the bladder neck.

These medicines can affect blood pressure, heart rate, and other body functions. A healthcare professional needs to consider your medical history and other medicines before deciding whether one is appropriate.

If having a biological child is a goal, fertility treatment may be helpful even when forward ejaculation cannot be restored. Options can include:

The most suitable approach is based on the number and quality of sperm recovered, how long pregnancy has been attempted, and fertility factors affecting both partners. A fertility clinic or specialist in male reproductive health can explain the options.

Counselling or sex therapy may help if the change has affected confidence, relationships, or sexual wellbeing. This does not mean that the condition is psychological.

- Collecting sperm from urine passed immediately after orgasm
- Preparing the bladder and urine to protect sperm before collection
- Using collected sperm for intrauterine insemination
- Using in vitro fertilisation, sometimes with sperm injection into an egg
- Retrieving sperm directly from the reproductive tract when necessary

Complications

Retrograde ejaculation does not usually injure the bladder. Semen in the urine does not normally cause infection or damage.

The main medical complication is reduced fertility. Pregnancy through penis-in-vagina sex may be difficult when little or no sperm leaves the penis, but biological parenthood may still be possible with treatment.

The change can also cause worry, embarrassment, or dissatisfaction with sex. Some people avoid intimacy because they are concerned about the lack of visible semen. Clear information and open communication with a partner may help.

An underlying condition, rather than retrograde ejaculation itself, may need ongoing care. Diabetes-related nerve damage, for example, can affect the bladder and erections as well as ejaculation.

Prevention

Retrograde ejaculation cannot always be prevented. It may follow necessary surgery or permanent nerve damage, and some people are born with conditions that affect ejaculation.

Managing diabetes may reduce the chance of further nerve damage, but it cannot guarantee that ejaculation will remain unchanged.

Before an operation involving the prostate, bladder neck, pelvis, or nearby nerves, ask how the procedure could affect ejaculation and fertility. If you may want biological children later, sperm banking before treatment may be worth discussing.

If a medicine affects ejaculation, speak with its prescriber. Do not stop it suddenly in an attempt to prevent or reverse symptoms.

Living with it and what to expect

Retrograde ejaculation often remains stable rather than steadily worsening. The likely course is mainly shaped by its cause.

A medicine-related change may improve after treatment is safely adjusted. Recovery from temporary nerve irritation after surgery may take time. Changes caused by permanent surgery or established nerve damage are more likely to continue.

You may still have erections and orgasms. The orgasm may feel the same, less intense, or simply different. Cloudy urine after orgasm can continue because semen is leaving the bladder with urine.

The condition does not usually limit exercise, work, travel, or everyday activities. It also does not directly lower testosterone or remove sexual desire. Erectile dysfunction can occur at the same time, particularly with diabetes or nerve disease, but it is a separate problem.

If future fertility matters, it can help to seek advice before you are ready to try for a pregnancy. This allows time to review the cause, test semen and post-orgasm urine, and discuss sperm

storage or assisted reproduction. No fertility treatment can promise pregnancy, but several ways of recovering and using sperm may be available.

When to get medical help

Arrange a routine appointment if you repeatedly release little or no semen, especially if this is a new change. An assessment is also appropriate if you are worried about the effect on your sex life or are having difficulty causing a pregnancy.

Seek prompt medical advice if the change occurs with:

Retrograde ejaculation itself is not an emergency. Go to urgent or emergency care if you cannot pass urine, have sudden severe testicular pain or swelling, or have serious pelvic or genital trauma. These symptoms suggest another problem that needs rapid assessment.

- Pain during ejaculation or urination
- Blood in semen or urine, particularly if it returns
- Fever, pelvic pain, or unusual discharge
- New weakness, numbness, or loss of bladder control
- Symptoms that began after an injury or medical procedure

Common questions

Can I still have an orgasm with retrograde ejaculation?

Yes. Orgasm and ejaculation are related but separate processes. You may still feel an orgasm even though little or no semen leaves the penis. The sensation may feel unchanged or less intense.

Is semen going into my bladder harmful?

Usually not. Semen mixes with urine and leaves the body when you next urinate. It does not normally damage the bladder. Your first urine after orgasm may look cloudy.

Can I get someone pregnant if I have retrograde ejaculation?

Natural pregnancy may be less likely because fewer sperm reach the vagina. It is not always impossible if some semen still comes out. When no sperm are released forward, specialists may be able to recover sperm from post-orgasm urine or directly from the reproductive tract for fertility treatment.

Can retrograde ejaculation be reversed?

Sometimes. Improvement is more likely when a medicine can be safely changed or when the bladder neck can still respond to treatment. Changes caused by permanent surgery or severe nerve damage may not reverse. Fertility treatment can still be considered if ejaculation cannot be restored.

Is retrograde ejaculation the same as erectile dysfunction?

No. Erectile dysfunction means difficulty getting or keeping an erection. Retrograde ejaculation affects where semen goes during orgasm. A person can have one condition without the other, although both may occur with diabetes, nerve damage, or after pelvic treatment.

Can medicines cause dry orgasms?

Yes. Some medicines for prostate symptoms, blood pressure, or mental health conditions can reduce or stop visible ejaculation. This may be retrograde ejaculation or another form of ejaculation difficulty. Ask the prescriber to review the timing and possible alternatives rather than stopping treatment yourself.

Sources

- European Association of Urology. Sexual and Reproductive Health Guidelines: Disorders of Ejaculation
- American Urological Association and American Society for Reproductive Medicine. Diagnosis and Treatment of Infertility in Men Guideline
- NHS. Ejaculation problems
- Mayo Clinic. Retrograde ejaculation: Symptoms and causes
- Parnham A, Serefoglu EC. Retrograde ejaculation, painful ejaculation and hematospermia. Translational Andrology and Urology . 2016;5(4):592-601. doi:10.21037/tau.2016.06.05