CONDITION

Rubber cement poisoning

Rubber cement poisoning is illness caused by swallowing, breathing, or prolonged contact with the solvents in rubber cement.

WHAT IT IS
Toxic effects from exposure to rubber cement solvents

BODY SYSTEM
Nervous system, lungs, heart, eyes, and skin

MAIN SYMPTOMS
Dizziness, headache, nausea, breathing trouble

WHAT CAUSES IT
Swallowing, inhaling, or contacting chemical solvents

HOW IT IS DIAGNOSED
Exposure history, examination, and targeted heart and lung tests

MAIN TREATMENT
Supportive care, fresh air, decontamination, and monitoring

CAN IT BE CURED?
Sometimes - severe exposure may leave lasting damage

IS IT AN EMERGENCY?
Sometimes - breathing trouble, collapse, confusion, or intentional inhalation

LONG-TERM OUTLOOK
Brief exposures often resolve; repeated inhalation may damage nerves

Overview

Rubber cement poisoning means harmful effects from the chemicals in rubber cement. Exposure can happen by breathing its fumes, swallowing the liquid, or getting it in the eyes or on the skin.

Rubber cement is usually made from rubber mixed with fast-drying chemical solvents. The exact solvents differ by product and may include heptane, hexane, toluene, acetone, or related chemicals. The product label or Safety Data Sheet can help a poison specialist identify the ingredients.

A small accidental exposure may cause only temporary irritation or dizziness. Larger exposures can affect the brain, lungs, heart, and other organs. Swallowing is especially concerning if the liquid enters the lungs during choking or vomiting. This is called aspiration , and it can cause

chemical inflammation of the lungs.

Repeated intentional inhalation, sometimes called glue sniffing or huffing, is particularly dangerous. It can cause loss of consciousness, abnormal heart rhythms, nerve damage, and substance use problems. Serious effects may occur even without a long history of use.

Symptoms

Symptoms depend on the product, amount, route of exposure, and how long the exposure lasted.

Breathing rubber cement fumes may cause:

Swallowing rubber cement may cause nausea, vomiting, stomach discomfort, coughing, or choking. If some enters the lungs, coughing and breathing problems may begin quickly or develop over several hours.

Eye exposure can cause burning, redness, pain, and watering. Skin contact may cause dryness, redness, or irritation because solvents remove protective oils from the skin.

Repeated exposure to some solvents, especially n-hexane, can damage peripheral nerves. These are the nerves outside the brain and spinal cord. Symptoms may include tingling, numbness, weakness, reduced grip strength, or trouble walking. They usually develop gradually rather than after a brief accidental smell.

- Headache, dizziness, or feeling light-headed
- Nausea or vomiting
- Drowsiness, confusion, or poor coordination
- Slurred speech or unusual behavior
- Cough, chest discomfort, or shortness of breath
- Fainting, seizures, or unconsciousness after severe exposure

Causes and risk factors

Rubber cement poisoning is caused by exposure to its solvents rather than the dried rubber itself. Fresh liquid cement and concentrated vapors present the greatest risk.

Exposure is more likely when rubber cement is used in a small, poorly ventilated space. Vapors can build up near the floor because many solvents are heavier than air.

Other factors that raise the risk include:

Some solvents can make the heart more sensitive to stress hormones such as adrenaline. Running, struggling, or being suddenly frightened during intentional inhalation may add to the risk of a dangerous heart rhythm.

- Swallowing the product, especially in young children
- Intentionally breathing concentrated fumes
- Using large amounts without adequate ventilation
- Working with rubber cement regularly without following label precautions
- Mixing it with other solvent-containing products
- Smoking or using flames near the product
- Storing it in a drink bottle or another unlabelled container

Diagnosis

There is no single test that confirms every case of rubber cement poisoning. Diagnosis is usually based on what happened, the product involved, symptoms, and a physical examination.

If possible, take the container, label, or a photo of the ingredients when seeking help. Do not delay emergency care to find it.

A healthcare team may check oxygen levels, breathing, alertness, heart rate, blood pressure, coordination, and muscle strength. An electrocardiogram, or ECG, records the heart's electrical activity and may be used after a significant inhalation or when heart symptoms are present.

A chest X-ray may be needed if rubber cement was swallowed and there is coughing, choking, low oxygen, or breathing difficulty. Lung changes from aspiration may not appear immediately, so observation can be important even when an early test is normal.

Blood tests are chosen according to the exposure and symptoms. Routine drug screening does not reliably identify all rubber cement solvents. Long-term numbness or weakness may require nerve conduction studies, which measure how well nerves carry electrical signals.

Treatment and management

There is no general antidote that reverses rubber cement poisoning. Treatment removes the person from exposure, limits further contact, and supports breathing, circulation, and other body functions while the chemicals clear.

Immediate first-aid steps include:

Do not give food, drink, or home remedies to someone who is very drowsy, vomiting, having a seizure, or unable to swallow safely.

In hospital, treatment may include oxygen, airway support, heart monitoring, intravenous fluids, and medicines for specific complications. People with possible aspiration may be watched for worsening cough, fever, low oxygen, or breathing difficulty. Antibiotics like amoxicillin or doxycycline do not routinely treat the initial chemical irritation itself, although they may be used if a separate bacterial infection is later identified.

Someone with repeated intentional exposure may benefit from confidential substance use and mental health support. Treatment can address cravings, triggers, housing or social pressures, and other substance use. Physical therapy or occupational therapy may help if nerve damage has affected strength, balance, or daily activities.

- Move away from fumes and into fresh air without entering an unsafe space
- Remove contaminated clothing and wash exposed skin with soap and water
- Rinse an exposed eye promptly with plenty of lukewarm water
- Do not make someone vomit after swallowing rubber cement
- Contact a poison center for advice based on the exact product and exposure

Complications

Many brief exposures do not cause lasting harm, but severe or repeated exposure can lead to complications.

Aspiration can cause chemical pneumonitis, meaning inflammation of the lungs caused by chemicals rather than infection. This may produce worsening cough, low oxygen, fever, or breathing difficulty.

Concentrated inhalation can severely slow brain activity. Possible results include unconsciousness, choking on vomit, seizures, injury from falls, or brain damage if oxygen levels remain low.

Certain solvents can contribute to abnormal heart rhythms. These may cause palpitations, chest pain, collapse, or sudden cardiac arrest.

Repeated exposure to n-hexane can cause peripheral neuropathy, or damage to the nerves in the arms and legs. Recovery may take months after exposure stops, and severe damage may not fully reverse.

Frequent solvent exposure can also affect memory, concentration, mood, work, school, and

relationships. The specific long-term risks depend on the ingredients and pattern of exposure.

Prevention

Most rubber cement poisoning can be prevented by controlling access and limiting exposure to the liquid and its fumes.

Use the product only as directed on its label. Work outdoors or in a well-ventilated space, and close the container when it is not being used. Do not deliberately smell the container or use rubber cement to become intoxicated.

Keep rubber cement in its original labelled container, away from children and pets. Do not transfer it into cups, bottles, or food containers. Choose a lower-solvent or solvent-free adhesive when it is suitable for the task.

Rubber cement vapors can catch fire. Keep the product away from cigarettes, sparks, flames, heaters, and other ignition sources. Follow workplace safety instructions when exposure is part of a job.

If intentional inhalation is happening, simply removing one container may not address the reason for use. Early, nonjudgmental help for inhalant use can reduce repeated exposure and connect the person with appropriate support.

Living with it and what to expect

After a mild, brief inhalation, headache or dizziness often improves after reaching fresh air. Eye and skin irritation usually settles after the chemical has been rinsed away, although persistent pain or redness needs assessment.

Symptoms after swallowing or aspiration can change over the next several hours. Someone who initially seems well may later develop coughing or breathing trouble. Follow the observation and follow-up instructions provided by the poison center or healthcare team.

Recovery from a severe exposure may take longer, especially after lung injury, loss of consciousness, or low oxygen. Follow-up may include checking breathing, heart symptoms, memory, concentration, or physical function.

Nerve damage from repeated n-hexane exposure usually develops over weeks or months. Symptoms may continue to worsen for a period after exposure stops because injured nerves recover slowly. Improvement may take months, and some people are left with weakness or altered sensation.

People recovering from inhalant use can return to school, work, relationships, and usual activities, but the course is not the same for everyone. Ongoing support may be useful because cravings, stress, or contact with inhalants can lead to renewed use.

When to get medical help

Call emergency services now if someone exposed to rubber cement:

Do not enter a space with concentrated fumes unless trained and properly equipped. Emergency responders can manage the rescue without creating another exposure.

In the United States, call Poison Help at 1-800-222-1222 after any swallowing exposure, concerning inhalation, or uncertain exposure. It is available 24 hours a day. Elsewhere, contact your local poison information service. Have the product container or ingredient list available if possible.

Arrange routine follow-up if cough or irritation persists, or if numbness, weakness, balance problems, memory changes, or repeated inhalant use develops. Routine follow-up is also appropriate after emergency treatment when the discharge instructions recommend another lung, heart, or nerve check.

- Has trouble breathing, blue or grey lips, or severe choking
- Collapses, cannot be awakened, or has a seizure
- Has severe confusion, agitation, or loss of coordination
- Develops chest pain, a racing or irregular heartbeat, or faints
- Is in an enclosed area filled with fumes

Common questions

Is smelling rubber cement once dangerous?

A brief accidental smell in a ventilated area is less likely to cause serious poisoning than concentrated or prolonged inhalation. Move to fresh air if you develop headache, dizziness, nausea, or irritation. Seek poison-center advice if symptoms continue or the amount of exposure is uncertain.

What should I do if my child swallowed rubber cement?

Remove the container and gently clear any product from the mouth. Do not cause vomiting. Call a poison center promptly with the product name and ingredients. Call emergency services

if the child coughs repeatedly, has trouble breathing, becomes very sleepy, or has a seizure.

Why shouldn't I make someone vomit?

Vomiting increases the chance that the liquid will enter the lungs. Even a small amount in the airways can cause significant chemical inflammation. Poison specialists can advise whether observation or medical assessment is needed.

Can rubber cement fumes damage the brain?

A severe exposure can reduce oxygen or strongly suppress brain activity, which may cause lasting injury. Repeated solvent inhalation may also affect memory, attention, mood, and coordination. A brief exposure that causes no symptoms is less likely to produce these effects.

Can nerve damage from rubber cement improve?

It can improve after exposure stops, but nerves heal slowly. Recovery may take months, and severe damage may be incomplete. New tingling, numbness, weakness, or difficulty walking should be medically assessed.

Is dried rubber cement still poisonous?

Dried cement generally releases much less solvent than fresh liquid cement. Risk is greatest while it is wet, during drying, or when the container is open. Continue to ventilate the area and follow the product label because drying times and ingredients differ.

Sources

- MedlinePlus Medical Encyclopedia. Rubber cement poisoning . U.S. National Library of Medicine.
- Agency for Toxic Substances and Disease Registry. Toxicological Profile for n-Hexane . U.S. Department of Health and Human Services.
- Agency for Toxic Substances and Disease Registry. Toxicological Profile for Toluene . U.S. Department of Health and Human Services, 2017.
- Merck Manual Professional Edition. Hydrocarbon Poisoning .
- Royal Children's Hospital Melbourne. Clinical Practice Guideline: Hydrocarbon Poisoning .
- America's Poison Centers. Poison Help: 1-800-222-1222 .