CONDITION

Simple goiter

A simple goiter is a noncancerous enlarged thyroid that does not initially change thyroid hormone levels.

WHAT IT IS
Noncancerous thyroid enlargement with initially normal hormone levels

BODY SYSTEM
Hormone system

MAIN SYMPTOMS
Neck swelling, throat pressure, swallowing difficulty, cough

WHAT CAUSES IT
Iodine shortage, thyroid growth stimulation, or unknown factors

HOW IT IS DIAGNOSED
Neck examination, thyroid blood tests, and ultrasound

MAIN TREATMENT
Monitoring, cause-specific treatment, radioiodine, or surgery

CAN IT BE CURED?
Sometimes - treatment can shrink or remove it

IS IT AN EMERGENCY?
Sometimes - breathing difficulty needs emergency care

LONG-TERM OUTLOOK
Often stable, but may enlarge or become nodular

Overview

A goiter is an enlarged thyroid gland. The thyroid is a butterfly-shaped gland at the front of your lower neck. It makes hormones that help control energy use, body temperature, heart rate, digestion, and many other body functions.

A simple goiter, also called a simple nontoxic goiter, is an enlarged thyroid that is not caused by cancer or inflammation and does not initially produce too much or too little thyroid hormone. Blood tests are usually normal.

"Simple" describes the thyroid's function and the absence of another identified thyroid condition. It does not mean that every neck swelling is harmless or that an enlarging goiter should be ignored. A healthcare professional may need to rule out thyroid nodules, hormone problems, inflammation, and cancer before using this diagnosis.

A simple goiter may enlarge evenly, which is called a diffuse goiter. Over time, some goiters develop separate lumps called nodules. This may then be described as a nontoxic multinodular goiter.

Many goiters are small and do not affect daily life. Larger ones can cause visible neck swelling or press on the throat, food pipe, or windpipe. Monitoring matters because the size, structure, and hormone activity of a goiter can change over the years.

Symptoms

A small simple goiter may cause no symptoms. It may be noticed during a routine examination, in a photograph, or when clothing begins to feel tighter around the neck.

Possible symptoms include:

The swelling often moves when you swallow because the thyroid is attached to structures around the windpipe.

Pain is not typical of a simple goiter. Sudden pain or rapid swelling may have another cause, such as bleeding into a thyroid nodule or inflammation, and should be assessed promptly.

Symptoms such as unexplained weight change, a racing or unusually slow heartbeat, shaking, heat or cold intolerance, marked tiredness, or changes in bowel habits may suggest altered thyroid hormone levels. These symptoms do not confirm a thyroid problem, but they are reasons to arrange an assessment.

- A smooth swelling at the front of the lower neck
- A feeling of fullness, tightness, or pressure in the throat
- Coughing or frequent throat-clearing
- Difficulty swallowing, especially solid food
- Shortness of breath when lying flat or during activity
- A hoarse or changed voice

Causes and risk factors

The thyroid may grow when it needs more stimulation to make enough hormone. This can happen when the body does not receive enough iodine. Iodine is a nutrient the thyroid uses to make thyroid hormones.

Iodine deficiency remains an important cause in parts of the world where food and salt do not

provide enough iodine. It is less common in places with effective salt iodization or other reliable dietary sources.

In areas with enough iodine, the cause of a simple goiter is often unclear. Several factors may contribute, including differences in how thyroid cells respond to normal growth signals.

Factors linked with goiter formation include:

Foods such as cabbage, broccoli, and soy contain substances that can affect iodine use in laboratory settings. Normal portions are not usually a concern for people who eat a balanced diet and receive enough iodine. Avoiding nutritious foods is not a standard treatment for goiter.

Autoimmune thyroid conditions, in which the immune system attacks or stimulates the thyroid, can also cause enlargement. However, these are separate diagnoses rather than a simple goiter.

- Living in an area where iodine deficiency is common
- Increased thyroid hormone needs during puberty or pregnancy
- A family tendency toward thyroid enlargement
- Certain medicines that affect iodine handling or thyroid function
- Regular exposure to very large amounts of substances that interfere with thyroid hormone production
- Previous thyroid nodules or long-standing thyroid enlargement

Diagnosis

Assessment usually begins with questions about how long the swelling has been present, whether it has changed, and whether you have pressure or hormone-related symptoms. Your neck may be examined while you swallow. Nearby lymph nodes may also be checked.

A thyroid-stimulating hormone, or TSH, blood test shows whether the thyroid is being stimulated normally. TSH is made by the pituitary gland in the brain and tells the thyroid how much hormone to produce. Tests for free T4, and sometimes T3, provide more information if TSH is outside the expected range.

An ultrasound uses sound waves to show the thyroid's size and structure. It can tell whether the enlargement is even or contains nodules. Ultrasound does not use radiation.

Finding a nodule does not mean it is cancer. Many thyroid nodules are noncancerous. Its size and ultrasound appearance help determine whether it needs monitoring or a fine-needle

aspiration biopsy. During this procedure, a thin needle removes cells for laboratory examination.

Other tests may be used in particular situations:

Diagnosis involves confirming the enlargement and excluding other causes. Normal blood tests alone cannot show the thyroid's size or whether nodules are present.

- Thyroid antibody tests if an autoimmune condition is suspected
- A radionuclide thyroid scan if TSH is low, to assess hormone activity
- CT or MRI if the goiter may extend into the chest or press on nearby structures
- A camera examination of the voice box if there is persistent hoarseness or concern about vocal-cord movement

Treatment and management

A small simple goiter that is not causing symptoms may not need active treatment. Monitoring may include neck examinations, thyroid blood tests, and repeat ultrasound when clinically appropriate. The timing is based on the goiter's size, ultrasound findings, symptoms, and whether it is changing.

If iodine deficiency is confirmed or strongly suspected, iodine replacement may help, particularly before the goiter has been present for many years. This should follow local medical and public health advice. Taking large amounts of iodine without guidance can trigger an overactive or underactive thyroid, especially when nodules are present.

Thyroid hormone treatment is appropriate when testing shows an underactive thyroid. Using thyroid hormone solely to suppress TSH and shrink a goiter is not routinely suitable for everyone. It may cause an overly fast heartbeat, bone loss, or other effects if the dose makes thyroid hormone levels too high.

Radioactive iodine can shrink some goiters. The thyroid absorbs the iodine, which gradually reduces thyroid tissue. This treatment may take months to have its full effect and can later lead to an underactive thyroid that requires hormone replacement.

Surgery may be considered when a goiter:

Surgery removes part or all of the thyroid. Possible risks include bleeding, changes to the voice from nerve injury, and damage to nearby parathyroid glands, which control calcium levels. Removing the whole thyroid means lifelong thyroid hormone replacement will be needed.

- Causes breathing or swallowing problems

- Extends into the chest
- Continues to enlarge
- Contains a concerning or cancerous nodule
- Causes substantial discomfort or distress about appearance

Complications

Most small simple goiters cause few problems. Complications are more likely when the thyroid becomes large or develops nodules.

A large goiter can press on the windpipe and make breathing uncomfortable. Pressure on the food pipe may make swallowing difficult. A goiter that grows downward behind the breastbone can cause symptoms even if the visible neck swelling is modest.

Over time, parts of a multinodular goiter may begin producing thyroid hormone without normal control. This can lead to an overactive thyroid, also called hyperthyroidism. Other goiters may eventually be associated with an underactive thyroid.

Bleeding into a nodule can cause sudden swelling and pain. It often settles, but urgent assessment is important if the swelling affects breathing or swallowing.

A simple goiter is not itself cancer. However, a new or changing nodule still needs appropriate assessment because thyroid cancer can also present as a lump. Persistent hoarseness, rapid growth, firm neck lymph nodes, or a hard fixed lump deserve prompt attention.

Prevention

Not every simple goiter can be prevented. There may be no clear preventable cause, especially when inherited tendencies or individual thyroid growth patterns are involved.

Adequate iodine intake helps prevent goiter caused by iodine deficiency. In regions with salt iodization, choosing iodized salt may help provide iodine without increasing the total amount of salt eaten. Seafood, dairy foods, eggs, and some fortified foods can also contain iodine, although amounts differ.

Iodine needs increase during pregnancy and breastfeeding. The right source may depend on diet, local food fortification, prenatal vitamins, and existing thyroid conditions.

Do not use kelp, seaweed concentrates, or high-dose iodine supplements as a general goiter treatment. Their iodine content can be high or unpredictable, and excess iodine can disrupt

thyroid function.

Living with it and what to expect

A small goiter may remain stable for years and require only occasional review. Some become less noticeable, while others slowly enlarge. A diffuse goiter can become nodular over time, particularly when it has been present for many years.

Normal thyroid blood tests may stay normal. They can also change, which is why follow-up may include repeat TSH testing. New symptoms do not always mean the goiter has changed, but they provide a reason for reassessment.

Most people can continue their usual work, exercise, and diet. There is generally no need to avoid turning your neck, speaking, or physical activity unless the swelling causes symptoms during those activities.

Noticeable neck swelling can affect confidence or make clothing uncomfortable. These effects are valid reasons to discuss management, even when the thyroid is functioning normally. Treatment decisions can take account of symptoms, appearance, test results, and your preferences.

Keep follow-up appointments if monitoring has been recommended. Arrange an earlier review if the swelling grows, becomes painful, feels different, or starts affecting swallowing, breathing, or your voice.

When to get medical help

Arrange a routine appointment if you notice a new swelling at the front of your neck, gradual thyroid enlargement, ongoing throat pressure, or a neck lump. These findings often have noncancerous causes, but an examination is needed to identify what they represent.

Seek prompt medical assessment if you have:

Seek emergency help for sudden or severe difficulty breathing, noisy high-pitched breathing, or swelling that makes it difficult to swallow saliva. These signs can mean the airway is being narrowed and should not wait for a routine appointment.

- Rapid enlargement of the swelling
- New or persistent hoarseness
- Increasing difficulty swallowing

- Significant neck pain or tenderness
- A hard lump or enlarged lymph nodes in the neck
- Symptoms suggesting an overactive or underactive thyroid

Common questions

Can a simple goiter go away on its own?

Some goiters become smaller, especially if a temporary increase in thyroid demand or iodine deficiency is corrected. Others remain unchanged or slowly enlarge. A long-standing nodular goiter is less likely to disappear without treatment.

Can I have a goiter with normal thyroid tests?

Yes. "Goiter" describes thyroid size, not hormone function. In a simple goiter, TSH and thyroid hormone levels are usually normal. Ultrasound may still be needed to assess the gland's structure.

Is a simple goiter cancer?

No. A simple goiter is a noncancerous enlargement. However, this description is used after other causes have been considered. Any new, growing, or suspicious thyroid nodule may need separate testing.

Should I take iodine for a goiter?

Not unless iodine deficiency is likely and supplementation has been recommended. Too much iodine can worsen thyroid function in some people. This is particularly important if the thyroid contains nodules or if you already have a thyroid condition.

Will I need surgery?

Many people do not. Surgery is mainly considered for pressure symptoms, extension into the chest, continued growth, concerning nodules, or substantial effects on comfort and appearance. Radioactive iodine may be an alternative in some situations.

Can a goiter grow back after treatment?

It can recur after partial thyroid surgery or after treatment that shrinks rather than removes the gland. Continued monitoring helps identify regrowth, new nodules, or changes in thyroid hormone production.

Sources



- American Thyroid Association. "Goiter." Patient thyroid information
- Merck Manual Professional Edition. "Simple Nontoxic Goiter."
- National Institute for Health and Care Excellence. Thyroid disease: assessment and management (NG145)
- Durante C, Hegedüs L, Czarniecka A, et al. "2023 European Thyroid Association Clinical Practice Guidelines for thyroid nodule management." European Thyroid Journal . 2023;12(5):e230067
- Hegedüs L, Bonnema SJ, Bennedbæk FN. "Management of simple nodular goiter: current status and future perspectives." Endocrine Reviews . 2003;24(1):102-132
- World Health Organization. Guideline: Fortification of food-grade salt with iodine for the prevention and control of iodine deficiency disorders . 2014