CONDITION

Spinocerebellar ataxia

Spinocerebellar ataxia is a group of inherited conditions that gradually affect balance, coordination, speech, and movement.

WHAT IT IS
Inherited conditions causing progressive coordination and balance problems

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Unsteady walking, poor coordination, slurred speech, abnormal eye movements

WHAT CAUSES IT
Inherited genetic changes

HOW IT IS DIAGNOSED
Neurological assessment, genetic testing, and brain imaging

MAIN TREATMENT
Rehabilitation, symptom management, and supportive care

CAN IT BE CURED?
No - treatment supports function and comfort

IS IT AN EMERGENCY?
Sometimes - sudden new symptoms may indicate stroke

LONG-TERM OUTLOOK
Usually progressive, with course shaped by genetic subtype

Overview

Spinocerebellar ataxia, often shortened to SCA, is a group of inherited conditions affecting the brain and nervous system. It mainly damages the cerebellum, the part of the brain that coordinates balance and movement. Some types also affect the spinal cord, nerves, eyes, or other brain areas.

"Ataxia" means poor coordination. It can make walking unsteady, speech less clear, and precise hand movements difficult. SCA usually develops gradually and progresses over time, although the age of onset and rate of change differ widely.

There are many genetic types, identified by names such as SCA1, SCA2, SCA3, and SCA6. These types overlap, but some have additional features. For example, certain types may affect vision, sensation, muscle stiffness, memory, or swallowing.

SCA is not the only possible cause of ataxia. Stroke, multiple sclerosis, vitamin deficiencies, medicines, alcohol, infections, immune conditions, and other inherited conditions can produce similar symptoms. New balance or coordination problems therefore need a medical assessment rather than being assumed to be SCA.

Symptoms

Symptoms may begin in childhood, early adulthood, or later life. Even relatives with the same genetic change may develop symptoms at different ages or experience a different rate of progression.

The most common early problem is difficulty with balance and walking. You may feel unsteady on turns, stumble more often, or need to hold a railing on stairs. Uneven ground and low light can make walking harder.

Other possible symptoms include:

Some subtypes can cause changes in hearing, eyesight, sleep, bladder control, thinking, or mood. These are not present in everyone. SCA7, for example, may damage the retina at the back of the eye and cause progressive sight loss.

Fatigue is common. Movement can require more concentration and effort, while disrupted sleep or low mood may add to tiredness.

Symptoms usually develop on both sides of the body and worsen gradually. Sudden one-sided weakness, facial drooping, severe dizziness, or abrupt loss of coordination is not the usual pattern of SCA and requires emergency assessment.

- Poor coordination of the hands and arms
- Slurred, slow, or uneven speech
- Tremor or difficulty controlling precise movements
- Abnormal, slow, or jerky eye movements
- Blurred or double vision
- Trouble swallowing or coughing during meals
- Muscle stiffness, spasms, weakness, or involuntary movements
- Numbness, tingling, or reduced awareness of foot position

Causes and risk factors

SCA is caused by a harmful change in a gene. Genes contain the instructions used by cells to function. Different genetic changes affect nerve cells in different ways, but many eventually interfere with the survival or communication of cells in the cerebellum.

Most named spinocerebellar ataxias are inherited in an autosomal dominant pattern. This means a person needs one altered copy of the gene to develop the condition. If a parent has a dominant SCA-causing genetic change, each child usually has a 1 in 2 chance of inheriting that change. Inheriting it does not always predict the exact age symptoms will begin or how severe they will be.

Several common types are caused by a repeated section of DNA that becomes too long. In some families, the repeat can expand when passed to the next generation. This may lead to symptoms starting earlier or becoming more severe, a pattern called genetic anticipation.

A person can have SCA without knowing of another affected relative. A parent may have had mild or unrecognised symptoms, died before symptoms appeared, or carried a newly occurring genetic change. Family information may also be incomplete.

Sex, diet, exercise, or parenting do not cause inherited SCA. Lifestyle may affect general health and day-to-day function, but it does not create the underlying genetic change.

Diagnosis

Diagnosis usually begins with a detailed history and neurological examination. The assessment looks at walking, balance, eye movements, speech, reflexes, sensation, strength, and coordination. Your clinician may also ask when symptoms began and whether relatives had similar problems.

A brain MRI can show shrinkage of the cerebellum or related structures. This is called atrophy. MRI can support the diagnosis and help exclude problems such as a tumour, stroke, inflammation, or structural damage, but it usually cannot identify the exact SCA type by itself.

Blood tests may look for treatable causes of ataxia, including vitamin deficiencies, thyroid problems, immune conditions, or metabolic disorders. Medication and alcohol use may also be reviewed.

Genetic testing is used to confirm an inherited SCA and identify its subtype. Testing may target a suspected gene or examine a panel of genes. Some SCA types involve DNA repeat expansions that require a specific test and may not be detected by standard gene sequencing.

A genetic test can also find a change whose meaning is uncertain. This is not the same as confirming SCA. Results need to be interpreted alongside your symptoms, examination, and family history.

Genetic counselling before and after testing can help you understand what the result may mean for you and your relatives. Testing an adult who has no symptoms raises separate emotional, insurance, employment, and family-planning issues, which differ by country.

Treatment and management

There is currently no cure for most forms of SCA, and treatment cannot replace nerve cells already lost. Care focuses on maintaining independence, reducing complications, and managing the symptoms that matter most to you.

Physiotherapy can work on balance, strength, flexibility, posture, and safer ways to move. A therapist may recommend a walking stick, walker, ankle support, or wheelchair when appropriate. Using an aid is intended to reduce falls and conserve energy, not to indicate failure.

Occupational therapy can help with dressing, bathing, writing, cooking, work, and home safety. Adapted utensils, grab rails, seating, and changes to a workspace may make daily tasks easier.

A speech and language therapist can support clearer communication and assess swallowing. Communication devices may help if speech becomes difficult to understand. If swallowing is unsafe or exhausting, changes in food texture, eating technique, or nutritional support may be discussed.

Medicines may be used for particular problems such as muscle stiffness, tremor, painful spasms, bladder symptoms, sleep disturbance, or depression. The best option depends on the SCA subtype, other health conditions, and possible side effects. There is no single medicine that treats every form of SCA.

Regular reviews may include nutrition, weight, swallowing, mobility, falls, mood, vision, and hearing. Certain subtypes need monitoring for their specific effects, such as retinal disease. Research studies and clinical trials are testing treatments aimed at particular genetic changes, but an experimental treatment is not the same as an established benefit.

Complications

Poor balance can lead to falls, fractures, and reduced confidence. Less movement may then contribute to muscle weakness, joint stiffness, constipation, blood clots, or loss of physical conditioning.

Swallowing problems can cause dehydration, weight loss, or food entering the airway. Food or liquid that reaches the lungs can lead to aspiration pneumonia, a lung infection. Repeated

coughing during meals, a wet-sounding voice after drinking, or unexplained weight loss should be reported promptly.

Speech changes may make social situations and work more difficult even when thinking remains clear. Depression, anxiety, and isolation can occur and deserve direct support.

Advanced SCA can increase dependence on other people for personal care and mobility. However, the type and timing of complications differ substantially. Not everyone develops every problem, and some subtypes progress slowly over many years.

Prevention

Inherited SCA cannot currently be prevented through diet, exercise, or lifestyle changes. Nothing a parent did during pregnancy causes the genetic change.

If a familial genetic change is known, genetic counselling can explain the chance of passing it on. Reproductive options may include testing during pregnancy or in vitro fertilisation with testing of embryos. These are personal choices and may not be available, acceptable, or appropriate for everyone.

Falls and some secondary complications can sometimes be reduced. Helpful measures may include removing trip hazards, improving lighting, using recommended mobility equipment, staying physically active within safe limits, and having swallowing problems assessed early. These steps do not stop the underlying condition from progressing.

Living with it and what to expect

Most forms of SCA change gradually over months and years rather than from day to day. Symptoms may feel worse when you are tired, stressed, unwell, or affected by alcohol or sedating medicine. A difficult day does not always mean that the underlying condition has suddenly progressed.

Over time, walking often becomes less secure and daily tasks may take longer. Some people eventually use a walker or wheelchair, while others remain independently mobile for many years. Speech or swallowing may become more affected, but the timing cannot be predicted from the diagnosis alone.

The genetic subtype, age at onset, specific genetic change, and areas of the nervous system involved all influence the course. Some types tend to progress slowly and may have little effect on life expectancy. Others can cause broader disability or complications that shorten life. A specialist familiar with your subtype can give more relevant information without guaranteeing

an exact course.

Regular follow-up is used to notice changes early and adjust support. Rehabilitation may not stop SCA, but it can help you preserve skills, reduce avoidable complications, and adapt as your needs change.

Many people continue relationships, hobbies, education, and employment, sometimes with modified duties or equipment. Planning ahead for transport, home access, communication, finances, and future care can provide more control. Psychological support and contact with an ataxia organisation may also help with the uncertainty of a progressive condition.

When to get medical help

Arrange a routine medical appointment if you develop gradually worsening balance, speech, coordination, vision, or swallowing problems. Assessment is also appropriate if you have a family history of SCA and want information about symptoms or genetic testing.

Contact your healthcare team promptly if you have:

Seek emergency help for symptoms that begin suddenly, especially facial drooping, one-sided weakness or numbness, new severe trouble speaking, sudden loss of coordination, or a severe unusual headache. These may be signs of a stroke or another urgent condition rather than the usual progression of SCA.

Emergency help is also needed for severe choking, inability to breathe, loss of consciousness, or a serious injury after a fall.

- Increasing falls or a new difficulty walking safely
- Coughing, choking, or a wet voice during meals
- Unplanned weight loss or difficulty drinking enough
- New vision loss, severe muscle spasms, or major sleep problems
- Persistent low mood, anxiety, or difficulty coping

Common questions

Is spinocerebellar ataxia the same as multiple sclerosis?

No. Both can affect balance and coordination, but they have different causes. SCA is usually caused by an inherited genetic change that damages certain nerve cells. Multiple sclerosis involves immune-system damage to the protective covering of nerves.

Will my children inherit spinocerebellar ataxia?

For an autosomal dominant SCA, each child of a parent carrying the genetic change usually has a 1 in 2 chance of inheriting it. The chance may be different for other inherited ataxias. Genetic testing and counselling can clarify the pattern in your family.

How quickly does spinocerebellar ataxia progress?

There is no single timetable. Progression is influenced by the SCA subtype, age at onset, genetic change, and individual health. Some people notice slow changes over decades, while others lose function more quickly.

Can exercise stop SCA from getting worse?

Exercise does not remove the genetic cause or reliably stop progression. It may help maintain strength, flexibility, fitness, and confidence with movement. A physiotherapist can help you choose activities that account for balance problems and fall risk.

Does alcohol make spinocerebellar ataxia worse?

Alcohol can temporarily worsen balance, speech, reaction time, and coordination. It may also increase the risk of falls or interact with medicines. Your healthcare team can help you consider what is safest for you.

Can I still drive with spinocerebellar ataxia?

Some people can drive safely in the earlier stages, while others develop problems with coordination, vision, reaction time, or judgment. Driving rules and reporting duties differ by location. A formal driving assessment can evaluate your current abilities.

Sources

- National Institute of Neurological Disorders and Stroke. Spinocerebellar Ataxias . Neurological condition information, National Institutes of Health
- Klockgether T, Mariotti C, Paulson HL. Spinocerebellar ataxia . Nature Reviews Disease Primers . 2019;5:24. doi:10.1038/s41572-019-0074-3
- Bird TD. Hereditary Ataxia Overview . GeneReviews® . University of Washington, Seattle
- National Center for Advancing Translational Sciences. Spinocerebellar ataxia . Genetic and Rare Diseases Information Center
- Zesiewicz TA, Wilmot G, Kuo SH, et al. Comprehensive systematic review summary: Treatment of cerebellar motor dysfunction and ataxia . Neurology . 2018;90(10):464-471

- Ataxia UK. Management of the ataxias: towards best clinical practice . Clinical guidance for inherited and acquired ataxias