CONDITION

Subcortical dementia

Subcortical dementia is cognitive decline caused by damage to brain networks involved in thinking speed, planning, movement, and motivation.

WHAT IT IS
Cognitive decline from disrupted deep-brain networks

BODY SYSTEM
Nervous system

MAIN SYMPTOMS
Slowed thinking, planning difficulty, apathy, walking changes

WHAT CAUSES IT
Brain vessel disease or other neurological conditions

HOW IT IS DIAGNOSED
Cognitive testing, brain imaging, and cause-specific assessment

MAIN TREATMENT
Treating the cause, rehabilitation, and supportive care

CAN IT BE CURED?
Sometimes - treatable causes may partly reverse symptoms

IS IT AN EMERGENCY?
Sometimes - sudden symptoms may signal stroke

LONG-TERM OUTLOOK
Often progressive; course follows the underlying condition

Overview

Subcortical dementia describes a pattern of cognitive decline linked to damage in networks below the brain's outer layer. These networks connect deep brain structures with the frontal lobes, which help with planning, attention, motivation, movement, and self-control.

"Subcortical" means beneath the brain's outer layer, or cortex. "Dementia" means that changes in thinking have become severe enough to interfere with everyday independence. Health professionals may instead use the term major neurocognitive disorder .

Subcortical dementia is a condition or pattern, not one specific disease. It can occur with small blood vessel disease in the brain, Parkinson's disease, Huntington's disease, progressive supranuclear palsy, multiple sclerosis, HIV-related brain disease, and several other conditions.

The distinction between cortical and subcortical dementia is not exact. Brain networks are closely connected, and many disorders affect both areas over time. A person may therefore have a mixture of symptoms rather than a purely subcortical pattern.

Finding the cause matters because it affects treatment and the likely course. Some causes can be treated or controlled. Others cannot currently be cured, but symptoms, safety, and quality of life can still be supported.

Symptoms

Symptoms usually develop gradually, although they may appear in steps when repeated small strokes are involved.

Slowed thinking is common. You may need more time to answer a question, make a decision, or move from one task to another. This is different from not understanding the question.

Planning and organising can become difficult. Tasks such as preparing a meal, managing appointments, or following several instructions may feel overwhelming. This is called executive dysfunction , meaning difficulty managing and directing your thoughts and actions.

Common features include:

Memory problems can occur, but they may look different from the early memory loss often associated with Alzheimer's disease. A person may have trouble bringing stored information to mind but remember it after a hint or prompt. This distinction is not reliable enough to identify the cause by itself.

Speech may become quieter or slower. Some people have trouble finding words or keeping up with fast conversations. Movement symptoms can include short steps, stiffness, shaking, falls, or difficulty starting to walk.

Symptoms depend on which networks are affected and why. Not everyone develops movement problems, and apathy should not automatically be mistaken for depression or unwillingness to participate.

- Slower thinking and processing of information
- Difficulty planning, organising, or switching between tasks
- Reduced attention or mental flexibility
- Loss of motivation, known as apathy
- Walking, balance, stiffness, or coordination changes
- Mood changes, including depression or irritability

Causes and risk factors

A common cause is cerebral small vessel disease , which damages the brain's smallest blood vessels and the white matter carrying signals between brain regions. Repeated small strokes can also interrupt these networks. This may be described as subcortical vascular cognitive impairment or subcortical ischemic vascular dementia.

Factors linked with vascular brain injury include high blood pressure, diabetes, smoking, high cholesterol, irregular heart rhythms, and previous stroke. These factors do not mean that dementia will occur, but they can increase the chance of blood vessel damage.

Other possible causes include:

More than one cause may be present. For example, someone may have both Alzheimer-type changes and blood vessel disease. This is called mixed dementia or mixed neurocognitive disorder.

Age can increase the likelihood of several underlying diseases, but subcortical cognitive problems are not a normal part of ageing. Family history matters for some inherited conditions, especially Huntington's disease, but many cases are not directly inherited.

- Parkinson's disease and related movement disorders
- Huntington's disease, an inherited neurological disease
- Progressive supranuclear palsy
- Multiple sclerosis
- Some infections affecting the brain, including HIV
- Normal-pressure hydrocephalus, involving excess fluid in brain spaces
- Brain injury, inflammation, or exposure to certain toxic substances

Diagnosis

There is no single test that confirms "subcortical dementia" or identifies every possible cause. Diagnosis combines your symptoms, changes in daily function, examination findings, cognitive testing, and brain imaging.

The assessment usually includes questions about when changes began and whether they developed gradually, suddenly, or in steps. With your agreement, information from someone who knows you well can help show how your abilities have changed.

Cognitive tests examine memory, attention, language, processing speed, problem-solving, and

planning. Brief screening tests can identify areas needing more assessment. Detailed neuropsychological testing may give a clearer picture of strengths and difficulties.

A neurological examination checks movement, reflexes, balance, eye movements, sensation, and speech. Magnetic resonance imaging, or MRI, can show small strokes, white-matter damage, fluid buildup, tumours, and patterns associated with particular diseases. A CT scan may be used when MRI is unsuitable or not available.

Blood tests can look for conditions that may cause or worsen cognitive symptoms, such as thyroid problems, vitamin deficiencies, infection, or organ dysfunction. Other tests, including spinal fluid testing, genetic testing, or specialised scans, are used only when the history suggests a particular cause.

Sudden confusion is not usually diagnosed as dementia. It may be delirium , a rapid change in attention and awareness often caused by illness, medication, dehydration, or another medical problem. Delirium needs prompt assessment.

Treatment and management

Treatment is based on the underlying cause and the problems affecting daily life. There is no single medicine for all forms of subcortical dementia.

For vascular causes, management may focus on reducing further damage from strokes and small vessel disease. This can include treatment of high blood pressure, diabetes, cholesterol problems, smoking, or heart conditions. Medication choices are individual and require medical review.

If another neurological condition is responsible, treatment may target that condition. For example, movement symptoms related to Parkinson's disease may be managed differently from cognitive changes caused by normal-pressure hydrocephalus or an infection.

Medicines used for Alzheimer's disease are sometimes considered for certain diagnosed dementias or mixed disease. Their benefit is not established for every subcortical condition, and they can have side effects. No one should start or stop cognitive, movement, mood, or blood-thinning medication without appropriate medical advice.

Non-drug support is often important:

Regular reviews can check cognition, mood, movement, falls, swallowing, sleep, medication effects, and the level of support needed. Family members and other caregivers may also need practical information and respite support.

- Occupational therapy can simplify tasks and improve home safety

- Physiotherapy can support walking, strength, balance, and transfers
- Speech and language therapy can help communication or swallowing
- Psychological treatment may help with depression or adjustment
- Routines, written prompts, and quieter surroundings can reduce overload

Complications

As cognitive and movement problems increase, managing medicines, money, cooking, driving, or personal care may become harder. Reduced judgment or slower reactions can create safety risks even when memory seems fairly good.

Walking and balance changes may lead to falls and injuries. Swallowing problems can cause weight loss, dehydration, or food and liquid entering the lungs. This can lead to a chest infection called aspiration pneumonia.

Apathy, depression, anxiety, irritability, or loss of emotional control may affect relationships and participation in daily life. These symptoms can come from changes in brain networks, the emotional impact of illness, or both.

People with dementia are also more likely to develop delirium during infections, hospital stays, surgery, or medication changes. A sudden decline should not be assumed to be expected progression.

Later complications depend heavily on the cause. Some people mainly need help with complex tasks, while others develop substantial movement, communication, or personal-care needs.

Prevention

Subcortical dementia cannot always be prevented. There is no established way to prevent inherited conditions such as Huntington's disease, and there is no proven prevention for every degenerative neurological disease.

Some vascular brain damage may be preventable. Managing blood pressure, diabetes, cholesterol, smoking, and heart disease can reduce the risk of stroke and may help protect brain health. Regular physical activity, a balanced diet, adequate sleep, and limiting alcohol can support overall vascular health.

Preventing head injuries and treating infections that can affect the nervous system may reduce specific risks. These measures cannot guarantee that cognitive decline will not develop.

If you already have an underlying condition, attending routine follow-up and reporting new cognitive, mood, or movement changes may allow treatable problems to be identified earlier.

Living with it and what to expect

The course of subcortical dementia is determined mainly by its cause. Degenerative conditions often worsen gradually over years. Vascular cognitive impairment may progress slowly, remain stable for a time, or worsen in steps after additional strokes. Treatable causes may improve, although recovery is not always complete.

Abilities do not always change at the same rate. Thinking may become slow while language and recognition remain useful. Movement problems may be more limiting than memory loss. Fatigue, pain, poor sleep, depression, infection, or unfamiliar surroundings can temporarily make symptoms more noticeable.

Many people continue to take part in family life, hobbies, exercise, and social activities, especially with a steady routine and enough time to complete tasks. Breaking activities into smaller steps and reducing distractions may help.

Planning ahead can protect your choices. This may include discussing driving, work, finances, future care preferences, legal decision-making, and who should help if your judgment or communication changes. These conversations are often easier before urgent decisions are needed.

Routine follow-up is used to monitor progression and adjust support. New symptoms should still be assessed rather than assumed to be part of dementia, particularly if they appear quickly.

When to get medical help

Arrange a routine medical assessment if you or someone close to you notices persistent slowing of thought, planning problems, loss of motivation, walking changes, or increasing difficulty with everyday tasks. Earlier assessment can identify medication effects, depression, sleep problems, vitamin deficiencies, or other treatable contributors.

Seek prompt medical advice for a noticeable decline over days or weeks, repeated falls, new swallowing problems, hallucinations, major behaviour changes, or difficulty managing medicines, meals, or personal safety.

Call emergency services for:

These symptoms can indicate stroke or another urgent condition. Do not wait to see whether

they pass.

- Sudden weakness or numbness, especially on one side
- Sudden trouble speaking, understanding, seeing, or walking
- A sudden severe headache or loss of consciousness
- New seizure activity
- Rapid confusion with severe illness, breathing difficulty, or inability to wake

Common questions

Is subcortical dementia the same as Alzheimer's disease?

No. Alzheimer's disease usually begins with changes affecting memory-related brain networks, while a subcortical pattern often involves thinking speed, planning, motivation, and movement. However, the patterns can overlap, and both conditions may occur together.

Does subcortical dementia always get worse?

Not always in the same way. Some causes are progressive, while vascular symptoms may remain stable for periods or worsen after further strokes. Treating an infection, fluid-pressure problem, medication effect, or another reversible contributor may produce improvement.

Can an MRI show subcortical dementia?

MRI can show damage that supports a diagnosis, such as small strokes, white-matter disease, or excess fluid. It cannot diagnose the condition by itself. Scan findings must be compared with symptoms, cognitive testing, and daily function.

Can you have it without severe memory loss?

Yes. Early problems may involve slowed thinking, planning, attention, motivation, or walking more than memory. Dementia is defined by the effect on everyday independence, not by memory loss alone.

Is apathy the same as depression?

No. Apathy is reduced motivation or initiative, while depression usually involves persistent low mood, hopelessness, or loss of pleasure. They can occur together, and both deserve assessment because their management may differ.

Can someone with subcortical dementia still live independently?

Some people remain independent during earlier stages, especially with routines, reminders, and support for complex tasks. Independence may decrease if thinking, judgment, walking, or the underlying disease worsens. Safety and support needs should be reviewed over time.

Sources

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-5-TR . Fifth edition, text revision. American Psychiatric Association Publishing; 2022.
- Bonelli RM, Cummings JL. Frontal-subcortical circuitry and behavior. Dialogues in Clinical Neuroscience . 2007;9(2):141-151.
- Gorelick PB, Scuteri A, Black SE, et al. Vascular contributions to cognitive impairment and dementia: a scientific statement from the American Heart Association and American Stroke Association. Stroke . 2011;42(9):2672-2713.
- O'Brien JT, Thomas A. Vascular dementia. The Lancet . 2015;386(10004):1698-1706.
- Román GC, Erkinjuntti T, Wallin A, Pantoni L, Chui HC. Subcortical ischaemic vascular dementia. The Lancet Neurology . 2002;1(7):426-436.
- National Institute for Health and Care Excellence. Dementia: assessment, management and support for people living with dementia and their carers . NICE guideline NG97; 2018, updated 2025.