CONDITION

Synergistic necrotizing fasciitis

Synergistic necrotizing fasciitis is a rapidly spreading deep-tissue infection caused by several types of bacteria acting together.

WHAT IT IS
Rapid mixed-bacterial infection destroying deep soft tissue

BODY SYSTEM
Skin, fat, connective tissue, and muscles

MAIN SYMPTOMS
Severe pain, swelling, skin colour change, fever

WHAT CAUSES IT
Several bacteria acting together in deep tissue

HOW IT IS DIAGNOSED
Urgent surgical inspection with deep tissue cultures

MAIN TREATMENT
Emergency surgery, intravenous antibiotics, intensive support

CAN IT BE CURED?
Yes - prompt treatment can eliminate the infection

IS IT AN EMERGENCY?
Yes - every suspected case needs immediate hospital care

LONG-TERM OUTLOOK
Recovery can be prolonged; severe cases may be fatal

Overview

Synergistic necrotizing fasciitis is a rare but serious bacterial infection of the tissues beneath the skin. "Necrotizing" means causing tissue death. "Fascia" is the layer of connective tissue surrounding muscles and other structures.

The word "synergistic" means that several kinds of bacteria are acting together. One group may use oxygen in the tissue, creating conditions in which other bacteria can grow. This mixed infection is often described medically as a type I necrotizing soft-tissue infection.

The infection can spread along the fascia faster than changes appear on the skin. It may damage fat, blood vessels, nerves, and muscle. Antibiotics like amoxicillin or doxycycline alone usually cannot reach all infected areas because dead tissue has a poor blood supply.

This is a medical and surgical emergency. Early surgery to remove infected and dead tissue, combined with antibiotics and hospital support, offers the best chance of controlling it. Treatment should not wait for the skin to become severely discoloured or blistered.

Symptoms

Symptoms often develop over hours or a few days. Early signs can resemble a less serious skin infection, which can make the condition difficult to recognise.

Severe pain is an important warning sign. The pain may be much greater than expected from the visible skin changes. It may also extend beyond the area that looks red or swollen.

Possible symptoms include:

Not everyone has every symptom. Gas under the skin is particularly associated with some mixed bacterial infections, but its absence does not rule out necrotizing fasciitis.

As nerves are damaged, an area that was extremely painful may become numb. This is not necessarily an improvement. It can be a sign of deeper tissue injury.

- Rapidly increasing pain, tenderness, warmth, or swelling
- Red, purple, grey, or darkening skin
- Fever, chills, weakness, or feeling seriously unwell
- Blisters or fluid leaking from the skin
- A crackling feeling caused by gas beneath the skin
- Dizziness, confusion, fast breathing, or reduced urination

Causes and risk factors

Synergistic necrotizing fasciitis occurs when several bacteria enter deeper tissue and grow together. The mixture may include bacteria that need oxygen and bacteria that grow with little or no oxygen.

Bacteria may enter through a surgical wound, cut, puncture, pressure sore, burn, injection site, or long-standing skin ulcer. An infection near the bowel, anus, urinary tract, or genital area can also spread into nearby tissue. Sometimes there is no obvious wound.

Factors associated with a higher risk include:

Having a risk factor does not mean that this infection will occur. It can also affect people without known health problems.

Synergistic necrotizing fasciitis does not usually spread between people as a complete condition. Individual bacteria may sometimes pass from one person to another, but ordinary contact does not usually lead to a mixed deep-tissue infection.

- Diabetes, particularly when blood glucose is difficult to control
- Reduced blood flow to the legs or other tissues
- Kidney or liver disease
- Cancer or treatment that weakens the immune system
- Medicines that suppress the immune system
- Recent surgery, major injury, or childbirth
- Chronic wounds or pressure injuries
- Injecting non-prescribed drugs
- Severe obesity or poor nutrition

Diagnosis

Diagnosis begins with an urgent assessment of symptoms, the affected area, and how quickly the problem is changing. Clinicians may mark the edge of visible redness to check whether it is spreading.

Blood tests can show inflammation, infection, abnormal blood clotting, kidney injury, or raised lactate, which can be a sign that tissues are not receiving enough oxygen. Blood cultures may identify bacteria in the bloodstream.

CT or MRI scans may show fluid, swelling, gas, or changes along the fascia. However, an early scan can appear less serious than the actual infection. Imaging should not delay surgery when clinical findings strongly suggest a necrotizing infection.[1,2]

The most reliable confirmation comes from urgently examining the deeper tissues during surgery. Surgeons look for dead fascia, fluid, poor bleeding, or tissue that separates unusually easily. Deep tissue and fluid samples are sent for microscopy and culture to identify the bacteria.

A laboratory scoring tool called LRINEC is sometimes mentioned in relation to necrotizing fasciitis. It uses routine blood results. It cannot safely rule out the condition by itself, especially early in the illness.[3]

Treatment and management

Treatment takes place in hospital and usually involves a surgical team, infection specialists, and critical care staff.

Emergency surgery is the central treatment. The surgeon removes infected and dead tissue, a procedure called debridement. The wound is left open so that it can be checked. Further operations are often needed because tissue that initially appears healthy may later show damage.[1,2]

Broad intravenous antibiotics are started as soon as possible. They are chosen to cover several types of bacteria, including bacteria that grow without oxygen. Once culture results identify the organisms, the antibiotic plan may be narrowed or changed. The length of treatment is based on infection control, further surgery, and overall recovery rather than a single schedule for everyone.

Hospital support may include intravenous fluids, oxygen, pain relief, nutrition, blood-pressure support, and treatment for kidney or breathing problems. Some people need care in an intensive care unit.

If an arm or leg is extensively affected, amputation may occasionally be the safest way to stop the infection. This is not required in every case.

After the infection is controlled, wound care continues. Skin grafts, reconstructive surgery, or specialised dressings may be needed. Physiotherapy and occupational therapy can help restore movement and independence.

Hyperbaric oxygen, which involves breathing oxygen in a pressurised chamber, is sometimes used as an additional treatment. Evidence for its benefit is uncertain, and it must never delay surgery or antibiotics.[2,4] Intravenous immune globulin is also reserved for selected situations; it is not routine treatment for every mixed infection.

Complications

The infection can enter the bloodstream and cause sepsis, the body's extreme response to infection. Sepsis can lead to very low blood pressure, abnormal clotting, and failure of the kidneys, lungs, heart, or other organs.

Local complications may include:

A severe infection can be fatal despite treatment. Risk is greater when diagnosis or surgery is delayed, when a large area is affected, or when major organ failure develops.[3,5]

Recovery can also affect emotional health. Distressing memories, low mood, anxiety, sleep

problems, or concerns about body image may follow intensive care, major surgery, or amputation. Psychological support can be part of rehabilitation.

- Loss of skin, fat, muscle, or fascia
- Nerve injury, numbness, or weakness
- Reduced movement or limb loss
- Large scars or changes in appearance
- Long-lasting pain or sensitivity
- Problems with wound healing

Prevention

Synergistic necrotizing fasciitis cannot always be prevented. It may develop after a minor injury or without a clear entry point.

Basic wound care may reduce opportunities for bacteria to enter deeper tissue. Clean cuts and grazes with water, cover open wounds with a clean dressing, and follow the wound instructions provided after surgery. Avoid swimming or soaking an open wound unless the surgical or wound-care team says it is safe.

Seek timely care for wounds that become increasingly painful, swollen, warm, or discoloured. People with diabetes, poor circulation, reduced immunity, or chronic ulcers may need earlier assessment because infections can progress with fewer obvious skin changes.

Managing diabetes and long-term wounds may lower risk, but it cannot guarantee prevention. Avoid injecting non-prescribed drugs. If this is difficult, confidential harm-reduction and substance-use services can help reduce infection risks.

There is no vaccine that prevents all the bacteria involved in synergistic necrotizing fasciitis.

Living with it and what to expect

The immediate hospital phase may last days or weeks. Operations can continue until no further dead or infected tissue is found. The wound then begins a slower period of healing or reconstruction.

During the following months, dressing changes, surgical reviews, rehabilitation, and treatment of underlying conditions may take up significant time. Fatigue and loss of strength are common after severe infection or intensive care. Progress may be uneven rather than steady.

Some people recover with limited lasting effects. Others have reduced movement, numbness, chronic pain, altered appearance, or loss of a limb. Skin grafts and scars can change over time and may need continued care. The infection itself does not usually become a lifelong illness once it has been eliminated, but its physical and emotional effects can be long-lasting.

Routine follow-up may include wound checks, monitoring for recurrent infection, rehabilitation, and reviews after reconstructive surgery. Appointments are also a chance to discuss pain, sleep, mood, mobility, or returning to work.

A new episode is not inevitable. Future risk is influenced by ongoing wounds, circulation, diabetes, immune suppression, and other health conditions.

When to get medical help

Call emergency services or go to an emergency department immediately for severe or rapidly worsening pain with swelling, redness, or skin colour change. This is especially important when the pain seems much worse than the skin looks.

Other emergency signs include:

Do not wait for a routine appointment if these symptoms are present. Do not rely on home antibiotics or pain relief to see whether the problem settles.

After hospital treatment, contact the surgical or wound-care team promptly for increasing wound pain, new drainage, fever, worsening redness, or problems with a graft or dressing. Stable scars, mobility concerns, persistent tiredness, and ongoing rehabilitation needs can usually be discussed at planned follow-up unless they suddenly worsen.

- Rapidly spreading swelling or discolouration
- Blisters, black or grey skin, or fluid leaking from a wound
- Fever with confusion, faintness, fast breathing, or extreme weakness
- New numbness in a previously painful area
- A crackling feeling under the skin
- Rapid deterioration after surgery, injury, or a wound infection

Common questions

Is synergistic necrotizing fasciitis the same as flesh-eating disease?

"Flesh-eating disease" is an informal and imprecise term for necrotizing soft-tissue infections.

The bacteria do not literally eat flesh. They damage tissue through infection, toxins, inflammation, and loss of blood supply. "Synergistic" indicates that several bacterial species are acting together.

How quickly does it spread?

It can progress over hours, although the speed is not the same in every case. The deeper infection may extend beyond the visible skin changes. Rapidly worsening pain or swelling needs emergency assessment.

Can antibiotics cure it without surgery?

Usually not. Dead tissue has little blood flow, so antibiotics may not reach it well. Surgery removes infected and dead tissue, while antibiotics treat bacteria in living tissue and the bloodstream.

Is it contagious?

The condition itself is not usually passed from person to person. Some bacteria involved can spread through contact, but they do not commonly cause necrotizing infection in others. Hospitals may use infection-control precautions depending on the bacteria found.

Can it come back after treatment?

The treated infection can be eliminated, but another infection is possible, particularly if chronic wounds, poor circulation, diabetes, or immune suppression remain. New rapidly worsening symptoms require fresh assessment rather than waiting for a follow-up visit.

How long does recovery take?

Recovery may take weeks to many months. It is shaped by the amount of tissue removed, organ complications, wound reconstruction, and rehabilitation needs. Some people regain their previous level of function, while others have lasting physical or emotional effects.

Sources

- Stevens DL, Bisno AL, Chambers HF, et al. "Practice Guidelines for the Diagnosis and Management of Skin and Soft Tissue Infections: 2014 Update by the Infectious Diseases Society of America." Clinical Infectious Diseases . 2014;59(2):e10-e52.
- Sartelli M, Guirao X, Hardcastle TC, et al. "2018 WSES/SIS-E Consensus Conference: Recommendations for the Management of Skin and Soft-Tissue Infections." World Journal of Emergency Surgery . 2018;13:58.

- Bonne SL, Kadri SS. "Evaluation and Management of Necrotizing Soft Tissue Infections." Infectious Disease Clinics of North America . 2017;31(3):497-511.
- Hakkarainen TW, Kopari NM, Pham TN, Evans HL. "Necrotizing Soft Tissue Infections: Review and Current Concepts in Treatment, Systems of Care, and Outcomes." Current Problems in Surgery . 2014;51(8):344-362.
- Stevens DL, Bryant AE. "Necrotizing Soft-Tissue Infections." New England Journal of Medicine . 2017;377(23):2253-2265.
- Centers for Disease Control and Prevention. "About Necrotizing Fasciitis." Group A Streptococcal Diseases, reviewed 2025.