CONDITION

Tapeworm infection - beef or pork

An intestinal infection caused by eating beef or pork containing tapeworm larvae.

WHAT IT IS
Intestinal infection with an adult beef or pork tapeworm

BODY SYSTEM
Digestive system

MAIN SYMPTOMS
Tapeworm segments in stool, abdominal discomfort, nausea, weight loss

WHAT CAUSES IT
Eating undercooked beef or pork containing larvae

HOW IT IS DIAGNOSED
Stool microscopy and identification of tapeworm segments

MAIN TREATMENT
Antiparasitic medicine and follow-up stool testing

CAN IT BE CURED?
Yes - treatment usually clears the intestinal worm

IS IT AN EMERGENCY?
Sometimes - with seizures or severe neurological symptoms

LONG-TERM OUTLOOK
Usually excellent after treatment and confirmed clearance

Overview

Beef and pork tapeworm infections are intestinal infections caused by adult Taenia tapeworms. The beef tapeworm is called Taenia saginata . The pork tapeworm is called Taenia solium .

People become infected after eating raw or undercooked beef or pork containing tapeworm larvae. In the intestine, a larva attaches to the bowel wall and grows into a long, flat adult worm. The worm releases small body sections called segments, which may appear in bowel movements.

Many people have no symptoms or only mild digestive symptoms. However, identifying the species matters. A person carrying an adult pork tapeworm can release microscopic eggs that may cause a more serious infection called cysticercosis if the eggs are swallowed by that person or someone else.

Cysticercosis is not caused directly by eating pork. It happens when pork tapeworm eggs from human stool enter the mouth, usually through contaminated hands, food, or water. The eggs can develop into cysts in body tissues. Cysts affecting the brain are called neurocysticercosis.

Both beef and pork intestinal tapeworm infections can usually be cured with antiparasitic medicine.

Symptoms

Most intestinal tapeworm infections cause few or no symptoms. Some people first notice an infection when a pale, flat tapeworm segment appears in their stool, underwear, or bedding. Beef tapeworm segments may move for a short time after leaving the body.

Possible symptoms include:

These symptoms can have many other causes. Seeing something unusual in your stool does not always mean it is a tapeworm.

Symptoms such as seizures, severe headaches, confusion, vision changes, or new weakness are not typical of an adult worm confined to the intestine. They may suggest cysticercosis or another neurological problem and need urgent assessment.

- Passing tapeworm segments in bowel movements
- Mild abdominal pain or discomfort
- Nausea, diarrhoea, or constipation
- Changes in appetite
- Unexplained weight loss
- Tiredness or weakness

Causes and risk factors

Beef tapeworm infection develops when someone eats beef containing T. saginata larvae that have not been killed by proper cooking. Pork tapeworm infection develops in the same way after eating pork containing T. solium larvae.

The adult worm lives only in the human intestine. It produces eggs that leave the body in stool. Cattle or pigs become infected when they swallow these eggs from contaminated soil, water, or feed. Larvae then form cysts in the animals' muscles, completing the cycle when undercooked meat is eaten.

Risk is higher with:

The beef tapeworm does not normally cause cysticercosis in people. Pork tapeworm eggs can.

A person does not get cysticercosis simply from eating pork containing larvae. They get it by swallowing T. solium eggs from human faecal contamination. A person with an intestinal pork tapeworm may also contaminate their own hands and swallow eggs, especially if hand hygiene is poor.

- Eating raw or undercooked beef or pork
- Eating meat that has not been safely inspected or handled
- Living in or travelling to places where human waste may contaminate food or animal feed
- Limited access to clean water, toilets, and handwashing facilities
- Living with someone who carries a pork tapeworm

Diagnosis

A healthcare professional will usually ask about symptoms, recent travel, food exposure, and whether you have noticed segments in your stool.

The main test is examination of stool under a microscope for tapeworm eggs or segments. Eggs may not be released every day, so samples collected on different days may be needed.

Beef and pork tapeworm eggs look alike under a microscope. Identifying the species may require examination of an intact segment or, less commonly, the worm's head. If you find a possible segment, a healthcare service can explain how to collect and transport it safely.

Blood tests do not usually confirm a simple adult intestinal tapeworm infection.

If pork tapeworm infection is possible and you have neurological symptoms, further assessment may be needed before treatment. Tests for cysticercosis can include brain imaging with a CT or MRI scan, an eye examination, and selected blood tests. A stool test cannot rule out cysts elsewhere in the body.

Treatment and management

Intestinal beef or pork tapeworm infection is treated with an antiparasitic medicine. Praziquantel is commonly used. Niclosamide or albendazole may be alternatives in some settings. The choice depends on local availability, the tapeworm species, other health conditions, and whether cysticercosis is a concern.

Treatment is often brief, but it should be arranged through a healthcare professional. This is particularly important if pork tapeworm infection is possible. Killing tapeworm larvae in the brain or other tissues can trigger inflammation, so suspected cysticercosis needs a different assessment and treatment plan.

You may or may not see the worm pass after treatment. Not seeing it does not mean treatment failed. Follow-up stool testing is used to check that eggs and segments are no longer present. The US Centers for Disease Control and Prevention advises repeat stool examination after treatment, including at one and three months.

During and after treatment, careful handwashing helps prevent pork tapeworm eggs from spreading. Wash your hands with soap and water after using the toilet, after changing nappies, and before preparing or eating food.

If T. solium is confirmed, public health or healthcare teams may consider whether close household contacts need assessment. This is because one person carrying an adult pork tapeworm can expose others to eggs.

Complications

Complications from an adult intestinal tapeworm are uncommon. A very large worm or a migrating segment can rarely contribute to blockage or inflammation involving the bowel, appendix, bile ducts, or pancreatic duct.

The main concern with T. solium is cysticercosis. After pork tapeworm eggs are swallowed, larvae may spread through the bloodstream and form cysts under the skin, in muscles, in the eyes, or in the nervous system.

Neurocysticercosis can cause:

Symptoms may develop months or years after exposure. Their type and severity are influenced by the number, location, and stage of the cysts.

Cysticercosis requires specialist assessment. Treatment may involve antiparasitic medicines, medicines to control inflammation or seizures, and sometimes surgery. It is not managed in the same way as an uncomplicated intestinal tapeworm.

- Seizures
- Persistent or severe headaches
- Increased pressure around the brain
- Problems with balance, speech, movement, or vision

- Confusion or changes in alertness

Prevention

Beef and pork tapeworm infections are preventable in many cases.

Cook beef and pork thoroughly, using a food thermometer and current local food-safety guidance. Colour alone does not reliably show whether meat has reached a safe temperature. Avoid tasting raw meat while preparing it.

Other protective steps include:

Freezing can kill some tapeworm larvae under controlled conditions, but household freezing is not a reliable substitute for safe cooking because temperature and storage time differ.

If you have been diagnosed with a pork tapeworm, strict hand hygiene is especially important until treatment and follow-up are complete. It reduces the chance of spreading eggs to yourself or other people.

- Wash hands with soap and water after using the toilet
- Wash hands before preparing food or eating
- Use safe drinking water
- Wash or peel produce when contamination is possible
- Keep raw meat separate from ready-to-eat food
- Buy meat from regulated or reliably inspected sources
- Use proper toilets and safe sewage disposal

Living with it and what to expect

An untreated adult tapeworm can remain in the intestine for years. Symptoms may stay mild, come and go, or remain absent. Passing segments can continue while the worm is present.

Most people can continue their usual activities while being assessed and treated. The infection is not spread through coughing, touching, or ordinary social contact. The concern with pork tapeworm is microscopic egg contamination after using the toilet, which is why handwashing matters.

After effective treatment, digestive symptoms and passage of segments should settle. Follow-up stool testing is important because symptoms alone cannot confirm that the worm

has gone. Reinfection is possible if you later eat contaminated, undercooked meat. Treatment does not provide lasting immunity.

Cysticercosis has a different course. Cysts can remain for years, and neurological symptoms may need longer-term monitoring or treatment. The outlook is shaped by where the cysts are located and whether they have caused seizures, pressure, or tissue injury. Many people can manage the condition with appropriate care, but no single course can be promised.

When to get medical help

Arrange a routine medical appointment if:

Seek urgent medical advice if you develop a new persistent headache, repeated vomiting, vision changes, unusual balance problems, or new numbness or weakness.

Seek emergency care for:

These symptoms can have several causes. They need prompt assessment rather than being assumed to come from a tapeworm.

- You notice a flat, pale segment or worm-like material in your stool
- You have ongoing digestive symptoms after eating raw or undercooked beef or pork
- You have unexplained weight loss
- Someone in your household has been diagnosed with pork tapeworm
- Symptoms or segments continue after treatment
- You are due for follow-up stool testing
- A first seizure or repeated seizures
- Loss of consciousness
- Sudden confusion, difficulty speaking, or one-sided weakness
- A severe headache with reduced alertness
- Severe abdominal pain with swelling, persistent vomiting, or inability to pass stool or gas

Common questions

Can a beef or pork tapeworm go away on its own?

An adult tapeworm can survive in the intestine for years, so waiting for it to disappear is not reliable. Antiparasitic treatment is usually effective, and follow-up testing checks whether it

has cleared.

How will I know whether the tapeworm is gone?

You may not see the whole worm pass. The most reliable check is follow-up examination of stool samples for eggs or segments. Feeling better alone does not confirm clearance.

Can I catch a tapeworm from another person?

You do not get an adult beef or pork tapeworm through ordinary contact. However, a person carrying a pork tapeworm can release eggs in stool. If those eggs contaminate hands, food, or water and are swallowed, they can cause cysticercosis.

Does eating pork cause tapeworm cysts in the brain?

Eating undercooked pork containing larvae can cause an adult intestinal pork tapeworm. Brain cysts develop after swallowing pork tapeworm eggs from human faecal contamination, not directly from eating pork.

Can pets give me beef or pork tapeworm?

Beef and pork tapeworms mainly follow a human-cattle or human-pig life cycle. Dogs and cats can carry other tapeworm species, so pets should still receive appropriate veterinary care and parasite control.

Can I get tapeworm infection again after treatment?

Yes. Treatment clears the current intestinal worm but does not create lasting protection. Eating contaminated undercooked meat can cause another infection.

Sources

- Centers for Disease Control and Prevention. Clinical Overview of Taeniasis
- Centers for Disease Control and Prevention. Clinical Treatment of Taeniasis
- Centers for Disease Control and Prevention. DPDx: Taeniasis
- World Health Organization. Taeniasis and cysticercosis
- White AC Jr, Coyle CM, Rajshekhar V, et al. Diagnosis and Treatment of Neurocysticercosis: 2017 Clinical Practice Guidelines by the IDSA and ASTMH. Clinical Infectious Diseases. 2018;66(8):e49-e75
- Garcia HH, Nash TE, Del Brutto OH. Clinical symptoms, diagnosis, and treatment of neurocysticercosis. The Lancet Neurology. 2014;13(12):1202-1215 70094-8)