CONDITION

Tar remover poisoning

Tar remover poisoning is harmful exposure to the solvents used in products that dissolve tar, grease, or asphalt.

WHAT IT IS
Harmful exposure to solvents used to dissolve tar

BODY SYSTEM
Lungs, brain, skin, eyes, heart, and digestive system

MAIN SYMPTOMS
Coughing, breathing trouble, dizziness, nausea

WHAT CAUSES IT
Swallowing, breathing, or skin contact with solvent chemicals

HOW IT IS DIAGNOSED
Exposure history, examination, oxygen monitoring, and targeted tests

MAIN TREATMENT
Decontamination, observation, and supportive hospital care

CAN IT BE CURED?
Sometimes - severe injuries may not resolve

IS IT AN EMERGENCY?
Sometimes - breathing trouble, collapse, seizure, or severe drowsiness

LONG-TERM OUTLOOK
Mild exposures often resolve; severe organ injury can persist

Overview

Tar remover poisoning happens when a tar-removing product harms the body after being swallowed, breathed in, splashed into the eyes, or left on the skin. Products sold for vehicles, tools, roofing, or industrial cleaning may contain strong solvents that dissolve tar, asphalt, oil, or grease.

The ingredients are not the same in every product. Some contain petroleum-based hydrocarbons, such as mineral spirits. Others may contain chemicals such as toluene, xylene, or chlorinated solvents. The label or safety data sheet is important because the ingredient affects the risks and treatment.

Swallowing a tar remover is especially concerning if the liquid enters the lungs. This is called aspiration. It can happen during swallowing or vomiting and may cause chemical pneumonitis,

meaning inflammation and injury in the lungs.

Breathing concentrated fumes can affect the brain and breathing. Large or prolonged exposures may also affect the heart or other organs. Skin and eye contact usually causes local irritation, but some solvents can also pass through the skin.

A suspected exposure should be discussed promptly with a poison centre, even if symptoms are mild or have not started. Serious symptoms require emergency care.

Symptoms

Symptoms depend on the product, amount, route of exposure, ventilation, and time since the exposure. They may begin quickly, but lung symptoms can sometimes develop or worsen over several hours.

After swallowing tar remover, possible symptoms include:

Coughing after swallowing is important because it may mean some liquid entered the airway. Vomiting can increase the chance of aspiration.

Breathing fumes may cause headache, dizziness, nausea, eye or throat irritation, sleepiness, or confusion. A heavy exposure in an enclosed space can lead to poor coordination, loss of consciousness, slowed breathing, or seizures.

Lung injury may cause ongoing coughing, chest discomfort, wheezing, fast breathing, or shortness of breath. The lips or skin may look bluish or grey when oxygen levels are dangerously low.

Skin exposure can cause redness, dryness, cracking, pain, or a burning feeling. Long contact under soaked clothing may cause a chemical burn. Eye exposure can cause pain, watering, redness, blurred vision, and sensitivity to light.

Some solvents can disturb the heart's rhythm. This may feel like a racing or irregular heartbeat, weakness, light-headedness, or fainting. The exact risks depend on the ingredients.

- Burning or irritation in the mouth and throat
- Nausea, vomiting, or stomach pain
- Coughing, choking, or gagging
- Drowsiness, dizziness, confusion, or poor coordination

Causes and risk factors

Tar remover poisoning is caused by contact with the chemicals in the product. Common situations include accidental swallowing, using a product in a poorly ventilated area, transferring it into an unlabelled drink container, or spilling it onto clothing or skin.

Young children may swallow a product that has been left within reach. Adults may be exposed while cleaning vehicles, tools, machinery, flooring, or roofing materials. Workplace exposures can occur when ventilation and protective equipment are inadequate.

Risk may be greater when:

Mixing tar remover with other cleaners can create additional hazards. Never assume two cleaning products are safe to combine.

- The product is used in an enclosed space
- A large amount is spilled or swallowed
- Contaminated clothing is not removed promptly
- The person vomits after swallowing the product
- The exposure lasts a long time
- Alcohol, sedatives, or other substances add to drowsiness
- Existing lung or heart problems make the body less able to tolerate exposure

Diagnosis

There is no single test that confirms every case of tar remover poisoning. Diagnosis is usually based on the product involved, how the exposure happened, symptoms, and a physical examination.

The product container, ingredient list, or safety data sheet can help the poison centre and hospital identify the likely chemicals. Useful details include when the exposure happened, whether the product was swallowed or inhaled, and whether coughing or vomiting occurred.

A healthcare team may check breathing rate, alertness, heart rhythm, temperature, blood pressure, and blood oxygen using a finger sensor. An electrocardiogram, or ECG, records the heart's electrical activity.

A chest X-ray may be used when there is coughing, low oxygen, or concern that liquid entered the lungs. Early X-rays can sometimes appear normal even when lung irritation is developing, so observation and repeat assessment may be needed.

Blood tests may check acid levels, electrolytes, blood sugar, or liver and kidney function after a significant exposure. Testing for particular chemicals is only useful in selected cases. For

example, products containing methylene chloride may lead clinicians to test for carbon monoxide exposure because the body converts part of that chemical into carbon monoxide.

Treatment and management

The first step is to stop the exposure without putting anyone else at risk. Move away from fumes and into fresh air if this can be done safely. Do not enter an enclosed contaminated area without appropriate rescue equipment.

Remove clothing soaked with the product. Wash exposed skin with plenty of lukewarm running water and mild soap. Rinse an exposed eye with clean, lukewarm water, removing contact lenses if they come out easily. A poison centre can advise how long rinsing should continue for the specific product.

Do not make someone vomit after swallowing tar remover. Vomiting can draw the liquid into the lungs. Do not give food, milk, oil, alcohol, activated charcoal, or another home remedy unless a poison specialist specifically recommends it.

Hospital treatment is supportive, meaning it protects breathing and other body functions while the chemical is cleared. It may include oxygen, breathing support, intravenous fluids, treatment for seizures, and heart monitoring. Wheezing may be treated with an inhaled medicine that opens the airways.

Activated charcoal and stomach pumping are generally avoided for hydrocarbon products because they can increase aspiration risk and often remove little chemical. A toxicology specialist may consider different measures for an unusual ingredient or a very specific situation.

There is no single antidote for most tar remover mixtures. Antibiotics like amoxicillin or doxycycline do not treat chemical lung inflammation and are not routinely used unless there is evidence of an infection. Steroids like prednisone or hydrocortisone are also not routine treatment for hydrocarbon aspiration.

People without symptoms may still need a period of observation because breathing problems can be delayed. The required monitoring depends on the product, symptoms, and route of exposure.

Complications

The main serious complication after swallowing is aspiration pneumonitis. The solvent irritates lung tissue and can cause inflammation, fluid leakage, and reduced oxygen. Severe cases may

progress to respiratory failure, meaning the lungs cannot provide enough oxygen without support.

Other possible complications include an abnormal heart rhythm, seizures, loss of consciousness, or injury caused by falling while dizzy. Prolonged skin contact can produce dermatitis or a chemical burn.

Certain ingredients can affect the liver, kidneys, blood, or nervous system after a substantial exposure. These effects are product-specific and are not expected after every tar remover exposure.

Long-term breathing problems or brain injury are possible after severe poisoning, particularly when there was prolonged low oxygen. Mild exposures that are treated promptly are less likely to leave lasting effects.

Prevention

Most tar remover exposures can be prevented. Keep the product in its original, labelled container with the cap secured. Never pour it into a cup, bottle, food jar, or other container that could be mistaken for a drink.

Store tar remover away from children, food, medicines, heat, sparks, and flames. Many solvent vapours can catch fire.

Use the product only as directed on the label. Work outdoors or in an area with effective ventilation. Household fans may not make a small enclosed space safe if they only move contaminated air around.

Wear the gloves and eye protection recommended by the manufacturer. Replace contaminated clothing promptly and wash exposed skin. Do not eat, drink, smoke, or vape while handling the product.

For workplace use, follow the safety data sheet, ventilation controls, protective-equipment requirements, and spill procedures. Dispose of leftover product and contaminated materials according to local hazardous-waste rules.

Living with it and what to expect

Tar remover poisoning is usually an acute event rather than an ongoing condition. What happens over the following days or months depends mainly on the chemical, the amount of exposure, whether liquid entered the lungs, and how severely breathing or consciousness was

affected.

Minor skin or eye irritation often improves after thorough rinsing, although dryness or soreness may last for a time. Mild headache or dizziness from fumes may improve after leaving the exposure area. Symptoms that continue or return need medical review.

After aspiration, coughing and breathing problems may develop during the first several hours. Recovery from lung inflammation can take days or longer. Severe lung injury may require intensive care and a longer period of reduced stamina.

Follow discharge instructions about activity and follow-up. A routine follow-up appointment may be recommended after significant lung injury, an abnormal chest X-ray, ongoing cough, or exposure to a chemical that can affect the liver, kidneys, or nervous system.

Some people feel anxious about using cleaning chemicals after a poisoning event. Reviewing safer storage, ventilation, and protective equipment can help reduce the chance of another exposure.

When to get medical help

Call emergency services now if the exposed person has trouble breathing, collapses, cannot be awakened normally, has a seizure, becomes severely confused, or develops blue or grey lips or skin. Move away from fumes only if it is safe to do so.

Do not wait for symptoms after a known or suspected swallowing. Contact a poison centre promptly for advice based on the exact product. In the United States, Poison Help is available at 1-800-222-1222. Elsewhere, use your national or regional poison information service.

Get urgent medical assessment for repeated coughing or choking after swallowing, persistent vomiting, chest pain, wheezing, increasing drowsiness, fainting, or an irregular heartbeat. Eye pain or blurred vision that continues after rinsing also needs prompt assessment.

After initial treatment, seek urgent reassessment if breathing becomes harder, cough worsens, fever develops with respiratory symptoms, alertness changes, or symptoms return. Routine follow-up is appropriate for stable, gradually improving symptoms when this was recommended at discharge.

Common questions

What should I do if I swallowed tar remover?

Do not make yourself vomit. Rinse your mouth gently, move away from the product, and

contact a poison centre immediately. Have the container or label available. Call emergency services for coughing with breathing trouble, severe drowsiness, collapse, or a seizure.

Can symptoms start hours after the exposure?

Yes. Lung irritation can develop after the initial swallowing or aspiration, even if you first feel well. This is why poison specialists may recommend observation rather than relying only on how you feel immediately afterward.

Will drinking milk help?

Milk does not neutralise tar remover and may not be safe if you are drowsy, vomiting, or at risk of aspiration. Do not take food, drinks, charcoal, or home remedies unless a poison specialist advises it.

Is smelling tar remover enough to cause poisoning?

A brief smell in a well-ventilated place may cause irritation without serious poisoning. Concentrated fumes, prolonged exposure, or use in an enclosed area can cause headache, dizziness, confusion, and breathing problems. Leave the area if symptoms begin.

What if tar remover gets on my skin?

Remove contaminated clothing and wash the skin thoroughly with lukewarm running water and mild soap. Contact a poison centre if there is ongoing pain, redness, blistering, a large exposed area, or concern about a highly concentrated product.

Can tar remover poisoning cause permanent damage?

It can after a severe exposure, especially if there is major lung injury, prolonged unconsciousness, or low oxygen. Lasting harm is less likely after a mild exposure that resolves without breathing or neurological problems.

Sources

- U.S. National Library of Medicine. "Tar Remover Poisoning." MedlinePlus Medical Encyclopedia
- Merck Manual Professional Edition. "Hydrocarbon Poisoning." Injuries; Poisoning
- Agency for Toxic Substances and Disease Registry. "Medical Management Guidelines for Toluene." U.S. Department of Health and Human Services
- Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. "Toluene." NIOSH Pocket Guide to Chemical Hazards

- Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. "Xylene (All Isomers)." NIOSH Pocket Guide to Chemical Hazards
- Royal Children's Hospital Melbourne. "Hydrocarbon Poisoning." Clinical Practice Guidelines