CONDITION

Tolmetin overdose

Tolmetin overdose happens when someone takes more of this anti-inflammatory medicine than their body can safely handle.

WHAT IT IS
Excess tolmetin causing harmful effects

BODY SYSTEM
Digestive system, kidneys, nervous system, and circulation

MAIN SYMPTOMS
Nausea, stomach pain, drowsiness, vomiting

WHAT CAUSES IT
Taking more tolmetin than the body can safely handle

HOW IT IS DIAGNOSED
Exposure history, examination, and blood and urine tests

MAIN TREATMENT
Poison-centre guidance and supportive medical care

CAN IT BE CURED?
Yes - most recover with prompt supportive care

IS IT AN EMERGENCY?
Sometimes - breathing trouble, collapse, seizure, or inability to wake

LONG-TERM OUTLOOK
Most people recover without lasting effects

Overview

Tolmetin is a prescription nonsteroidal anti-inflammatory drug like ibuprofen or naproxen , or NSAID. It reduces pain, swelling, and stiffness, including symptoms caused by some forms of arthritis.

A tolmetin overdose happens when someone takes more than their body can safely process. This may involve one large amount, several extra doses, or repeated dosing errors. It can be accidental or intentional.

Too much tolmetin can irritate the stomach and intestines, reduce blood flow through the kidneys, and affect the brain and breathing. Most small overdoses cause mild or no symptoms, but the amount taken does not always predict the outcome.

There is no specific antidote that cancels tolmetin's effects. Treatment supports breathing, circulation, kidney function, and other body systems while the medicine is cleared.

If an overdose may have happened, contact a poison centre promptly even if there are no symptoms. In the United States, Poison Help is available at 1-800-222-1222 . Elsewhere, use your local poison information service. Call emergency services if the person collapses, has a seizure, has trouble breathing, or cannot be awakened.

Symptoms

Symptoms often begin within several hours, although kidney problems or bleeding may become clear later. Early symptoms can be similar to common medicine side effects.

Mild or moderate symptoms may include:

More serious poisoning can affect several body systems. Possible signs include confusion, marked sleepiness, very little urine, low blood pressure, rapid breathing, or blood in vomit or stool.

Severe overdoses have rarely caused seizures, loss of consciousness, slowed or difficult breathing, acute kidney injury, and major disturbances in the blood's acid balance. An allergic-type reaction, including swelling or wheezing, is also possible.

Do not wait for symptoms before seeking poison-centre advice. Someone who initially feels well may still need assessment based on the amount, timing, age, health conditions, and other substances involved.

- Nausea, vomiting, or upper stomach pain
- Heartburn or indigestion
- Drowsiness, dizziness, or unusual tiredness
- Headache or ringing in the ears

Causes and risk factors

The direct cause is taking too much tolmetin. Accidental overdose can happen when someone repeats a dose after forgetting they already took it, misunderstands the label, or uses more than one container.

Risk can also increase when tolmetin is taken with another NSAID. Examples include ibuprofen, naproxen, diclofenac, and high-dose aspirin. Many cold, flu, and pain products contain an

NSAID, so it is possible to combine them without realizing it.

The effects may be more serious in someone who has kidney disease, heart failure, stomach ulcers, or previous digestive-tract bleeding. Dehydration can make the kidneys more vulnerable because they already receive less blood flow.

Older adults and very young children may have less reserve when dealing with poisoning. Taking alcohol, sedatives, or other medicines at the same time may change the symptoms or increase the chance of complications.

An intentional overdose may occur during a mental health crisis. This requires both medical assessment and compassionate mental health support, even if physical symptoms seem mild.

Diagnosis

There is no routine blood test that measures tolmetin and reliably shows how severe an overdose will be. Diagnosis is mainly based on what was taken, how much may have been taken, when it happened, and the person's symptoms.

Bring the medicine container or a clear photograph of its label if it is safe to do so. Healthcare professionals may ask about other prescription medicines, nonprescription products, alcohol, or recreational substances. This helps identify a mixed overdose, which may require different treatment.

An examination usually includes checking alertness, breathing, pulse, blood pressure, temperature, and hydration.

Tests may include:

Testing is guided by the situation. Someone with a minor dosing error and no symptoms may not need the same investigations as someone who is drowsy, dehydrated, or unwell.

- Blood tests for kidney function, salts, blood sugar, and acid balance
- A blood count to look for significant bleeding
- Urine testing and measurement of urine output
- An electrocardiogram, or ECG, to check heart rhythm
- Tests for other common overdoses when the substances are uncertain

Treatment and management

Contact a poison centre rather than trying to manage a possible overdose alone. Poison

specialists can assess the specific product, amount, timing, symptoms, age, and medical history. They may recommend monitoring at home or evaluation in an emergency department.

Do not make the person vomit. Vomiting does not reliably remove the medicine and can lead to choking, especially if the person becomes drowsy. Do not give activated charcoal unless a poison specialist or healthcare professional directs it.

In hospital, treatment is supportive. This means treating the effects of the overdose while the body clears tolmetin. Care may include monitoring, intravenous fluids, medicine for nausea or stomach irritation, oxygen, and treatment of abnormal blood pressure or seizures.

Activated charcoal may be considered by clinicians after some recent, significant overdoses. It binds medicine inside the digestive tract. It is not suitable in every case, particularly when someone cannot protect their airway.

People with suspected digestive bleeding may need repeated blood counts and additional treatment. Kidney function and urine output may also be followed because kidney injury can develop after the initial symptoms.

Tolmetin is strongly attached to proteins in the blood. For this reason, dialysis does not usually remove much of the drug. Dialysis may still be used to support someone who develops severe kidney failure or a dangerous acid or salt imbalance.

Complications

Most people recover, particularly after a small overdose and early assessment. Complications are more likely after a large amount, a delayed presentation, dehydration, or an overdose involving several substances.

Possible complications include:

Acute kidney injury is a sudden decline in the kidneys' ability to filter the blood. It may improve with supportive treatment, but severe injury can require longer monitoring and, occasionally, dialysis.

Major digestive bleeding may cause vomiting of blood, black sticky stools, weakness, or fainting. Significant blood loss may require hospital treatment.

Lasting problems are uncommon after a mild overdose. A severe episode can sometimes leave ongoing kidney problems or other effects related to low oxygen, bleeding, or prolonged loss of consciousness.

- Stomach or intestinal bleeding

- Acute kidney injury
- Severe dehydration or low blood pressure
- Abnormal blood acidity
- Seizures, coma, or breathing failure
- Choking on vomit while drowsy

Prevention

Tolmetin overdose is often preventable. Keep the medicine in its original, labelled container rather than moving it into an unmarked bottle or bag. Store it where children and other people cannot access it.

Use a written schedule, medication organizer, or reminder system if missed or repeated doses are a concern. A medication organizer should still be stored securely.

Before using another pain, cold, or flu product, check whether it contains an NSAID. Do not combine tolmetin with another NSAID unless a clinician has specifically advised it.

Do not take an extra dose to make up for a forgotten one without checking the prescription instructions or asking a pharmacist. If a dosing error occurs, contact a poison centre rather than waiting for symptoms.

If there is concern that someone may take medicine intentionally, secure the available supply and seek mental health support. During an immediate crisis, stay with the person if it is safe and contact emergency or crisis services.

Living with it and what to expect

Tolmetin overdose is usually an acute event rather than a long-term condition. The first several hours focus on watching for drowsiness, vomiting, stomach pain, breathing problems, and changes in blood pressure.

After a minor exposure, a poison centre may recommend observation at home with clear instructions about what symptoms require medical care. A person assessed in hospital may be discharged once they remain well and any necessary test results are reassuring.

Recovery from a more significant overdose may take several days. Kidney function, hydration, and blood counts may need to be followed after the immediate symptoms improve. Routine follow-up is especially important if there was kidney injury, digestive bleeding, or an existing

kidney or stomach condition.

Do not restart tolmetin or substitute another NSAID after an overdose until a healthcare professional has reviewed what happened. Future pain treatment may need to account for any kidney or digestive complications.

If the overdose was intentional, physical recovery is only part of care. A mental health assessment can help address the crisis, improve safety, and arrange ongoing support. This should be offered without blame or judgment.

When to get medical help

Contact a poison centre promptly after any suspected tolmetin overdose, even if the person feels well. Have the container, strength, possible amount, and approximate time available, but do not delay the call if some details are unknown.

Call emergency services now if the person:

Urgent medical assessment is also appropriate for repeated vomiting, worsening stomach pain, very little urine, marked drowsiness, or an overdose involving other medicines.

Routine follow-up may be needed after discharge to repeat kidney function or blood-count tests. Keep that follow-up even if symptoms have resolved, because some complications are found through testing rather than how you feel.

- Has trouble breathing, wheezing, or blue or grey lips
- Collapses, has a seizure, or cannot be awakened
- Is severely confused or becoming rapidly less responsive
- Vomits blood or passes black, sticky stool
- Has severe weakness, fainting, or signs of shock

Common questions

I took one extra tolmetin dose. Is that an overdose?

It is a dosing error, but it does not always cause poisoning. The risk depends on the tablet strength, total amount, timing, age, health conditions, and other medicines. Contact a poison centre for advice specific to the situation.

How much tolmetin is dangerous?

There is no single amount that is safe to use as a cutoff for everyone. Do not rely on an online

dose calculation. A poison centre can assess the actual product and circumstances without waiting for symptoms.

What should I do if I feel fine?

Call a poison centre promptly and follow its advice. Do not take more tolmetin or another NSAID while waiting. Feeling well at first does not rule out later stomach or kidney effects.

Can food, milk, or water stop the overdose?

No food or drink neutralizes tolmetin. Do not force fluids or give anything by mouth to someone who is very drowsy, vomiting repeatedly, or having trouble swallowing. Follow poison-centre instructions.

Will dialysis remove tolmetin?

Usually not effectively, because most tolmetin in the bloodstream is attached to proteins that dialysis cannot easily remove. Dialysis may still support the body if severe kidney failure or a dangerous chemical imbalance develops.

Can I take tolmetin again after an overdose?

That decision should be made after medical review. It may be reasonable after a minor error, but kidney injury, digestive bleeding, or an intentional overdose can change whether tolmetin remains appropriate.

Sources

- DailyMed, U.S. National Library of Medicine. Tolmetin sodium capsule-prescribing information . Sections on overdose, warnings, and clinical pharmacology.
- MedlinePlus, U.S. National Library of Medicine. Tolmetin: Drug Information .
- Hunter LJ, Wood DM, Dargan PI. "The patterns of toxicity and management of acute nonsteroidal anti-inflammatory drug overdose." Open Access Emergency Medicine . 2011;3:39-48.
- Royal Children's Hospital Melbourne. Clinical Practice Guideline: Nonsteroidal Anti-inflammatory Drug Poisoning .
- Chyka PA, Seger D, Krenzelok EP, Vale JA. "Position paper: Single-dose activated charcoal." Clinical Toxicology . 2005;43(2):61-87.
- Poison Help, Health Resources and Services Administration. Get Help in a Poison Emergency .