CONDITION

Torsion dystonia

Torsion dystonia causes repeated muscle contractions that twist the body into unwanted movements or postures.

WHAT IT IS
Movement condition causing twisting movements and abnormal postures

BODY SYSTEM
Nervous system and muscles

MAIN SYMPTOMS
Twisting, abnormal postures, muscle spasms, pain

WHAT CAUSES IT
Inherited changes, brain injury, medicines, or unknown causes

HOW IT IS DIAGNOSED
Neurological examination, supported by tests for underlying causes

MAIN TREATMENT
Movement therapy, medicines, injections, or deep brain stimulation

CAN IT BE CURED?
No - treatment can reduce symptoms

IS IT AN EMERGENCY?
Sometimes - breathing difficulty or severe continuous spasms

LONG-TERM OUTLOOK
Often lifelong; severity and spread differ by pattern

Overview

Torsion dystonia is a form of dystonia, a movement condition in which muscles contract when you do not intend them to. These contractions can pull part of the body into a twisting movement or an unusual posture.

"Torsion" describes the twisting quality of the movement. The term is used less often today because dystonia is now classified by the body areas affected, the age symptoms began, and the underlying cause. You may still see torsion dystonia in older medical records or information about inherited, widespread dystonia.

Dystonia can affect one small area, several connected areas, or much of the body. For example, it may involve the neck, eyelids, jaw, voice, hand, foot, or trunk. Generalized dystonia affects the trunk and at least two other body areas.

The condition matters because sustained contractions can interfere with walking, speaking, eating, writing, driving, sleep, or work. They may also cause pain. Dystonia does not usually affect intelligence, but its physical and emotional effects can be significant.

Symptoms

Dystonia may cause slow twisting, repeated movements, fixed postures, or a mixture of these. Symptoms can be mild and intermittent or more persistent.

Common features include:

Symptoms may first appear during a specific action, such as writing, playing an instrument, speaking, or running. Over time, they may occur during other activities or at rest. Stress, tiredness, movement, and strong emotions can temporarily make symptoms more noticeable, but they do not mean the symptoms are imagined.

Some people find that lightly touching the face or another body area briefly reduces a posture. This is called a "sensory trick." Symptoms often improve during sleep, although this is not true for everyone.

- A foot that turns inward or drags while walking
- A neck that turns, tilts, or pulls backward
- Repeated blinking or forceful eyelid closure
- A hand that curls or tightens during writing or another task
- Twisting of the trunk, arms, or legs
- Shaking that may look like a tremor
- Muscle pain, stiffness, or tiredness

Causes and risk factors

Dystonia results from changes in the brain networks that control and coordinate movement. It is not simply a problem with the affected muscles.

In many people, no single cause is found. This is sometimes called idiopathic dystonia. "Idiopathic" means that the cause remains unknown after assessment.

Some forms are inherited. Changes in genes such as TOR1A can cause early-onset dystonia, but many other genes are also linked to dystonia. Having a gene change does not always mean symptoms will develop. The pattern of inheritance and the chance of passing it on differ

between genes.

Dystonia can also develop because of another condition or exposure, including:

Dystonia caused by dopamine-blocking medicine may appear during treatment or after long-term use. Do not stop a prescribed medicine suddenly. A clinician can review its risks and possible alternatives.

Early-onset dystonia often begins in an arm or leg and has a greater chance of spreading to other body areas. Dystonia beginning in adulthood more often starts in the neck, face, or hand and may remain more localized.

- Brain injury, stroke, or reduced oxygen reaching the brain
- Cerebral palsy or certain neurological conditions
- Wilson disease, which causes copper to build up in the body
- Some infections or immune-related conditions affecting the brain
- Medicines that block dopamine, including certain antipsychotic and anti-nausea medicines

Diagnosis

There is no single scan or blood test that confirms every form of torsion dystonia. Diagnosis is mainly clinical, meaning it is based on your symptoms, medical history, and a neurological examination.

A clinician may watch you walk, write, speak, or perform an action that brings on the movement. They will look at which body areas are affected, whether movements are patterned or twisting, and whether there are signs of another neurological condition.

You may be asked about:

Blood or urine tests may be used to look for treatable causes, such as Wilson disease. Brain MRI may be appropriate if symptoms started suddenly, follow an injury, or occur with other neurological signs. Genetic testing may identify an inherited cause, especially when symptoms began young or affect several relatives.

Some people are offered a supervised trial of levodopa, a medicine that replaces a brain chemical called dopamine. A strong response can help identify dopa-responsive dystonia. Tests also help distinguish dystonia from muscle cramps, tics, seizures, joint problems, and other movement conditions.

- When the movements began and how they have changed

- Medicines you currently take or previously used
- Birth complications, injuries, infections, or other health conditions
- Relatives with dystonia, tremor, or unexplained movement problems

Treatment and management

Treatment aims to reduce unwanted movements, pain, and the effect on daily life. The most suitable approach is influenced by the body areas involved, the cause, and how much symptoms interfere with your activities. More than one treatment may be used.

Botulinum toxin injections are commonly used for dystonia affecting a specific area, such as the neck, eyelids, jaw, or hand. The medicine weakens selected overactive muscles for a limited period, so repeat treatment is usually needed. Possible effects include temporary weakness. Injections around the neck or mouth can sometimes affect swallowing or speech.

Oral medicines may be considered when several body areas are affected. Options include medicines that alter acetylcholine, dopamine, or other chemical signals in the nervous system. Examples include anticholinergic medicines, baclofen, clonazepam, and levodopa for certain forms. Benefits and side effects differ, and some medicines must be adjusted gradually rather than stopped suddenly.

Physical therapy can help you maintain movement, balance, comfort, and safe ways of completing tasks. It does not remove the neurological cause, and repeatedly forcing a dystonic body part into a "normal" position may worsen pain. Occupational therapy can help with writing, computer use, dressing, mobility, equipment, and workplace changes.

Speech and language therapy may help if dystonia affects your voice, speech, chewing, or swallowing. Psychological support can be useful when pain, visible movements, work problems, or social reactions affect your mood or confidence.

Deep brain stimulation may be considered for severe dystonia that has not improved enough with other treatment. A surgeon places thin electrical leads in selected brain areas, usually those involved in movement control. A device under the skin supplies adjustable electrical stimulation. Improvement may develop gradually, and ongoing programming is required. Surgery has risks, including infection, bleeding, and device problems.

Complications

Repeated contractions can cause pain, exhaustion, falls, joint strain, or reduced movement. A

fixed posture may eventually shorten muscles and tendons, leading to contractures that make a joint difficult to straighten.

Dystonia affecting the mouth, throat, or neck may make speaking, chewing, or swallowing harder. Swallowing problems can increase the chance of food or liquid entering the airway.

Symptoms may also affect employment, education, driving, relationships, and emotional well-being. Anxiety and depression can occur and deserve direct care rather than being treated as an unavoidable part of dystonia.

Rarely, severe generalized dystonia becomes continuous and rapidly worse. This is sometimes called dystonic storm or status dystonicus. It can cause dehydration, muscle breakdown, fever, breathing problems, and exhaustion, and requires emergency treatment.

Prevention

Inherited torsion dystonia cannot currently be prevented. Genetic counselling can explain a confirmed genetic result and what it may mean for relatives or future pregnancies, but it cannot predict the exact severity of symptoms.

Many cases with no known cause also cannot be prevented. Avoiding unnecessary exposure to medicines that can cause dystonia may reduce some risk, but these medicines may be important treatments. Never stop or change a prescribed medicine without medical advice.

Early assessment may not prevent dystonia, but it can identify treatable causes and help limit pain, loss of movement, or other complications.

Living with it and what to expect

Torsion dystonia is often long term. Its course is not the same for everyone. Symptoms may remain in one body area, spread gradually, or settle into a stable pattern. Day-to-day intensity can change with sleep, stress, pain, illness, and activity.

When dystonia begins in a leg or arm during childhood, it is more likely to spread over the following years than dystonia that begins in adulthood. Adult-onset neck, eyelid, or task-specific dystonia often remains more localized, although spread is still possible.

Treatment usually manages symptoms rather than removing the condition. Finding the best combination may take time because benefit must be balanced against weakness, sleepiness, thinking problems, or other side effects. Injection appointments, medicine reviews, therapy, and device programming may form part of long-term care.

Many people continue working, studying, exercising, caring for family, and taking part in social activities. Practical adjustments may be needed. Pacing demanding tasks, using adapted equipment, planning rest, and explaining the condition to people you trust can make daily life easier.

Arrange routine follow-up if symptoms are changing, treatment is wearing off sooner than expected, pain is increasing, or daily tasks are becoming harder. New swallowing problems, falls, low mood, or medicine side effects should also be discussed.

When to get medical help

Arrange a routine medical appointment if you develop repeated twisting movements, muscle contractions, or abnormal postures that do not have an obvious short-lived cause. Assessment is also appropriate if a movement began after starting or changing a medicine.

Seek prompt medical advice if:

Call emergency services if you have difficulty breathing, are choking, or cannot swallow saliva. Emergency care is also needed for severe, continuous spasms accompanied by fever, marked sweating, confusion, extreme exhaustion, or dark urine. Sudden abnormal movement with facial drooping, weakness, or trouble speaking could be a stroke and also needs emergency assessment.

- Symptoms worsen quickly over hours or days
- Swallowing becomes difficult or you repeatedly cough while eating
- New weakness, numbness, confusion, or loss of coordination appears
- Pain, falls, or reduced movement are increasing
- A medicine seems to be causing new movements

Common questions

Is torsion dystonia the same as Parkinson's disease?

No. Both are movement conditions, but they are different. Dystonia causes patterned contractions and abnormal postures. Parkinson's disease more often causes slowed movement, stiffness, and a characteristic tremor. Dystonia can occur in someone with Parkinson's, but having dystonia does not mean you have Parkinson's disease.

Does torsion dystonia get worse with age?

It can change over time, but worsening is not inevitable. Early-onset limb dystonia is more likely to spread than adult-onset focal dystonia. The underlying cause also matters. A neurological assessment gives more useful information than age alone.

Can stress cause dystonia?

Stress does not usually cause inherited or neurological dystonia, but it can make movements more noticeable. Tiredness, pain, illness, and focused activity can have a similar effect. Managing stress may reduce flare-ups without removing the underlying condition.

Is torsion dystonia painful?

It can be. Pain is especially common when contractions repeatedly pull the neck, back, or a joint into an uncomfortable position. Treatment directed at the overactive muscles, along with movement therapy and pain management, may help.

Can exercise make dystonia worse?

Exercise does not usually damage the nervous system or make the condition permanently worse. Some movements may trigger temporary contractions, while other activities feel easier. A physical therapist familiar with neurological movement conditions can help you adapt exercise without repeatedly forcing painful positions.

Can torsion dystonia be cured with surgery?

Deep brain stimulation is not considered a cure. It can substantially reduce symptoms in some people, particularly with certain inherited or isolated generalized forms, but the response is not guaranteed. The implanted system also needs long-term monitoring and adjustment.

Sources

- National Institute of Neurological Disorders and Stroke. Dystonia . Neurological condition information, US National Institutes of Health
- Albanese A, Bhatia K, Bressman SB, et al. Phenomenology and classification of dystonia: a consensus update . Movement Disorders . 2013;28(7):863-873
- Balint B, Mencacci NE, Valente EM, et al. Dystonia . Nature Reviews Disease Primers . 2018;4:25
- Simpson DM, Hallett M, Ashman EJ, et al. Practice guideline update summary: Botulinum neurotoxin for the treatment of blepharospasm, cervical dystonia, adult spasticity, and headache . Neurology . 2016;86(19):1818-1826

- Bressman SB, de Leon D, Brin MF. DYT-TOR1A . GeneReviews , University of Washington, Seattle
- National Health Service. Dystonia . NHS health information