CONDITION

# X-linked bulbospinal muscular atrophy

An inherited condition that causes slowly progressive muscle weakness, usually in adult men.

**WHAT IT IS**
Inherited condition causing slowly progressive muscle weakness

**BODY SYSTEM**
Nervous system and muscles

**MAIN SYMPTOMS**
Limb weakness, muscle cramps, twitching, swallowing difficulty

**WHAT CAUSES IT**
Inherited change in the AR gene

**HOW IT IS DIAGNOSED**
Genetic testing for an AR gene repeat expansion

**MAIN TREATMENT**
Supportive therapy, rehabilitation, and symptom management

**CAN IT BE CURED?**
No - the genetic cause cannot be removed

**IS IT AN EMERGENCY?**
Sometimes - sudden breathing difficulty or choking

**LONG-TERM OUTLOOK**
Gradual progression over decades, often with near-normal lifespan

## Overview

X-linked bulbospinal muscular atrophy is an inherited condition affecting the nerves that control voluntary muscles. These nerves are called lower motor neurons. Over time, damage to them causes weakness, muscle shrinkage, cramps, and twitching.

"Bulbar" refers to muscles used for speaking, chewing, and swallowing. "Spinal" refers to motor neurons in the spinal cord that control the arms, legs, and trunk.

The condition usually becomes noticeable in adulthood and progresses slowly over many years. It mainly affects people with one X and one Y chromosome, who are typically male. People with two X chromosomes can carry the genetic change and may occasionally develop mild symptoms.

This condition can resemble amyotrophic lateral sclerosis, or ALS, but they are not the same. X-linked bulbospinal muscular atrophy usually progresses more slowly. It can also cause sensory and hormonal changes that are not typical of ALS.

## Symptoms

Symptoms often begin between early and later adulthood, although the timing differs from person to person. Early symptoms may be subtle, such as frequent cramps, hand tremor, or difficulty climbing stairs.

Muscle weakness usually affects the muscles closest to the centre of the body first. These include the hips, thighs, shoulders, and upper arms. Weakness may later spread to the hands, lower legs, face, and throat.

Common symptoms include:

Some people have reduced sensation, particularly in the feet. This happens because sensory nerves can also be affected, although numbness may be mild.

The altered gene affects how the body responds to androgens, which are hormones such as testosterone. Related features can include breast tissue enlargement, reduced fertility, smaller testicles, erectile problems, or reduced body hair. Diabetes, abnormal blood fats, and fatty liver are also reported more often in people with this condition.

Thinking, memory, and bladder control are generally not major areas of difficulty.

- Muscle cramps, especially after activity
- Muscle twitching, also called fasciculations
- Tremor in the hands
- Difficulty running, climbing stairs, or rising from a chair
- Frequent trips or falls
- Weakness in the shoulders, arms, hips, or legs
- Muscle shrinking, known as muscle atrophy
- Slurred or quieter speech
- Difficulty chewing or swallowing
- Coughing during meals or drinking
- Weakness of the facial or tongue muscles

## Causes and risk factors

X-linked bulbospinal muscular atrophy is caused by an expanded section of DNA in the AR gene. This gene provides instructions for making the androgen receptor, a protein that allows cells to respond to hormones such as testosterone.

The expansion causes the androgen receptor protein to fold and function abnormally. The altered protein gradually damages motor neurons and muscle cells. Androgens appear to contribute to this process, which helps explain why substantial muscle weakness occurs mainly in people with one X and one Y chromosome.

The condition follows an X-linked inheritance pattern. A person with one altered X chromosome and one Y chromosome will usually develop symptoms, although the age and severity cannot be predicted precisely.

A person with two X chromosomes and one altered copy is called a carrier. The second working copy usually offers protection. Some carriers develop cramps, twitching, tremor, or mild weakness, especially later in life.

A family history may not be obvious. Older relatives may have had mild or unrecognised symptoms, and the altered gene may have passed through several carriers.

## Diagnosis

A blood test that looks for the repeat expansion in the AR gene confirms the diagnosis. Genetic counselling can help you understand what the result means for you and your biological relatives.

Before genetic testing, an assessment may include questions about weakness, swallowing, cramps, hormonal symptoms, and family history. A physical examination can look for muscle loss, twitching, reduced reflexes, tremor, and weakness in particular muscle groups.

Other tests can support the diagnosis or check for complications:

These findings are not enough to confirm the condition without genetic testing. They may also help distinguish it from ALS, other motor neuron conditions, muscle disorders, or treatable causes of weakness.

Testing relatives who do not have symptoms requires careful discussion. A result can affect emotional wellbeing, family planning, insurance, and employment rights, depending on local laws.

- Electromyography , or EMG, measures electrical activity in muscles and can show loss of nerve supply
- Nerve conduction studies assess how well motor and sensory nerves carry signals
- Creatine kinase blood testing measures a muscle enzyme that may be raised when muscle is damaged
- Swallowing, breathing, liver, glucose, and cholesterol tests assess specific effects of the condition

## Treatment and management

There is currently no cure or established treatment that removes the genetic cause. Care focuses on maintaining movement and independence, managing symptoms, and identifying swallowing or breathing problems early.

You may see a neurologist alongside rehabilitation, speech and language, nutrition, respiratory, hormone, or genetic specialists. The mix of support usually changes as your needs change.

Physical therapy can help preserve flexibility, balance, and safe movement. Moderate activity may support general fitness, but repeated exercise to exhaustion can worsen fatigue and may increase the risk of injury. An assessment can help identify a suitable level of activity.

Occupational therapy can suggest ways to make daily tasks safer and less tiring. Braces, walking aids, handrails, adapted seating, or a wheelchair may improve independence rather than signal a failure to stay active.

Speech and language therapy can assess speech and swallowing. Management may include safer swallowing techniques, changes to food texture, communication aids, or strategies for conserving the voice. If eating becomes unsafe or does not provide enough nutrition, tube feeding may be discussed.

Breathing is usually affected later or less severely than in some other motor neuron conditions, but it still needs attention. Breathing tests and sleep studies may be used if there is shortness of breath, morning headache, disturbed sleep, or daytime sleepiness. Some people benefit from assisted coughing or non-invasive ventilation, which supports breathing through a mask.

Medicines may be offered for particular problems such as cramps, pain, tremor, or excess saliva. There is no single medicine that is suitable for everyone.

Treatments designed to reduce androgen activity have been studied, but they are not routine curative therapy. Testosterone or anabolic steroid use may theoretically worsen the underlying process. Anyone with the condition or its genetic change should discuss hormone treatment

with a specialist before starting or changing it.

## Complications

Weakness can increase the risk of falls, fractures, and loss of mobility. Reduced activity may also contribute to stiffness, weight changes, or lower physical fitness.

Swallowing weakness can allow food, drink, or saliva to enter the airway. This is called aspiration. It can lead to choking, weight loss, dehydration, or aspiration pneumonia.

Weak cough and respiratory muscle weakness can make it harder to clear mucus during a chest infection. Breathing problems during sleep may develop before breathlessness is obvious during the day.

Hormonal and metabolic effects can include reduced fertility, sexual difficulties, breast tissue enlargement, diabetes, abnormal cholesterol or triglyceride levels, and fatty liver. These can be monitored and treated separately.

The condition can also affect work, driving, relationships, and emotional wellbeing. Practical support and counselling can be useful even when physical symptoms are changing slowly.

## Prevention

X-linked bulbospinal muscular atrophy cannot be prevented in someone who has inherited the expanded AR gene.

Genetic counselling can explain inheritance and reproductive choices. These may include testing during pregnancy or testing embryos created through in vitro fertilisation. Whether to use genetic testing is a personal decision.

If a biological relative receives a confirmed diagnosis, other adult relatives may choose to discuss carrier or predictive testing. Testing children for an adult-onset condition is generally approached cautiously unless there is a clear medical reason to know during childhood.

## Living with it and what to expect

The condition usually progresses slowly over decades. Weakness may begin in one area and gradually involve more muscles. Abilities often change in stages rather than from week to week.

Many people continue walking and working for years after symptoms begin. Some eventually

need a stick, walker, wheelchair, or help with personal care. It is not possible to predict from a genetic result alone when these changes will happen.

Speech and swallowing problems may become more noticeable with time. Regular review can identify changes before they cause major weight loss or repeated chest infections. Breathing should also be checked if symptoms suggest weak respiratory muscles or disturbed sleep.

Life expectancy is often close to normal, but serious swallowing or breathing complications can shorten life. Prompt assessment of choking, recurrent chest infections, unexplained weight loss, or breathing symptoms can reduce some of these risks.

Planning ahead can make changes easier to manage. This may include reviewing home access, mobility equipment, work adjustments, communication options, and future feeding or breathing support. Plans can be updated as your priorities change.

## When to get medical help

Arrange a routine appointment if you develop gradual unexplained weakness, repeated muscle cramps, frequent falls, persistent twitching, or changes in speech or swallowing. These symptoms have many possible causes and do not by themselves mean you have this condition.

If you have already been diagnosed, contact your usual care team promptly about:

Seek urgent medical care for a suspected chest infection with breathing difficulty, especially after choking or swallowing problems.

Call emergency services for sudden severe breathing difficulty, blue or grey lips, inability to speak because of breathlessness, or choking that does not clear. These are emergencies regardless of their cause.

- Increasing coughing or choking during meals
- Unplanned weight loss or difficulty maintaining hydration
- More frequent falls or a sudden decline in mobility
- Morning headaches, poor sleep, or unusual daytime sleepiness
- Shortness of breath with less activity than usual
- Repeated chest infections

## Common questions

Is X-linked bulbospinal muscular atrophy the same as ALS?

No. Both conditions affect motor neurons and can cause weakness and twitching, but they have different causes and typical courses. X-linked bulbospinal muscular atrophy is confirmed by an AR gene test, usually progresses more slowly, and may cause sensory and hormonal changes.

Can women develop symptoms?

People with two X chromosomes can carry the genetic change. Most carriers have no symptoms or only mild problems, but some develop cramps, tremor, twitching, or weakness later in life. A carrier with new symptoms should have an appropriate medical assessment rather than assuming the gene is the cause.

Will I eventually need a wheelchair?

Some people do, while others remain able to walk with or without support for many years. The condition usually progresses slowly, and genetic testing cannot accurately predict an individual timeline. Mobility equipment can reduce falls, fatigue, and pain while supporting independence.

Can I pass it on to my children?

A father with the condition passes his X chromosome to all biological daughters and his Y chromosome to all biological sons. His daughters therefore inherit the altered gene, while his sons do not inherit it from him.

A carrier with two X chromosomes has a 50% chance in each pregnancy of passing on the altered X chromosome. Genetic counselling can explain this in relation to your family.

Is exercise safe?

Moderate, low-impact activity is often appropriate and may help maintain fitness and mobility. Exercise should not repeatedly leave you exhausted, very sore, or weaker for a prolonged period. A physical therapist familiar with neuromuscular conditions can help adjust activity to your abilities.

Does it affect intelligence or memory?

Major changes in intelligence or memory are not typical features. Physical fatigue, sleep-related breathing problems, stress, or low mood can still affect concentration and should be discussed during follow-up.

## Sources

- Fischbeck KH. Spinal and Bulbar Muscular Atrophy . GeneReviews®, University of Washington, Seattle. Updated 2017

- National Institute of Neurological Disorders and Stroke. Kennedy's Disease . Neurological condition information
- MedlinePlus Genetics. Spinal and bulbar muscular atrophy . U.S. National Library of Medicine
- Grunseich C, Fischbeck KH. "Spinal and Bulbar Muscular Atrophy." Neurologic Clinics . 2015;33(4):847-854
- Rhodes LE, Freeman BK, Auh S, et al. "Clinical features of spinal and bulbar muscular atrophy." Brain . 2009;132(12):3242-3251