DIET & WEIGHT

Why Vitamin D Can Run Low

A low result can begin with sunlight, food, absorption, medicines, or the organs that make vitamin D usable.

The result is only the beginning

You open your test results and there it is: vitamin D, low. The number may look like a simple verdict. The story behind it rarely is. Your level reflects several steps-vitamin D reaching your body, entering your blood and being converted into a form you can use. A problem anywhere along that route can leave you short.

That is the first reason low vitamin D can seem to turn up everywhere. There is no single doorway. Your skin can form it after sunlight exposure; food and supplements can provide it; your digestive system must absorb it; and your liver and kidneys help convert it into its active form. Several ordinary biological routes all have to work.

Deficiency can result from too little vitamin D in the diet, too little sunlight, poor absorption, certain medicines, or difficulty in the liver or kidneys. Those explanations are not interchangeable. Someone who absorbs nutrients poorly is not in the same position as someone whose main limitation is sunlight.

The supplied evidence does not establish how many people are deficient, so the popular claim that "everyone is low" goes further than these sources allow. What the evidence does show is why low levels can appear across very different lives. The same laboratory finding can be the endpoint of several different stories.

Sunlight is not a simple solution

Vitamin D stands apart because your body can form it naturally after your skin is exposed to sunlight. That makes "get more sun" sound like the obvious answer. It is not a complete one. Excessive sun exposure can contribute to skin aging and skin cancer, which is why many people look to other sources.

Skin also does not produce vitamin D equally in everyone. Older adults make it less efficiently during sunlight exposure than younger people. Dark skin has less ability to produce vitamin D from the sun. These are biological differences, not proof that someone has made a poor lifestyle choice.

Age can create a second bottleneck. Along with reduced production in the skin, older kidneys are less able to convert vitamin D into its active form. One person may therefore face limitations at both the beginning and the end of the pathway.

This is where the easy story breaks down. Sunlight contributes to vitamin D, but it does not erase differences in skin, age, absorption, medicines, liver function or kidney function. A low level cannot automatically tell you which link in the chain is responsible.

The same laboratory finding can be the endpoint of several different stories.

Food has limits too

Perhaps you turn to your kitchen next. Vitamin D occurs naturally in some foods, including fatty fish such as salmon, tuna and mackerel, as well as beef liver, cheese, mushrooms and egg yolks. Fortified foods can add another route; milk, breakfast cereals, orange juice, yogurt and soy drinks may contain added vitamin D.

The important word is "may." Fortification is something you can check on a food label rather than assume. And even when vitamin D is present, eating it is not the final step. Your body still has to absorb it.

Crohn's disease, ulcerative colitis and celiac disease can make it harder to absorb nutrients from food. Gastric bypass surgery can also make vitamin D absorption more difficult because the operation bypasses part of the small intestine, where vitamin D is absorbed. In these situations, simply choosing a vitamin D-containing food may not address the whole problem.

Obesity introduces another route to a low blood level. Body fat binds some vitamin D and prevents it from entering the blood. Again, the visible result is the same, but the underlying process is different. A food list alone cannot account for that difference.

Your body must finish the job

Vitamin D arriving in your body is not the same as vitamin D being ready for use. Your liver and kidneys must convert it into an active form. Chronic liver or kidney disease can interfere with that conversion, leaving you unable to use vitamin D in the usual way.

Medicines can also affect vitamin D levels. Certain cholesterol-lowering, anti-seizure, steroid and weight-loss medicines are possible contributors because they can interfere with conversion or absorption. That does not mean you should stop a medicine. It means the medicine belongs in the conversation when a low result is being interpreted.

These details matter because vitamin D helps your body absorb calcium, one of the main building blocks of bone. It also plays roles in the nervous, muscle and immune systems. Deficiency can reduce bone density and contribute to osteoporosis and fractures.

Severe deficiency carries clearer consequences. In children, it can cause rickets, a rare condition in which bones become soft and bend. In adults, severe deficiency can cause osteomalacia, with weak bones, bone pain and muscle weakness. At that point-or if you have a condition or medicine that raises your risk-this is no longer just a lifestyle puzzle. It belongs with your health care provider.

More is not automatically safer

Once you hear that your level is low, it is tempting to treat vitamin D as a harmless substance that can only help. The evidence does not support that assumption. Supplements are widely available, including in multivitamins, pills and liquid products, but too much vitamin D can be harmful.

Vitamin D toxicity can cause nausea, vomiting, poor appetite, constipation, weakness and weight loss. Very high levels can damage the kidneys and raise calcium in the blood. High blood calcium can lead to confusion, kidney failure and an irregular heartbeat.

Most vitamin D toxicity occurs when someone overuses supplements. Sun exposure does not cause the same problem because the skin limits how much vitamin D it makes. That distinction is another reason not to improvise: the source of vitamin D changes the risk.

For the next week, keep the task small. Check labels on foods and supplements you already use, write down relevant medicines and health conditions, and bring that information to your provider if your result is low or you think you are at risk. A blood test can measure vitamin D in your body; the right response depends on why yours is low.

Where this comes from

- Vitamin D Deficiency

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/vitaminddeficiency.html
- Intermittent fasting strategies and their effects on body weight and other cardiometabolic risk factors: systematic review and network meta-analysis of randomised clinical trials.

  BMJ (Clinical research ed.), 2025 · https://pubmed.ncbi.nlm.nih.gov/40533200/

- Diet, Food, and Nutritional Exposures and Inflammatory Bowel Disease or Progression of Disease: an Umbrella Review.

  Advances in nutrition (Bethesda, Md.), 2024 · https://pubmed.ncbi.nlm.nih.gov/38599319/
- Dietary fibre and whole grains in diabetes management: Systematic review and meta-analyses.

  PLoS medicine, 2020 · https://pubmed.ncbi.nlm.nih.gov/32142510/
- Effect of low glycaemic index or load dietary patterns on glycaemic control and cardiometabolic risk factors in diabetes: systematic review and meta-analysis of randomised controlled trials.

  BMJ (Clinical research ed.), 2021 · https://pubmed.ncbi.nlm.nih.gov/34348965/
- Mediterranean diet in the management and prevention of obesity.

  Experimental gerontology, 2023 · https://pubmed.ncbi.nlm.nih.gov/36792040/