SLEEP & STRESS

The Sleep Apnea Signs You Might Miss

The loudest clues happen at night, but daytime sleepiness, headaches and concentration problems can complete the picture.

The clue you may never hear

Someone tells you that you stopped breathing in your sleep. Perhaps they heard silence after a stretch of loud snoring, followed by a gasp, snort or choking sound. You feel embarrassed, doubtful or both. You were asleep, after all. Yet this second-hand account may be the most important clue you receive.

Sleep apnea repeatedly stops or reduces breathing during sleep. In obstructive sleep apnea (OSA), the most common type, the upper airway becomes blocked and airflow falls or stops. Breathing may restart with a snort or choking sound. Central sleep apnea is different: the brain does not send the signals needed to breathe.

The nighttime pattern to listen for is not simply noise. It is breathing that starts and stops, frequent loud snoring and gasping for air. Pauses can last from seconds to minutes and may happen 30 or more times in an hour. Because you are asleep when this occurs, another person may notice it first.

Snoring is a clue, not a verdict. People with sleep apnea often snore loudly, but not everyone who snores has the condition. The more concerning report is repeated snoring interrupted by apparent pauses, gasps or choking sounds. That pattern deserves a conversation with a healthcare provider rather than another joke about the noise.

Daytime can tell the story

You may wake without any memory of struggling to breathe. Instead, the evidence follows you into the day: sleepiness, tiredness or fatigue that does not fit the night you thought you had. Interrupted sleep can leave you drowsy even if you spent plenty of time in bed.

Poor-quality sleep can also show up in your thinking. You may struggle to learn, focus or react. Concentrating, making decisions, remembering things and controlling your behavior can become harder when sleep apnea is untreated. These problems are easy to blame on stress or a demanding week, which is one reason the nighttime clues matter.

A dry mouth, reduced libido or sexual dysfunction can also occur. Other possible signs include

waking frequently during the night to urinate and insomnia. Headache, fatigue and insomnia are reported more commonly in women, while frequent loud snoring is more common in men. That difference can make a quieter pattern easier to overlook.

None of these daytime experiences proves that sleep apnea is responsible. Headache in particular is not a simple shortcut to an answer: a 2020 systematic review found that the relationship between migraine and sleep disorders is complex and not completely understood. The useful question is whether daytime problems sit beside reports of disturbed breathing at night.

Snoring is a clue, not a verdict.

Risk changes how loudly clues speak

Sleep apnea is common, but estimates vary sharply according to the population studied and the threshold used. A 2018 systematic review of 24 studies found OSA prevalence estimates ranging from 9% to 38% at one commonly used threshold. The wide range reflected substantial differences in study methods, so it should not be read as one tidy figure for everyone.

That review found OSA was more prevalent with advancing age, male sex and higher body mass index. General guidance also identifies family history and small airways as risk factors. Obesity, large tonsils and changes in hormone levels can narrow the airway and raise the likelihood of obstruction during sleep.

Risk is context, not confirmation. Being male, older or living in a larger body does not establish that your nighttime breathing stops. Equally, someone outside those groups should not dismiss witnessed pauses, gasping or persistent daytime sleepiness. The signs still need proper assessment.

Children can have sleep apnea too, including children with enlarged tonsils or adenoids. Their daytime picture may look different: overactivity, trouble paying attention at school, bedwetting and worsening asthma are among the reported signs. A child with these concerns and disturbed nighttime breathing needs assessment by a healthcare provider.

A watch cannot settle it

If the pattern sounds familiar, this is where guessing should stop. A healthcare provider can consider your symptoms, medical and family histories and a physical examination. A sleep study may then be needed to find out whether sleep apnea is present. The concern becomes

medical rather than merely inconvenient when breathing repeatedly stops or restarts, or daytime sleepiness affects safe driving, work or ordinary functioning.

Wearable devices may appear to offer a quicker answer. A 2024 systematic review and meta-analysis of 38 studies found that wearable artificial intelligence (AI) showed potential for identifying and classifying sleep apnea. But its performance varied by factors including the algorithm, the data collected, the type of apnea and where the device was worn.

The review judged current wearable AI performance suboptimal for routine clinical use and recommended using it alongside traditional assessment until stronger evidence supports its reliability. In practical terms, a wearable alert can be something to mention. It should not overrule witnessed breathing pauses, and it should not be treated as a confirmed result by itself.

Getting assessed matters because untreated sleep apnea is not just noisy sleep. It can raise the risk of car crashes and work-related accidents, and it is associated with increased risks of stroke, heart attack and other serious problems. Treatments can include lifestyle changes, positive airway pressure devices, custom-fitted oral devices, orofacial therapy and surgery, depending on the person and the findings.

Make a one-page sleep note

This week, use one sheet of paper and a pen to gather the clues that are otherwise scattered between night and day. Write down whether anyone has noticed loud snoring, breathing pauses, gasping, snorting or choking. Add the daytime problems you have noticed, such as sleepiness, fatigue, poor concentration, dry mouth, headache or waking often to urinate.

Keep the note factual. "Breathing seemed to stop, then restarted with a gasp" is more useful than "slept badly." If nobody shares your sleeping space, record what you can observe yourself: daytime sleepiness, tiredness, insomnia, dry mouth, headaches and nighttime waking. Do not invent certainty where there is none.

Take that sheet to a healthcare provider and say plainly what brought you in. Ask whether the combination of nighttime and daytime signs suggests that you need a sleep study. Mention any wearable alert as one piece of information, not the final answer. If sleepiness is interfering with driving, work or your ability to react, say that directly.

The step for Monday is small: one page, one pen and one honest account of what is happening. You do not need to decide whether you have sleep apnea. You need to make the pattern visible enough for the right assessment to begin.

Where this comes from

- Sleep Apnea

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/sleepapnea.html
- Sleep Apnea - What Is Sleep Apnea? | NHLBI, NIH

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/health/sleep-apnea
- Sleep Apnea - Symptoms | NHLBI, NIH

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/health/sleep-apnea/symptoms
- Sleep Apnea - Treatment | NHLBI, NIH

  National Heart, Lung, and Blood Institute · https://www.nhlbi.nih.gov/health/sleep-apnea/treatment
- Prevalence of obstructive sleep apnea in the general population: A systematic review.

  Sleep medicine reviews, 2018 · https://pubmed.ncbi.nlm.nih.gov/27568340/
- Migraine and sleep disorders: a systematic review.

  The journal of headache and pain, 2020 · https://pubmed.ncbi.nlm.nih.gov/33109076/
- Orofacial Myofunctional Therapy for Obstructive Sleep Apnea: A Systematic Review and Meta-Analysis.

  The Laryngoscope, 2023 · https://pubmed.ncbi.nlm.nih.gov/37606313/
- Detection of Sleep Apnea Using Wearable AI: Systematic Review and Meta-Analysis.

  Journal of medical Internet research, 2024 · https://pubmed.ncbi.nlm.nih.gov/39255014/
- Exercise and sleep: a systematic review of previous meta-analyses.

  Journal of evidence-based medicine, 2017 · https://pubmed.ncbi.nlm.nih.gov/28276627/