DIET & WEIGHT

Which Foods Are Really Making Your Reflux Worse?

The usual suspects matter, but meal size, timing and your own pattern may tell you more than a universal banned-food list.

The suspect may be dinner itself

You finish dinner, settle onto the sofa and soon feel that familiar burn rising through your chest or throat. Perhaps you blame the tomato sauce. But the more useful question is not simply what was on your plate. It is also how much you ate and what you did next.

Gastroesophageal reflux (GER) happens when stomach contents move back into the oesophagus, the tube connecting your mouth and stomach. The muscle at the bottom of the oesophagus normally helps stop that return journey. Reflux can occur when it becomes weak or relaxes when it should not.

An occasional episode is common. Repeated symptoms or damage over time may mean gastroesophageal reflux disease (GERD), a chronic condition rather than a single disagreeable meal. Heartburn and the taste of acid or food at the back of your mouth are common, but reflux can also be associated with a dry cough, hoarseness or trouble swallowing.

That distinction matters because food is only one part of the picture. Being overweight or having obesity, pregnancy, smoking or exposure to secondhand smoke, and a hiatal hernia are all associated with a greater likelihood of GERD. Medicines can make it worse, too. A perfect menu cannot necessarily solve a problem that is not mainly about the menu.

The usual suspects are not convictions

The commonly linked foods and drinks form a familiar lineup: acidic foods such as citrus fruits and tomatoes, alcohol, chocolate, coffee and other caffeine sources, high-fat foods, mint and spicy foods. General guidance often suggests reducing or avoiding a suspected item to see whether symptoms improve.

But "commonly linked" is not the same as "guaranteed to cause reflux in everyone." The dietary advice is explicitly personal: some people find that particular foods or drinks trigger symptoms or make them worse. That makes a universal blacklist a blunt instrument. It may

remove foods that never troubled you while leaving your actual pattern undiscovered.

It is also easy to turn one meal into a misleading verdict. A spicy, high-fat dinner accompanied by alcohol and coffee contains several usual suspects at once. If symptoms follow, you cannot tell which item mattered-or whether the large meal and the decision to lie down shortly afterwards were more important.

The honest conclusion is less dramatic than many reflux diets promise. The available guidance identifies foods worth investigating, not foods that everyone with reflux must permanently surrender. Your job is not to eat the blandest possible diet. It is to notice whether a repeatable connection exists between a particular food or drink and your symptoms.

The available guidance identifies foods worth investigating, not foods that everyone with reflux must permanently surrender.

The clock can matter as much as the plate

Reflux after dinner may look like evidence against dinner's ingredients when the timing is doing much of the work. Eating a large meal and lying down too soon afterwards can allow stomach contents to come back up and irritate the oesophagus.

For symptoms that appear at night or while you are lying down, guidance recommends eating at least three hours before you lie down or go to bed. Eating smaller meals is another commonly advised change. These are not promises of relief, but they let you test something more precise than "food is the problem."

This means the same dish may not tell the same story in every setting. A meal eaten earlier and in a smaller amount differs from a large meal followed quickly by bed. If you change the ingredients, portion and timing simultaneously, you may feel better without learning which change mattered.

Other measures may help some people manage symptoms: avoiding alcohol, wearing loose-fitting clothes, sleeping on the left side or raising the head of the bed, quitting smoking, and maintaining a healthy weight. If you are overweight or have obesity, a doctor may suggest weight loss and recommend a healthy eating plan.

Food stops being the whole question

If reflux happens occasionally, looking for a food or timing pattern may be reasonable. If symptoms occur two or more times a week, cause persistent discomfort or suggest damage to the oesophagus, the issue may be GERD. That deserves a medical conversation rather than an

ever-longer list of forbidden foods.

Untreated GERD can lead to inflammation, scarring or changes in the tissue lining of the oesophagus called Barrett's oesophagus. It can also worsen asthma or contribute to a long-lasting cough or hoarse voice. Trouble swallowing is another reason not to keep treating the problem as a simple disagreement with chilli or coffee.

Most cases are assessed first through your symptoms and medical history. If symptoms do not improve with lifestyle changes and medicines, or if another health problem or a complication is possible, a doctor may recommend tests. These can include an upper gastrointestinal endoscopy or oesophageal pH monitoring, which detects stomach acid in the oesophagus and relates reflux to diet, sleep and symptoms.

Chest pain needs particular care. Get medical help immediately if it occurs with shortness of breath or pain in your jaw or arm, because those can be signs of a heart problem. Do not assume that a burning feeling after food must be reflux.

Make one week tell you something

For the next week, use a plain notebook and a pen to record what you ate and drank, whether the meal was large or small, whether you lay down soon afterwards, and what symptoms followed. Keep the entries simple enough that you will actually make them.

Start by observing rather than banning the entire usual-suspect list. Look for repeated links involving citrus, tomatoes, alcohol, chocolate, coffee or other caffeine sources, high-fat foods, mint or spicy foods. A single uncomfortable evening is a clue; a recurring pattern is more useful.

If one item repeatedly appears before symptoms, reduce or avoid that item and see whether the pattern changes. Try not to overhaul every part of the meal at once. Separating food from meal size and bedtime timing gives you a clearer account to bring to a doctor if symptoms continue.

A notebook cannot diagnose GERD, and it should not delay care for frequent symptoms, trouble swallowing or concerning chest pain. What it can do is replace a borrowed list of "bad foods" with your own record. This Monday, put the notebook and pen where you eat: the first useful reflux diet is the one that shows you what actually happens.

Where this comes from

- GERD

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/gerd.html
- Reflux in Infants

  MedlinePlus, U.S. National Library of Medicine · https://medlineplus.gov/refluxininfants.html
- Acid Reflux (GER & GERD) in Adults

  National Institute of Diabetes and Digestive and Kidney Diseases · https://www.niddk.nih.gov/health-information/digestive-diseases/acid-reflux-ger-gerd-adults
- Diagnosis of GER & GERD

  National Institute of Diabetes and Digestive and Kidney Diseases · https://www.niddk.nih.gov/health-information/digestive-diseases/acid-reflux-ger-gerd-adults/diagnosis
- Eating, Diet, & Nutrition for GER & GERD

  National Institute of Diabetes and Digestive and Kidney Diseases · https://www.niddk.nih.gov/health-information/digestive-diseases/acid-reflux-ger-gerd-adults/eating-diet-nutrition
- The role of dietary supplements in inflammatory bowel disease: a systematic review.

  European journal of gastroenterology & hepatology, 2017 · https://pubmed.ncbi.nlm.nih.gov/27769076/
- Laparoscopic sleeve gastrectomy and gastroesophageal reflux disease: a systematic review and meta-analysis.

  American journal of surgery, 2016 · https://pubmed.ncbi.nlm.nih.gov/26341463/