SYMPTOM

Abdominal sounds

Abdominal sounds are noises made as food, fluid, and gas move through your intestines.

WHAT IT FEELS LIKE
Gurgling, rumbling, growling, or popping inside your belly

WHERE IT IS FELT
Across the belly, around the stomach and intestines

MOST COMMON CAUSES
Normal digestion, hunger, swallowed air

WHEN IT IS URGENT
Severe pain with vomiting, swelling, or inability to pass gas

USUALLY LASTS
Minutes to hours, based on meals and bowel activity

SEE A DOCTOR IF
Persistent changes with pain, bowel changes, or weight loss

Overview

Abdominal sounds are noises made when the muscles of your digestive tract move food, liquid, and gas through your intestines. This movement is called peristalsis . Although often called "stomach growling," many sounds come from the intestines.

Most abdominal sounds are a normal part of digestion. They may become louder after eating, when you are hungry, or when there is more gas or liquid in your intestines.

Sounds alone rarely show whether something is wrong. Their meaning depends on other symptoms, such as pain, vomiting, diarrhea, constipation, swelling, fever, or weight loss.

What it may feel or look like

Abdominal sounds may include:

They may seem to move across your belly. Some are loud enough for others to hear, while others feel like movement with little sound.

Noises may happen in short bursts or continue on and off for several hours. They can occur with hunger, after a meal, or before a bowel movement. Loudness does not necessarily reflect

how serious the cause is.

Doctors may describe sounds as active, reduced, or absent when listening with a stethoscope. This is difficult to judge at home, and a quiet belly alone is not a reliable sign of illness.

- Rumbling or growling
- Gurgling or bubbling
- Squeaking or high-pitched noises
- Popping or churning sensations

Common causes

Normal digestion

Your intestines contract to move their contents forward. Gas and liquid can make these contractions easier to hear, especially after eating.

Hunger

An empty stomach and small intestine still contract in cycles, creating growling noises that usually settle after eating.

Swallowed air and intestinal gas

Eating quickly, talking while eating, chewing gum, using a straw, or drinking fizzy beverages can increase swallowed air. Bacteria in the large intestine also produce gas when they break down certain carbohydrates.

Diarrhea or a short-term stomach infection

Extra liquid and faster intestinal movement can cause frequent gurgling, often with loose stools, cramps, nausea, or vomiting.

Food-related symptoms

Some people notice more gas and noise after certain foods. This may happen with lactose intolerance or other poorly absorbed carbohydrates. Sounds alone do not confirm an intolerance.

Less common causes

Persistent sounds with discomfort or bowel changes can occur with irritable bowel syndrome. Medicines that speed up or slow down the intestines may also affect sounds.

Rarely, unusually active or reduced sounds occur with an intestinal blockage or ileus , when the intestines temporarily stop moving normally. These conditions usually also cause pain,

vomiting, swelling, or inability to pass stool or gas.

When to seek medical care

Abdominal sounds without pain or other symptoms usually do not require medical care.

Seek urgent, same-day medical advice if the sounds occur with:

These symptoms can occur when intestinal contents are not moving through normally.

Call emergency services or go to an emergency department for sudden or severe abdominal pain, fainting, confusion, a hard or very tender abdomen, vomiting blood, or black or bloody stools.

Arrange a routine appointment if persistent changes occur with recurring pain, ongoing diarrhea or constipation, unexplained weight loss, reduced appetite, or symptoms that regularly disturb sleep. Also seek advice if symptoms began after starting a medicine. Do not stop prescribed medicine without consulting a healthcare professional.

- Repeated vomiting
- Increasing abdominal swelling
- Significant or worsening abdominal pain
- Inability to pass stool or gas
- Signs of dehydration, such as very little urine or marked dizziness

How the cause is found

A healthcare professional will ask when the sounds started and whether they relate to meals, foods, bowel movements, or medicines. They may also ask about pain, vomiting, gas, stool changes, previous abdominal surgery, and weight loss.

During an examination, they may feel your abdomen and listen with a stethoscope. Bowel sounds must be interpreted alongside your symptoms and examination; sound patterns alone cannot confirm a diagnosis.

Testing is often unnecessary when sounds are occasional and there are no concerning symptoms. Depending on other symptoms, tests may include blood or stool tests and imaging such as an X-ray, ultrasound, or CT scan. Persistent food-related symptoms may lead to testing for an intolerance or digestive condition.

Treatment and self-care

Normal abdominal sounds do not need treatment. If they are bothersome, try:

Do not remove many foods from your diet based only on abdominal sounds. Restrictive diets can make it harder to get enough nutrients. A clinician or dietitian can help if a food repeatedly causes pain, diarrhea, or bloating.

If you have diarrhea, replace lost fluids. Seek medical advice if you cannot keep fluids down or develop signs of dehydration.

Treatment for persistent sounds depends on the cause and may include managing constipation or diarrhea, reviewing medicines, treating an infection, or addressing a confirmed intolerance. A suspected intestinal blockage needs urgent medical assessment, not home treatment.

- Eating slowly and chewing food well
- Having smaller, regular meals rather than long gaps
- Reducing fizzy drinks if they increase gas
- Limiting chewing gum or straws if they worsen symptoms
- Keeping a short food and symptom record if episodes recur

What to expect

Sounds linked to hunger, meals, or swallowed air often come and go over minutes or hours. They may settle after eating, passing gas, or having a bowel movement.

Noisy digestion can recur without causing harm. Rather than monitoring every sound, watch for pain, vomiting, swelling, altered bowel habits, or a decline in your general health.

When another digestive problem is responsible, the noises usually improve as that problem settles or is treated.

Common questions

Why does my belly make noises after eating?

Eating activates the muscles that move food through your digestive tract. Gas and liquid moving with food can make this activity audible.

Why does my stomach growl when I am hungry?

The stomach and small intestine continue contracting between meals. With less solid food present, movement of air and fluid may sound louder.

Can loud abdominal sounds mean a blockage?

A blockage cannot be identified from sound alone. Seek urgent care if noises occur with worsening pain, repeated vomiting, marked swelling, or inability to pass stool or gas.

What does a very quiet abdomen mean?

A quiet abdomen can be normal, including during sleep. It may also occur when intestinal movement slows, but this cannot be assessed reliably without an examination. Quietness with severe pain, vomiting, or swelling needs urgent assessment.

Can stress make abdominal sounds louder?

Stress can affect intestinal movement and make you more aware of bodily sensations. However, do not assume new or persistent symptoms are due to stress, especially if other digestive symptoms are present.

Sources

- MedlinePlus Medical Encyclopedia: Abdominal sounds , US National Library of Medicine
- National Institute of Diabetes and Digestive and Kidney Diseases: Symptoms and causes of gas in the digestive tract
- National Institute of Diabetes and Digestive and Kidney Diseases: Symptoms and causes of intestinal pseudo-obstruction
- Jackson PG, Raiji MT. Evaluation and Management of Intestinal Obstruction , American Family Physician
- American College of Radiology Appropriateness Criteria: Suspected Small-Bowel Obstruction , American College of Radiology