SYMPTOM

Visceromegaly

Visceromegaly is enlargement of one or more internal organs, usually in the abdomen.

WHAT IT FEELS LIKE
Fullness, pressure, or a lump under the ribs

WHERE IT IS FELT
Upper abdomen, often beneath either rib cage

MOST COMMON CAUSES
Fatty liver disease, infections, blood-flow congestion

WHEN IT IS URGENT
Sudden severe belly pain, fainting, or pain after injury

USUALLY LASTS
Until the underlying cause improves or is treated

SEE A DOCTOR IF
Fullness, swelling, or an unexplained lump persists

Overview

Visceromegaly means that one or more internal organs are larger than expected. The term usually refers to organs in the abdomen, such as the liver, spleen, or kidneys. It describes a physical finding, not a disease or a diagnosis.

Doctors often use a more specific name. An enlarged liver is called hepatomegaly , while an enlarged spleen is called splenomegaly . Enlargement of both is called hepatosplenomegaly .

You may notice pressure or swelling, but many enlarged organs cause no symptoms. The change may instead be found during an examination or on an ultrasound, CT scan, or MRI performed for another reason.

Organ size alone does not show how serious the cause is. Some causes are temporary or treatable. Others need longer-term monitoring and care.

What it may feel or look like

Visceromegaly can cause a sense of fullness, pressure, or heaviness in the upper abdomen. You might feel full after eating a small amount because an enlarged organ is pressing on the

stomach.

An enlarged liver may cause discomfort beneath the right ribs. An enlarged spleen more often causes discomfort beneath the left ribs or pain that spreads toward the left shoulder. Kidney enlargement may cause pain or fullness in the side or back.

Other possible changes include:

These symptoms can occur for many reasons other than organ enlargement. A lump that you can feel should be examined rather than identified at home.

- A firm area or lump beneath the ribs
- Visible swelling or an increasing waist size
- Reduced appetite or feeling full quickly
- Tiredness, fever, or weight change related to the underlying cause
- Yellow skin or eyes when liver or bile flow is affected

Common causes

Common causes depend on which organ is enlarged.

Fat buildup in the liver, often called fatty liver disease, is a common reason for liver enlargement. Liver inflammation from alcohol, medicines, or viral infections can also increase its size.

Infections may enlarge the liver, spleen, or both. Examples include infectious mononucleosis, viral hepatitis, and some bacterial or parasitic infections. The likely infection depends on your symptoms, travel, exposures, and health history.

Blood-flow problems can make an organ congested with blood. Heart failure may contribute to liver enlargement. Scarring of the liver can raise pressure in the vein leading to the liver, which may enlarge the spleen.

Less common causes include blood disorders, immune conditions, cysts, noncancerous growths, and cancer. Certain inherited conditions cause substances to collect inside organs. Kidneys may enlarge because of cysts, urine-flow blockage, or inflammation.

Urgent causes are less common. A blow to an enlarged spleen can cause tearing and internal bleeding. A severe infection or a sudden blockage in an organ's blood supply may also cause rapid pain and illness.

When to seek medical care

Arrange a routine appointment if you have persistent upper-abdominal fullness, unexplained swelling, early fullness with meals, or a lump beneath your ribs. You should also follow up if an enlarged organ was found on a scan, even if you feel well.

Seek same-day medical advice for new or worsening abdominal pain with fever, repeated vomiting, yellow skin or eyes, dark urine, or marked abdominal swelling. These features may need prompt assessment but do not always mean there is a serious cause.

Get emergency help for:

If you know your spleen is enlarged, avoid contact sports and activities that could cause a blow to your abdomen until a clinician says they are safe.

- Sudden or severe abdominal pain
- Upper-left abdominal pain after a fall, collision, or blow
- Pain with fainting, severe dizziness, weakness, or clammy skin
- Severe breathing difficulty or rapidly worsening swelling

How the cause is found

Your clinician will ask where you feel discomfort, when it began, and whether you have fever, weight changes, early fullness, or digestive symptoms. They may ask about alcohol, medicines, supplements, infections, travel, family history, and previous liver, heart, blood, or kidney conditions.

During an examination, they may gently feel and tap your abdomen. It is not always possible to judge organ size accurately this way, especially when an enlargement is mild.

An ultrasound is often used to confirm which organ is enlarged and assess its appearance. CT or MRI scans can provide more detail when needed.

Blood tests may check:

Further testing depends on what these first results show. Some people need heart tests, blood-flow imaging, genetic testing, or assessment by a specialist. A biopsy, which removes a small tissue sample, is needed only in selected cases.

- Liver and kidney function
- Blood cell counts
- Signs of inflammation or infection

- Clotting and bile flow
- Tests for specific infections or immune conditions

Treatment and self-care

Treatment is directed at the cause rather than organ size alone. An infection may need monitoring or targeted treatment. Liver or heart conditions may require medicines and changes that reduce strain on the affected organ. A blockage, cyst, blood disorder, or growth may need specialist care.

Do not try to press, massage, or repeatedly check an enlarged area. This can increase discomfort and will not show whether the organ is improving.

Until you have been assessed, avoid starting herbal products or high-dose supplements marketed for "liver cleansing." Some can harm the liver or interact with medicines. If liver enlargement is suspected, ask about alcohol use and whether any prescription or nonprescription medicines need review. Do not stop prescribed medicine without advice.

If eating makes you feel very full, smaller meals may be more comfortable. This does not replace assessment of the cause.

What to expect

The outlook depends on the organ involved and why it became enlarged. Enlargement caused by a short-term infection may settle as the infection clears. Changes linked to long-term liver, heart, kidney, or blood conditions may persist and require monitoring.

Symptoms and organ size do not always improve at the same rate. Follow-up may include repeat blood tests, an examination, or another scan. Seek earlier review if pain, swelling, fever, jaundice, or your general condition worsens.

Common questions

Is visceromegaly the same as a swollen abdomen?

No. A swollen abdomen can result from gas, constipation, pregnancy, body-fat changes, fluid buildup, or a mass. Imaging may be needed to tell whether an organ is enlarged.

Can an enlarged organ return to normal size?

Sometimes. It may shrink when an infection, inflammation, congestion, or another reversible cause improves. Some long-term conditions leave lasting enlargement.

Can I feel an enlarged liver or spleen myself?

You may notice pressure or a firm area, but self-examination cannot reliably identify the organ or its size. Pressing deeply can be uncomfortable and may be unsafe if the spleen is enlarged.

Does visceromegaly mean cancer?

No. Cancer is one possible cause, but infections, fatty liver disease, blood-flow problems, and other noncancerous conditions can also enlarge organs. Testing is needed to identify the cause.

Sources

- Aldulaimi S, Mendez AM. Splenomegaly: Diagnosis and Management in Adults. American Family Physician . 2021;104(3):271-278.
- National Institute of Diabetes and Digestive and Kidney Diseases. Metabolic Dysfunction-Associated Steatotic Liver Disease.
- NHS. Spleen problems and spleen removal.
- Merck Manual Professional Edition. Splenomegaly.
- Rinella ME, Neuschwander-Tetri BA, Siddiqui MS, et al. AASLD practice guidance on the clinical assessment and management of nonalcoholic fatty liver disease. Hepatology . 2023;77(5):1797-1835.