TEST

Anti-smooth muscle antibody

A blood test that looks for antibodies associated with autoimmune hepatitis and some other conditions

WHAT IT MEASURES
Antibodies that bind smooth muscle structures, especially actin

SAMPLE
Blood

FASTING NEEDED?
No

HOW LONG IT TAKES
Usually less than 10 minutes

RESULTS IN
Usually several days; timing varies by laboratory

USED FOR
Supporting diagnosis of autoimmune hepatitis and related liver disease

MAIN LIMITATION
Positive results occur in several conditions, not only autoimmune hepatitis

Overview

An anti-smooth muscle antibody test looks for antibodies that react with proteins found in smooth muscle and other tissues. Antibodies are proteins made by the immune system. Smooth muscle works automatically in places such as the bowel and blood vessels.

Many anti-smooth muscle antibodies react with actin, a protein found throughout the body. Laboratories may call these smooth muscle antibodies, or SMA. Some use a more specific test for F-actin antibodies.

The test is mainly used when autoimmune hepatitis is suspected. Autoimmune hepatitis is long-term liver inflammation caused by the immune system attacking liver cells. A positive result can support this diagnosis, but it cannot confirm it by itself.

Some people with autoimmune hepatitis have a negative result. Positive results can also occur with other liver diseases, infections and autoimmune conditions. Your result therefore needs to be considered alongside liver blood tests, immunoglobulin G levels, other antibody tests and sometimes a liver biopsy.

Why the test is ordered

The test may be ordered when blood tests suggest liver inflammation without a clear cause. These tests may include alanine aminotransferase and aspartate aminotransferase, often shortened to ALT and AST.

It may also be requested if you have symptoms that could come from liver inflammation, such as persistent tiredness, nausea, itching, yellow skin or eyes, or discomfort under the right ribs. These symptoms have many possible causes, so the antibody test is only one part of the assessment.

Anti-smooth muscle antibody testing is often combined with:

The result is not generally used as a stand-alone screening test in people without symptoms or abnormal liver tests.

- Antinuclear antibody testing, also called ANA
- Immunoglobulin G, or IgG, measurement
- Tests for viral hepatitis and other causes of liver injury
- Liver imaging or biopsy when further evidence is needed

Who should have the test

You may be offered this test if your medical history, symptoms or liver test results raise the possibility of autoimmune hepatitis. It can also help investigate unexplained liver inflammation when more common causes have not provided an answer.

The test may be useful in adults and children, although antibody levels are interpreted differently across age groups. Testing may also be appropriate if you have another autoimmune condition and unexplained abnormal liver results.

A family history of autoimmune disease does not automatically mean that you need this test. The decision is usually based on your own results and clinical circumstances.

How to prepare

You do not need to fast for the anti-smooth muscle antibody test. You can usually eat, drink water and have coffee as normal.

Tell the healthcare professional about prescription medicines, over-the-counter products and

supplements. Some medicines can cause autoimmune-like liver inflammation, while immune-suppressing treatment may reduce antibody levels. Do not stop a medicine unless the clinician responsible for it advises you to do so.

If other blood tests are being collected at the same appointment, follow any preparation instructions for those tests. For example, a separate glucose or cholesterol test may require fasting.

What happens during the test

A healthcare professional places a band around your upper arm and cleans an area of skin. A small needle is inserted into a vein, usually inside your elbow, and blood is collected into a tube.

The blood draw normally takes less than 10 minutes. You may feel a brief scratch or stinging sensation. A dressing is placed over the site afterward, and the sample is sent to a laboratory.

Laboratories may examine the sample using indirect immunofluorescence, which shows whether antibodies attach to prepared cells or tissues. Another method measures antibodies against F-actin. The method used affects how the result is reported.

Understanding your results

A result may be reported as negative, positive, or as a titre. A titre shows how far the blood sample can be diluted while the antibody remains detectable. For example, a higher second number generally represents a stronger antibody reaction.

Other laboratories report F-actin antibodies in measurement units rather than as a titre. These results should be compared only with the reference information supplied by that laboratory.

A positive result means anti-smooth muscle antibodies were detected. It does not mean that you definitely have autoimmune hepatitis. The importance of the result increases when it occurs with compatible liver test abnormalities, raised IgG and other clinical evidence.

A weak positive result may be less specific. It can occur temporarily or alongside other liver, infectious or autoimmune conditions. A stronger result may provide more support for autoimmune hepatitis, but it is still not proof on its own.

A negative result makes antibody-positive autoimmune hepatitis less likely, but does not completely rule it out. Some people with autoimmune hepatitis do not have detectable smooth muscle antibodies or have different antibodies.

Reference ranges and interpretation vary by laboratory, testing method, age, sex, pregnancy

status, medicines and clinical context. Your report's own reference information is more useful than a range found online.

What affects the results

Anti-smooth muscle antibody results have relatively few day-to-day influences. Food, coffee, ordinary activity and the time of day do not usually change the result meaningfully.

Important influences include:

Kidney function and recent exercise do not usually change the antibody itself. However, strenuous exercise can affect liver enzymes collected at the same time, which may complicate the overall picture. Tell the healthcare team about pregnancy, recent transfusions and recent intravenous immunoglobulin so these can be considered.

- Testing method: Immunofluorescence and F-actin tests do not identify exactly the same antibodies. Results from different laboratories may not be directly comparable.
- Other illnesses: Viral hepatitis, some infections, other autoimmune diseases and several liver conditions can produce a positive result without autoimmune hepatitis.
- Medicines: Drugs including nitrofurantoin and minocycline can sometimes cause autoimmune-like liver injury with positive antibodies. Corticosteroids and other immune-suppressing medicines may reduce detectable antibody levels.
- Antibody-containing treatments: Recent intravenous immunoglobulin can temporarily introduce antibodies from donors and complicate interpretation.
- Sample quality: Severe breakdown of blood cells, cloudiness from fats or other collection problems may interfere with some laboratory methods.

Accuracy and limitations

This test can miss autoimmune hepatitis. A person may have the condition without smooth muscle antibodies, particularly if another antibody is present or the antibody level is below the method's detection limit.

The test can also be positive when autoimmune hepatitis is not present. This is more likely with a weak result or when liver enzymes and IgG are normal. Other liver diseases, infections and autoimmune conditions can produce similar antibodies.

F-actin testing may be more specific for autoimmune hepatitis than a broad smooth muscle antibody screen in some settings, but it can still give positive results for other reasons. Neither

version is accurate enough to be used alone.

When autoimmune hepatitis remains possible, tests such as ANA, anti-soluble liver antigen antibodies, IgG and viral hepatitis testing may provide more information. A liver biopsy can show the pattern and severity of inflammation and may be the better test when blood results do not give a clear answer.

Risks and complications

The risks are those of an ordinary blood draw. You may have a small bruise, brief soreness or feel faint; serious complications are uncommon.

What happens next

A negative result with otherwise reassuring liver tests may need no action beyond routine follow-up. If liver tests remain abnormal, further investigation may still be needed because a negative antibody result does not exclude every cause of liver inflammation.

After a positive result, the usual next step is to review it with your symptoms, medicines and other blood tests. Further testing may include IgG, ANA, viral hepatitis tests, liver imaging or a liver biopsy. Repeating the antibody test may help if the result is weak, unexpected or inconsistent with the rest of the assessment.

Anti-smooth muscle antibody levels are not usually used alone to monitor treatment. Liver enzymes, bilirubin, IgG, symptoms and sometimes repeat biopsy findings provide more useful information.

Arrange routine follow-up for an unexpected positive result or persistent abnormal liver tests. Seek urgent medical help for rapidly worsening yellowing, repeated vomiting, a swollen abdomen, or unusual bleeding. Confusion, severe drowsiness, vomiting blood or passing black, tar-like stool requires emergency assessment.

Common questions

Can I drink coffee before the test?

Yes. Coffee does not usually affect anti-smooth muscle antibodies. Check your instructions if other fasting blood tests are being taken at the same appointment.

Does the time of day matter?

No. Antibody levels are not known to have a clinically important daily pattern, so the sample can usually be collected at any convenient time.

Can I exercise beforehand?

Ordinary activity is fine. Strenuous exercise does not usually alter this antibody, but it can temporarily affect some liver and muscle blood tests taken alongside it.

Should I stop my medicines?

No, unless the clinician responsible for the medicine tells you to. Bring an up-to-date list because some medicines can cause autoimmune-like liver injury or reduce antibody levels.

Does a positive result mean I have autoimmune hepatitis?

No. It supports the diagnosis only when it fits with liver tests, IgG, symptoms and other findings. Positive results also occur in other conditions.

Is the test safe during pregnancy?

Yes. It is a standard blood draw. Pregnancy does not prevent testing, but it forms part of the clinical context used to interpret your result.

Sources

- National Institute of Diabetes and Digestive and Kidney Diseases. Diagnosis of Autoimmune Hepatitis
- Mack CL, Adams D, Assis DN, et al. Diagnosis and Management of Autoimmune Hepatitis in Adults and Children: 2019 Practice Guidance and Guidelines From the American Association for the Study of Liver Diseases . Hepatology . 2020;72(2):671-722.
- European Association for the Study of the Liver. EASL Clinical Practice Guidelines: Autoimmune hepatitis . Journal of Hepatology . 2015;63(4):971-1004.
- Vergani D, Alvarez F, Bianchi FB, et al. Liver autoimmune serology: a consensus statement from the committee for autoimmune serology of the International Autoimmune Hepatitis Group . Journal of Hepatology . 2004;41(4):677-683.
- ARUP Consult. Autoimmune Hepatitis
- Mayo Clinic Laboratories. Smooth Muscle Antibody Screen, Serum