TEST

Cryoglobulins

A cryoglobulin test checks blood for proteins that clump in the cold and dissolve when warmed.

WHAT IT MEASURES
Blood proteins that precipitate when cold and redissolve when warmed

SAMPLE
Blood

FASTING NEEDED?
Sometimes, if your laboratory requests an overnight fast

HOW LONG IT TAKES
Usually under 10 minutes

RESULTS IN
Several days to one week; varies by laboratory

USED FOR
Helping diagnose conditions linked to cryoglobulins

MAIN LIMITATION
Cooling before serum separation can cause a false-negative result

Overview

Cryoglobulins are antibodies or antibody-containing proteins that separate from blood when they become cold. This is called precipitation. They usually dissolve again when warmed.

The cryoglobulin test looks for these proteins in the liquid part of blood, called serum. The sample must be collected and handled under carefully controlled temperatures. If it cools too soon, cryoglobulins may be lost before testing.

Finding cryoglobulins does not diagnose a specific disease. Some are linked to infections, autoimmune diseases, or disorders involving antibody-producing blood cells. Small amounts can also be found without a clear related illness.

Laboratories may report whether cryoglobulins were found, estimate how much is present, and identify the type of antibody involved.

Why the test is ordered

The test is usually ordered when symptoms or other results suggest that cold-sensitive proteins may be affecting blood vessels, nerves, skin, or kidneys.

Reasons for testing may include:

The test may also help classify a previously detected cryoglobulin or follow a known cryoglobulin-related condition. Cryoglobulin levels do not always match symptom severity, so the result is not usually used by itself to judge improvement or worsening.

- A raised purple or red rash, especially on the legs
- Skin sores or fingers and toes that become painful in the cold
- Unexplained joint pain, weakness, numbness, or tingling
- Blood or protein found in urine
- Reduced kidney function without a clear explanation
- Signs of inflammation in small blood vessels, called vasculitis
- Investigation of hepatitis C, certain autoimmune diseases, or blood cell disorders

Who should have the test

Cryoglobulin testing is not a routine screening test for people without symptoms or relevant medical findings.

It may be suitable if your healthcare professional suspects cryoglobulinemia, meaning cryoglobulins are present in the blood, or cryoglobulinemic vasculitis, meaning these proteins are associated with blood vessel inflammation.

Not everyone with hepatitis, an autoimmune condition, kidney disease, cold fingers, or a skin rash needs this test. The decision is based on your symptoms, examination, medical history, and results from other blood or urine tests.

How to prepare

Follow the instructions from the laboratory collecting your sample. Some laboratories prefer an overnight fast because fat particles can make serum cloudy and harder to assess. Other laboratories do not require fasting.

If you are told to fast, drink only plain water during the fasting period. Do not stop prescribed medicines unless the healthcare professional who prescribed them tells you to.

Tell the person arranging the test about medicines that affect the immune system, recent

plasma exchange, treatment for a blood cell disorder, or a recent blood transfusion. These may influence interpretation.

You do not usually need to warm yourself before the test. Keeping the specimen warm after collection is the laboratory's responsibility.

What happens during the test

A healthcare worker takes blood from a vein, usually inside your elbow. A tourniquet is placed around your upper arm, the skin is cleaned, and a needle is inserted into the vein.

Cryoglobulin samples require special handling. The tube and collection equipment may be pre-warmed. After collection, the blood is kept close to body temperature while it clots and until the serum is separated from the blood cells.

In the laboratory, the serum is cooled for several days and checked for material that settles or forms a gel. Any precipitate is warmed to confirm that it dissolves again. Further testing, often immunofixation, can identify the antibodies involved.

Understanding your results

A result may be reported as negative, positive, or detected below a laboratory's measurement threshold.

A negative result means cryoglobulins were not found in that sample using that laboratory's method. It does not completely rule them out, especially if the sample cooled before separation or the amount was very low.

A positive result confirms that cold-precipitating proteins were detected. It does not show by itself whether they are causing symptoms or why they are present.

Further testing may classify them into three broad groups:

The amount may be reported as a cryocrit, which estimates the percentage of serum occupied by the precipitate. Cryocrit measurement is imprecise, and a larger amount does not always mean more severe illness.

There is no single universal reference range. Methods and reporting thresholds differ between laboratories. Interpretation also reflects age, sex, pregnancy status, medicines, kidney function, other test results, and the clinical context, even when these factors do not directly change the cryoglobulin measurement.

- Type I: One abnormal antibody, called a monoclonal immunoglobulin
- Type II: A mixture containing one monoclonal and other polyclonal antibodies
- Type III: A mixture of polyclonal antibodies produced by several groups of immune cells

What affects the results

Specimen handling is the most important influence. Cryoglobulins can attach to blood cells or precipitate if the blood cools before the serum is separated. This can produce a negative result even when cryoglobulins are present.

Other factors include:

Exercise and time of day are not known to have major direct effects. Age, pregnancy, and kidney function generally affect the clinical meaning more than the laboratory measurement itself.

- Too little incubation time: Some cryoglobulins take several days to appear
- Cloudy or fatty serum: A recent high-fat meal can make visual assessment difficult
- Incomplete clotting: Fibrin strands may be mistaken for cryoprecipitate
- Wrong sample type: Cryoglobulin testing requires serum collected and processed correctly
- Immune-suppressing treatment: Steroids like prednisone or hydrocortisone , antibody treatments, chemotherapy, or plasma exchange may lower detectable levels
- Recent transfusion: Donor antibodies or dilution of your own antibodies may complicate interpretation

Accuracy and limitations

There is no single reliable figure for how often cryoglobulin testing is right or wrong. Collection procedures, cooling time, detection methods, and reporting thresholds are not fully standardised between laboratories.

The main problem is a missed result. A sample that cools too soon can test negative because the proteins have already separated from the serum or attached to blood cells. When suspicion remains high, a carefully collected repeat sample may be more useful than accepting one negative result.

A reported positive can also be misleading. A small amount may be genuinely present but

unrelated to your symptoms. Fibrin, fat particles, or other material may resemble a cryoprecipitate. Confirming that the material dissolves when warmed helps reduce this problem.

Immunofixation is better for identifying the antibody type. Complement tests, rheumatoid factor, hepatitis testing, kidney tests, and urine testing help assess possible causes and effects. If blood vessel or kidney inflammation must be confirmed, a skin or kidney biopsy may provide information that a cryoglobulin blood test cannot.

Risks and complications

A blood draw may cause brief discomfort, a small bruise, or minor bleeding where the needle entered. Some people feel light-headed or faint, but serious complications are uncommon.

What happens next

A negative result may need no further action if your symptoms and other results do not suggest a cryoglobulin-related problem. If concern remains, your healthcare professional may discuss repeating the test with strict warm handling.

After a positive result, follow-up commonly includes antibody typing and tests for kidney involvement, inflammation, complement proteins, hepatitis, autoimmune disease, or abnormal blood cells. The choice of tests is guided by your history rather than the cryoglobulin result alone.

Routine follow-up is appropriate for discussing stable symptoms and laboratory findings. Seek prompt medical advice for a new painful skin ulcer, rapidly increasing rash, dark urine, much less urine than usual, or worsening weakness or numbness.

Call emergency services for sudden one-sided weakness, difficulty speaking, severe breathing difficulty, chest pain, or another rapidly developing serious symptom. Do not wait for a cryoglobulin result in an emergency.

Common questions

Can I drink coffee before the test?

If fasting is not required, coffee is usually allowed. If your laboratory requests fasting, have only plain water unless its instructions say otherwise.

Does the time of day matter?

Usually not. Correct warm handling of the sample matters much more than the collection time.

Can I exercise beforehand?

Normal activity is unlikely to change the result directly. Avoid unusually strenuous exercise if it worsens your symptoms or if you have been given different instructions.

Why does the result take several days?

Some cryoglobulins precipitate slowly. Laboratories may keep the serum cold for up to several days before deciding that no precipitate is visible.

Can cold weather change my result?

Cold exposure can trigger symptoms in some people, but it is not needed for the test. The crucial issue is preventing the blood sample from cooling before serum separation.

Should I stop my medicines first?

No. Some treatments can lower cryoglobulin levels, but stopping them may be unsafe. Tell your healthcare professional and the laboratory what you take.

Sources

- Sargur R, White P, Egner W. "Cryoglobulin evaluation: best practice?" Annals of Clinical Biochemistry . 2010;47(1):8-16.
- Motyckova G, Murali M. "Laboratory testing for cryoglobulins." American Journal of Hematology . 2011;86(6):500-502.
- Shihabi ZK. "Cryoglobulins: an important but neglected clinical test." Annals of Clinical and Laboratory Science . 2006;36(4):395-408.
- Brouet JC, Clauvel JP, Danon F, Klein M, Seligmann M. "Biologic and clinical significance of cryoglobulins: a report of 86 cases." The American Journal of Medicine . 1974;57(5):775-788.
- ARUP Consult. "Cryoglobulinemia - Choose the Right Test." Laboratory guidance on testing, specimen handling, and interpretation.
- Mayo Clinic Laboratories. "Cryoglobulins, Serum." Test catalog guidance on specimen collection, processing, and reporting.