TEST

Esophageal ph monitoring

Esophageal pH monitoring records how often and how long stomach acid enters your esophagus.

WHAT IT MEASURES
Acid exposure inside the esophagus over one or more days

SAMPLE
None; measurements come from a temporary esophageal sensor

FASTING NEEDED?
Sometimes, before sensor placement or sedated endoscopy

HOW LONG IT TAKES
Placement 15-30 minutes; recording usually 24-96 hours

RESULTS IN
Several days; varies by laboratory and specialist review

USED FOR
Diagnosing reflux and assessing persistent symptoms during treatment

MAIN LIMITATION
Reflux varies daily, so one recording may be unrepresentative

Overview

Esophageal pH monitoring measures acidity inside your esophagus, the tube carrying food to your stomach. It is mainly used to investigate gastroesophageal reflux, when stomach contents travel upward.

A sensor records periods of acid exposure. You also record meals, sleep, symptoms, and sometimes medicines so these events can be compared with reflux.

A catheter-based test uses a thin tube passed through your nose into your esophagus, usually for about 24 hours. Some catheters also use impedance to detect liquid, gas, and weakly acidic or non-acid reflux.

A wireless test uses a small capsule attached to the esophageal lining, often during an upper endoscopy. It sends readings to a portable receiver for about 48 to 96 hours, then detaches and passes through your digestive system.

Why the test is ordered

Your healthcare professional may order monitoring when it is unclear whether reflux explains heartburn, regurgitation, chest discomfort, chronic cough, or throat symptoms.

It may also be used when symptoms continue despite treatment, an endoscopy is inconclusive, before an anti-reflux procedure, or to assess whether acid suppression is working. It is not usually needed for straightforward reflux symptoms.

Who should have the test

The test may help if you have persistent symptoms whose cause remains uncertain. Testing can be performed while taking acid-reducing medicine or after temporarily stopping it. Testing without medicine looks for abnormal acid reflux; testing during treatment assesses ongoing symptoms in someone with established reflux disease.

Impedance-pH monitoring may be preferred when non-acid reflux is a concern. Longer wireless monitoring can help when symptoms are unpredictable.

Tell the testing team if you are pregnant, have difficulty swallowing, have had esophageal or stomach surgery, take blood-thinning medicines, or have an implanted medical device. These factors may affect the test.

How to prepare

Follow your testing unit's instructions. Preparation depends on the monitor, whether endoscopy is planned, and whether testing is being done on or off treatment.

You may be told to stop proton pump inhibitors like omeprazole or lansoprazole , H2 blockers, antacids, or other digestive medicines. Do not stop prescribed medicine unless the testing team tells you when and how. Taking or stopping these medicines changes what the test measures.

You may need to avoid food and drink for several hours, especially before sedation or endoscopy. Tell the team about all medicines and supplements, and ask whether someone must take you home after sedation.

Wear comfortable clothing. The receiver is generally carried on a belt or shoulder strap.

What happens during the test

For catheter monitoring, a clinician may numb your nose before guiding a thin tube through one nostril into your esophagus. You may be asked to swallow during placement. The catheter is secured to your face and connected to a recorder. Placement may briefly cause watering eyes, gagging, coughing, or discomfort.

For wireless monitoring, a capsule is attached to your esophagus, commonly during a sedated upper endoscopy. It transmits readings to a receiver you carry.

During monitoring, press the recorder buttons or complete a diary when you eat, lie down, take medicine, or have symptoms. Accurate entries help the laboratory compare symptoms with reflux.

Unless instructed otherwise, follow your usual routine and eat your typical foods. The catheter is removed at the clinic. A wireless capsule detaches by itself.

Understanding your results

The report describes how long your esophagus was exposed to acid. Acid exposure is conventionally recorded when pH falls below 4, although interpretation involves more than one number.

Results may include:

Symptom association does not prove reflux caused every episode. Results must be interpreted with your symptoms, medicines, endoscopy findings, and other tests. Reference ranges vary by laboratory and monitoring method.

- Acid exposure while upright, lying down, and overall
- The number and duration of reflux events
- Whether symptoms occurred near reflux events
- Non-acid reflux if impedance was used
- Whether the recording was technically reliable

What affects the results

Acid-reducing medicines can substantially change the findings. Taking medicine during a test intended to be performed off treatment-or stopping it during a test intended to assess treatment-may answer the wrong question.

Eating less than usual, avoiding trigger foods, changing sleep position, or limiting activity can

make the study unrepresentative. Acidic foods may temporarily lower pH, so record meal times accurately.

Other influences include sensor movement, early capsule detachment, lost signals, vomiting, equipment disconnection, missing diary entries, or monitoring on an unusually mild or severe day. Reflux varies daily, and even longer monitoring cannot capture every pattern.

Accuracy and limitations

Monitoring can miss reflux that does not occur during the recording, especially when symptoms are infrequent. Standard pH testing may also miss weakly acidic or non-acid reflux; impedance-pH monitoring is better for detecting these events.

Wireless monitoring records for longer and avoids a nasal tube but generally measures acidity rather than movement of non-acid material. A catheter can include impedance, although wearing it may alter eating, sleep, or activity.

Endoscopy is better for finding inflammation, narrowing, ulcers, or structural problems. Esophageal manometry assesses swallowing muscle function. These tests answer different questions and may be used together.

Risks and complications

A nasal catheter commonly causes temporary throat or nasal discomfort. Minor nosebleeding, gagging, coughing, or feeling faint can occur during placement.

A wireless capsule may cause temporary chest awareness or discomfort, especially when swallowing. Endoscopy and sedation have separate risks that should be explained beforehand.

Contact the testing unit promptly if the recorder stops working, the catheter comes out, you cannot swallow normally, or chest discomfort persists or worsens.

Seek urgent medical care for severe chest pain, difficulty breathing, vomiting blood, black stools, or a severe reaction after sedation. These are not routine effects.

A wireless capsule may affect when you can have an MRI. Tell imaging staff about the capsule and follow the device-specific instructions.

What happens next

The laboratory downloads and reviews the recording after you return the equipment. Your

healthcare professional will explain whether acid exposure was within the expected range and whether reflux matched your symptoms.

A normal result makes significant acid reflux less likely but does not identify another cause. Follow-up may include continuing or changing treatment, reviewing how you take medicine, arranging endoscopy or manometry, or investigating other causes. Do not change prescribed treatment based only on the numerical report.

Common questions

Can I drink coffee before the test?

You may need to fast before placement, so coffee might not be allowed that morning. During monitoring, you can often follow your usual diet. Follow your testing unit's instructions.

Does the time of day matter?

The monitor records continuously. Mark when you lie down and get up so daytime and nighttime reflux can be compared.

Can I exercise while wearing the monitor?

Usual light activity is often encouraged. Avoid swimming, showering, contact sports, or anything that could wet, pull out, or damage the equipment.

Can I eat normally during the test?

Usually, yes. Record the beginning and end of meals because eating can change pH readings.

Should I take my reflux medicine?

Only as instructed. Some tests require continued treatment; others require medicines to be stopped. Contact the testing unit if instructions are unclear.

Can I shower or sleep normally?

You can usually sleep in your normal position, but record when you lie down. External recorders generally must stay dry, so showering, bathing, and swimming are restricted.

Sources

- Katz PO, Dunbar KB, Schnoll-Sussman FH, et al. ACG Clinical Guideline for the Diagnosis and Management of Gastroesophageal Reflux Disease . American Journal of Gastroenterology . 2022;117(1):27-56.

- Yadlapati R, Gyawali CP, Pandolfino JE. AGA Clinical Practice Update on the Diagnosis and Management of Extraesophageal Gastroesophageal Reflux Disease . Gastroenterology . 2023;164(4):628-639.
- Gyawali CP, Yadlapati R, Fass R, et al. Updates to the Modern Diagnosis of GERD: Lyon Consensus 2.0 . Gut . 2024;73(2):361-371.
- Trudgill NJ, Sifrim D, Sweis R, et al. British Society of Gastroenterology Guidelines for Oesophageal Manometry and Oesophageal Reflux Monitoring . Gut . 2019;68(10):1731-1750.
- National Institute of Diabetes and Digestive and Kidney Diseases. Diagnosis of GER and GERD . Reviewed July 2020. Accessed August 9, 2026.